

a) Given: $\phi_1(x,y)$, $\phi_2(x,y)$, $p_1(x,y)$, $p_2(x,y)$
 $\nabla^2 \phi_1 = 0$, $\nabla^2 \phi_2 = 0$ (satisfy mass conservation)

Define: $\phi_3 = \phi_1 + \phi_2$

Determine p_3

$$\Rightarrow p_3 = p_0 - \frac{1}{2}\rho |\nabla \phi_3|^2 = p_0 - \frac{1}{2}\rho (\nabla \phi_3 \cdot \nabla \phi_3)$$

$$= p_0 - \frac{1}{2}\rho [(\nabla \phi_1 + \nabla \phi_2) \cdot (\nabla \phi_1 + \nabla \phi_2)]$$

$$p_3 = p_0 - \frac{1}{2}\rho [|\nabla \phi_1|^2 + |\nabla \phi_2|^2 + 2\nabla \phi_1 \cdot \nabla \phi_2]$$

Note: $p_3 \neq p_1 + p_2$!

b) Given $\phi_4 = \partial \phi_1 / \partial x$

Is it physically realizable?

Test: $\nabla^2 \phi_4 \stackrel{?}{=} 0$

$$\nabla^2 (\partial \phi_1 / \partial x) \stackrel{?}{=} 0$$

$$\frac{\partial}{\partial x} (\nabla^2 \phi_1) \stackrel{?}{=} 0$$

$\frac{\partial}{\partial x}()$ and $\nabla^2()$ commute.


Since $\nabla^2 \phi_1 = 0$ as given

Example: $\phi_1 = \frac{\Lambda}{2\pi} \ln \sqrt{x^2 + y^2} = \frac{\Lambda}{2\pi} \ln r$ source

$$\phi_4 = \frac{\partial \phi_1}{\partial x} = \frac{\Lambda}{2\pi} \frac{x}{x^2 + y^2} = \frac{\Lambda}{2\pi} \frac{\cos \theta}{r}$$
 doublet!

$$u_1 = \frac{y}{x^2+y^2} \quad v_1 = \frac{-x}{x^2+y^2}$$

" $\frac{1}{r}$ vortex"



$$u_2 = V_\infty \quad v_2 = 0$$

uniform flow



$$u_3 = u_1 + u_2 = \frac{y}{x^2+y^2} + V_\infty \quad v_3 = v_1 + v_2 = \frac{-x}{x^2+y^2}$$


$$P_3 = P_0 - \frac{1}{2} \rho (u_3^2 + v_3^2) = P_0 - \frac{1}{2} \rho \left[\frac{y^2}{(x^2+y^2)^2} + \frac{2yV_\infty}{x^2+y^2} + V_\infty^2 + \frac{x^2}{(x^2+y^2)^2} \right]$$

$$P_3 = P_0 - \frac{1}{2} \rho \left[\frac{1+2yV_\infty}{x^2+y^2} + V_\infty^2 \right]$$

Maximum pressure is where $u_3^2 + v_3^2$ is minimum.

Note that on y-axis where $x=0$, we have $v_3=0$.
Also, at $y = -1/V_\infty$ we also have $u_3=0$

$$\Rightarrow \text{max } P_3 \text{ at } x, y = 0, -\frac{1}{V_\infty}$$



Alternative mathematical approach (hard way)

$$\text{set } \frac{\partial P_3}{\partial x} = 0 \text{ and } \frac{\partial P_3}{\partial y} = 0$$

$$\frac{\partial P_3}{\partial x} = -\frac{1}{2} \rho \frac{1+2yV_\infty}{(x^2+y^2)^2} (-2x) = 0 \Rightarrow (1+2yV_\infty)x = 0 \quad (1)$$

$$\frac{\partial P_3}{\partial y} = -\frac{1}{2} \rho \frac{1+2yV_\infty}{(x^2+y^2)^2} (-2y) - \frac{1}{2} \rho \frac{2V_\infty}{x^2+y^2} = 0 \Rightarrow (1+2yV_\infty)y - (x^2+y^2)V_\infty = 0$$

$$\text{or } (1+yV_\infty)y - x^2V_\infty = 0 \quad (2)$$

Two possibilities from equation (1)

a) $1+2yV_\infty = 0, x \neq 0 \rightarrow y = -\frac{1}{2V_\infty}$

Plug into equation (2) $\rightarrow \frac{1}{2} \cdot \left(-\frac{1}{2V_\infty}\right) - x^2V_\infty = 0 \rightarrow x^2 = \frac{-1}{4V_\infty}$ no real solution

b) $1+2yV_\infty \neq 0, \boxed{x=0}$



Plug into equation (2) $\rightarrow (1+yV_\infty)y = 0 \rightarrow y = 0$ Nope. Inconsistent with (1).

$\rightarrow 1+yV_\infty = 0 \rightarrow \boxed{y = -\frac{1}{V_\infty}}$ ✓

13-732
42-381
42-382
42-383
42-384
42-385
42-386
42-387
42-388
42-389
42-390
42-391
42-392
42-393
42-394
42-395
42-396
42-397
42-398
42-399
42-400
500 SHEETS, FILLER, 5 SQUARE
50 SHEETS, EYE-EASE, 5 SQUARE
100 SHEETS, EYE-EASE, 5 SQUARE
200 SHEETS, EYE-EASE, 5 SQUARE
100 RECYCLED WHITE, 5 SQUARE
200 RECYCLED WHITE, 5 SQUARE
Made in U.S.A.



M17



Mohr's circles read

a) 60° rosette - plots as 120° on Mohr's circle

ϵ_a aligned with x_1 , \therefore reads $\epsilon_{11} = -200 \mu \epsilon \Leftarrow$

ϵ_b 60° ~~clockwise~~ counter clockwise

$$\text{angle } \phi = 120^\circ - 106.85^\circ = 13.15^\circ$$

$$\therefore \epsilon_b = \frac{100}{c} + \frac{361}{R} \sin 13.5^\circ = 182.1 \mu \epsilon \Leftarrow$$

$$\text{angle } \psi = 180 - 120 - 16.85 = 43.2^\circ$$

$$\epsilon_c = 100 + 361 \cos 43.2^\circ = 363.4 \mu \epsilon \Leftarrow$$

$$b) \quad \Sigma_{mn} = \begin{pmatrix} -200 & -200 & 0 \\ -200 & +400 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Eq eigenvalues solution to $\Delta = |M - \lambda I|$

$$(-200 - \lambda)(+400 - \lambda) - (-200)^2 = 0$$

$$\lambda^2 - 200\lambda - 120000 = 0$$

$$\lambda = \frac{+200 \pm \sqrt{200^2 + 4 \times 120000}}{2}$$

$$= 100 \pm 361$$

$$= -261 \mu\epsilon \text{ and } +461 \mu\epsilon !! \leftarrow$$

$$c) \text{ for } \Sigma_{33} = 300 \mu\epsilon \text{ and } \Sigma_{23} = \Sigma_{13} = 0$$

Σ_{33} is a principal strain. (no associated shear)

$$\therefore \Sigma_{\text{I}} = 461 \mu\epsilon \text{ + } \Sigma_{\text{II}} = -261 \mu\epsilon \text{ as before}$$

$$\Sigma_{\text{III}} = 300 \mu\epsilon. \leftarrow$$

M18 i) Young's modulus is controlled by bonding and crystal structure (atomic packing)

ii) The glass transition temperature is the temperature at which the polymer changes from being an elastic solid to a viscoelastic one (at higher temps) which is due to the breaking (melting) of the van-der-Waals bonds between the polymer molecules

iii) Metals have higher densities because they have higher atomic masses and close packed crystal structures

iv) Interatomic bond energy $U(r)$ ^{separation}

$$\text{Interatomic force} = \frac{dU(r)}{dr} \Leftarrow$$