





Massachusetts Institute of Technology  
Organic Chemistry 5.512


March 9, 2005  
Prof. Rick L. Danheiser

Problem Set 2  
Strategies for Synthesis of Acyclic Molecules Based on  
Desymmetrization, Chirality Transfer, the "Chiron Approach",  
and "Ring Template" Strategies

- I. Design a highly stereoselective synthesis of the following target molecules beginning with commercially available materials and employing one of the strategies listed above. For the compounds indicated, a route to the target molecule in racemic form will be sufficient. Be sure to explicitly identify all reagents necessary for each transformation. Enantiomerically enriched reagents may be used if they are commercially available; however, each stereogenic center in the target molecule must be generated in your synthetic route. In other words, the stereogenic carbons in the chiral reagents you employ cannot be directly incorporated in the final product.



- II. In 1944, Paranjape and coworkers in India announced a "total asymmetric synthesis" of the anthelmintic sesquiterpene santonin (see attached article). They reported that treatment of racemic 2-formylcyclohexanone with sodium ethoxide (to form the sodium enolate) followed by alkylation with methyl iodide (in either benzene, toluene, or ethanol as solvent) afforded 2-methyl-2-formylcyclohexanone as a liquid with an optical rotation of  $-26.2^\circ$ ! How do you account for these results?



difficulty. The  
ited to  $3 \times 10^8$   
um is combined  
leads directly to  
have been much

difficulty of all  
Jeans's theory  
er modification,  
ould need to be  
ities of thermal  
xpansion due to  
ave offered some  
not particularly  
does not appear

important in the  
have some idea  
olids at tempera-  
It seems clear  
asteroids could  
ous state under  
emperature was low  
to fall below the  
take place any-  
stended star, the  
of gas. In the  
tellites we have  
dium of the order  
nging from  $100^\circ$   
as above that of  
considered, there  
could grow no  
t, their gravita-  
ce more than an  
quired could be  
to find them in

OLD JEFFREYS.

lfvén<sup>1</sup> appears to  
the problem of  
resent time the  
is determined by  
n endowment of

which Jupiter is  
initially endowed  
any effect pro-  
be regarded as  
irrelevant to the  
is suggested that  
ired during the  
ssary to describe  
h the transfer of

nsisted originally  
the form of gas,  
l that the stars,  
lved by a process  
then an adequate  
ust explain three  
rtum of the solar  
gular momentum  
cloud; (2) why

the angular momentum (per unit mass) of the sun, referred to the centre of the system, is so much less than that of the planets; and (3) why the angular momentum (per unit mass) of a planet, such as Jupiter, referred to the centre of the planetary system, is so much less than that of its satellites.

K. E. EDGEWORTH.


Cherbury, Booterstown,  
Co. Dublin. Dec. 24.

<sup>1</sup> NATURE, 152, 721 (1943).


## A Case of Total Asymmetric Synthesis

We have recently announced a successful synthesis of santonin (I)<sup>1</sup>. We believed then that it was racemic. We have since found it to be optically active. The synthetic specimen has a rotation of  $[\alpha]_D = -150$  add m.p.  $171^\circ$ . Sodium santoninate prepared from this sample was fractionally precipitated as the strychnine salt. This salt on decomposition gave santonin with  $[\alpha]_D = -172$  identical with the rotation of natural santonin. The filtrates gave another product with identical m.p. ( $171^\circ$ ) and rotation  $[\alpha]_D = -108$ , which was not changed on further fractional precipitation by strychnine or quinine hydrochloride, but in no case was a dextro-rotatory product obtained. So far as we are aware, this is the first total asymmetric synthesis, apart from asymmetric synthesis carried out in the presence of polarized light, etc.

We have also prepared compounds II, III, IV and V. It has been found that asymmetric synthesis occurred only in those compounds (I, IV and V) which had an angular methyl group. The others were racemic and could be resolved into *d*- and *l*-forms through their strychnine salts.



We have also determined the stage at which the asymmetric synthesis occurs by carrying out the synthesis of V by the following reactions:



VII is an unstable liquid and cannot be purified by distillation, and also its solid derivative could not be prepared. However, the crude substance had an optical rotation  $[\alpha]_D = -26.22$ . The corresponding methylated formyl derivative used in the synthesis of santonin could not be examined for optical activity as it was highly coloured. Methylation of the formyl

cyclohexanone or the corresponding derivative appears to be the stage at which the asymmetric synthesis occurs.

All rotations have been determined in chloroform solution at  $28^\circ\text{C}$ .

Further work is in progress.

K. D. PARANJAPÉ.  
N. L. PHALNİKAR.  
B. V. BHIDE.  
K. S. NARGUND.

Maharaja Pratapsinh Chemical Laboratory,  
Sir Parashurambhau College,  
Poona, 2.  
Nov. 13.

<sup>1</sup> Current Science, 12, 153 (1943).

## Aerial Disinfection

THE letter from workers at the National Institute of Medical Research<sup>1</sup> was of interest to us, as we also find that lactic acid is an effective bactericidal aerosol, when the test organism is *C. xerosis*, emulsified in sterile saliva.

Bechhold<sup>2</sup> long ago advocated the use of hydroxy and carboxylic acids as germicidal aerosols. We found cinnamic and benzoic acids (both constituents of Peru balsam<sup>3</sup>) to be strong aerial bactericides, but acids, such as citric, fumaric, maleic, malic and phthalic tested recently proved relatively ineffective. On the other hand, maleic and phthalic anhydrides were found to be more active than their corresponding acids. In a concentration of 4 mgm./m<sup>3</sup>. and a relative humidity of about 60 per cent, maleic anhydride generally sterilized the air of our test organism within five minutes; the durability of lethal effectiveness being good, a 15-minute old mist only allowing survival of some 5 per cent of the bacteria beyond the 5-minute exposure time.

As regards the amount of germicide necessary to sterilize or nearly sterilize the air of our test organism within 5 minutes, it was found that of fifteen phenolic compounds examined, each required the vapour concentration to be of the order of 25 per cent saturation. Saturation at  $20^\circ\text{C}$ . was calculated from the vapour pressures deduced from the formula of Clausius and Clapeyron (which can be expressed as  $\log p = A + B/T$ , where *A* and *B* are constants). Other substances, for example, mercuric chloride, propylene and diethylene glycols, apparently require to be of an approximately similar concentration. Notable exceptions are iodine and maleic anhydride (below 1 per cent), suggesting that the mechanism of the lethal effect here is different from that of

the phenols. It will thus be understood that while we see no reason to alter our opinion<sup>4</sup> regarding the fundamental importance of the vapour pressure of phenols in relation to activity on air-borne bacteria, it appears that this particular characteristic is not a reflexion of the activity of all groups of substances. Again, we do not consider water solubility an essential