

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
Department of Physics

Physics 8.01

Fall 2003

EXAM 3
Friday, November 21, 2003

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

FAMILY (Last) NAME

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

GIVEN (First) NAME

--	--	--	--	--	--	--	--	--	--

Student ID Number

Your Recitation (check one)



Instructions:

1. **SHOW ALL WORK.** All work must be done in this booklet. Print your name on each sheet.
2. One 8 ½ x 11 sheet of notes allowed.
3. This is a closed book exam.
4. CALCULATORS, BOOKS, COMPUTERS and CELL PHONE are NOT ALLOWED.
5. Do all **FOUR (4)** problems.
6. Extra pages provided.
7. Exams will be collected 5 minutes before the hour.

Problem	Maximum	Score	Grader
1	15		
2	15		
3	15		
4	15		
TOTAL	60		


R17	TR 12:00		Maria Chan
R23	TR 11:00		Maria Chan
R12	TR 1:00		Min Chen
R13	TR 2:00		Min Chen
R14	TR 3:00		Min Chen
R01	MW 1:00		Bruno Coppi
R05	MW 2:00		Bruno Coppi
R09	MW 1:00		Qudsia Ejaz
R06	MW 2:00		Paul Joss
R07	MW 3:00		Paul Joss
R08	MW 4:00		Paul Joss
R02	MW 2:00		Young Lee
R03	MW 3:00		Young Lee
R15	TR 10:00		Hong Liu
R16	TR 11:00		Hong Liu
R21	TR 2:00		David Pritchard
R22	TR 3:00		Gunther Roland
R25	TR 1:00		Gunther Roland
R04	MW 1:00		Brian Ross
R19	TR 10:00		Brian Ross
R10	MW 2:00		Senthil Todadri
R11	MW 3:00		Senthil Todadri
R18	TR 9:00		Vladan Vuletic
R24	TR 12:00		Vladan Vuletic
R20	TR 11:00		Xiao-Gang Wen

Name: _____

Problem 1: Collision (15 Points)

Two masses M and $3M$ collide on a horizontal frictionless surface as shown. Before the collision the mass M has a velocity V_1 in the y -direction. The mass $3M$ has a velocity $\frac{5}{12}V_0$ making an angle θ to the x -axis as shown. Assume $\sin\theta = 3/5$ and $\cos\theta = 4/5$. After the collision the mass $3M$ is at rest. The mass M moves along the x -axis with the velocity V'_1 . Neglect gravity. Give all your answers to parts b), c), and d) in terms of M and V_0 . Be careful do not confuse your symbols.

- a) What are the x and y -components of the net linear momentum before the collision in terms of M , V_1 , V_0 and θ ?
- b) What is the speed V_1 of the mass M before the collision?
- c) What is the speed V'_1 of the mass M after the collision?
- d) What is the velocity of the center-of-mass?




Name: _____

Problem 2: Rotational Dynamics (15 points)

A spool of wire of mass M and radius R is unwound along a horizontal surface under a constant force \vec{F} . Assume the spool is a uniform solid cylinder that does not slip. The coefficient of static friction is μ_s . Assume that the radius of the spool does not decrease significantly while the spool is rolling. Give all your answers in terms of F , M , R , μ_s , g and L .

- a) State the moment of inertia, I , of the cylinder about its central axis.
- b) What is the force of friction, \vec{f} (magnitude AND direction) acting on the spool? Show the direction of f on the diagram.
- c) What is the acceleration of the center-of-mass?
- d) What is the angular acceleration?
- e) What is the total kinetic energy of the spool when it has rolled through a distance L ?




Name: _____

Problem 3: Rotational Collision (15 points)

A uniform cylindrical shell (hoop) sits on one of its flat sides on a frictionless surface. The hoop has mass M , radius R and height H . A bullet of mass μ moving horizontally with velocity V_0 strikes the hoop with impact parameter R at mid-height ($H/2$ from the surface). After the collision the bullet continues with velocity $V_0/2$ in its original direction. Ignore any hole the bullet creates. Give all your answers in terms of M , R , H , V_0 , and μ .

- What was the angular momentum, \vec{L} , of the system about the center of the hoop before the collision?
- What is the linear velocity, \vec{V} , of the center of the hoop after the collision?
- What is the angular velocity, $\vec{\omega}$, of the hoop after the collision?




Name: _____

Problem 4: Multiple Choice (15 Points)

a)

The only force acting on a 2.0-kg object moving along the x axis is shown. If the velocity v_x is -2.0 m/s at $t = 0$, what is the velocity at $t = 4.0$ s?



- a. -2.0 m/s
- b. -4.0 m/s
- c. -3.0 m/s
- d. $+1.0$ m/s
- e. $+5.0$ m/s


b)

A car of mass m_1 traveling at velocity v passes a car of mass m_2 parked at the side of the road. The momentum of the system of two cars is

- a. 0.
- b. $m_1 v$.
- c. $(m_1 - m_2)v$.
- d. $\frac{m_1 v}{m_1 + m_2}$
- e. $(m_1 + m_2)v$.

c)


Two forces of magnitude 50 N, as shown in the figure below, act on a cylinder of radius 4 m and mass 6.25 kg. The cylinder, which is initially at rest, sits on a frictionless surface. After 1 second, the velocity and angular velocity of the cylinder in m/s and rad/s are respectively



- a. $v = 0; \omega = 0.$
- b. $v = 0; \omega = 4.$
- c. $v = 0; \omega = 8.$
- d. $v = 8; \omega = 8.$
- e. $v = 16; \omega = 8.$


d)

Five objects of mass m move at velocity v at a distance r from an axis of rotation perpendicular to the page through point A, as shown below. The one that has zero angular momentum about that axis is




(a)


(b)



(c)




(d)



(e)

e)

A square of side $\frac{L}{2}$ is removed from one corner of a square sandwich that has sides of length L . The center of mass of the remainder of the sandwich moves from C to C' . The displacement of the x coordinate of the center of mass (from C to C') is



- a. $\frac{1}{12}L$.
- b. $\frac{\sqrt{2}}{12}L$.
- c. $\frac{1}{6}L$.
- d. $\frac{1}{8}L$.
- e. $\frac{\sqrt{2}}{8}L$.