


8.02X PSET 3 SOLUTIONS

PROBLEM 1

$$\begin{aligned} \text{①} \quad \text{⊕} \xrightarrow{r} \text{⊙} q \quad U_+ &= \frac{kQq}{r} \\ \text{⊕} \xrightarrow{r} \text{⊙} q \quad U_- &= -\frac{kQq}{r} \end{aligned} \quad U_{r \rightarrow \infty} = 0$$

- a) the charge $+q$ has greater potential energy at r than at ∞ .
- b) $U_+ - U_- = 2 \frac{kQq}{r}$
- c) the electric potential doesn't depend on q , $V = \frac{U_+}{q} = k \frac{Q}{r} \left(= \frac{U_-}{-q} \right)$

PROBLEM 2



conservation of energy:

$$\begin{aligned} \text{at infinity} \quad E_k &= \frac{1}{2} m v^2 \\ \text{at closest distance} \quad E_p &= k \frac{Qq}{r} \\ \hline E_{k(\infty)} &= E_{p(r)} \end{aligned}$$

giving $v = \sqrt{\frac{2kQq}{m r}}$

take $q = 1.6 \times 10^{-19} \text{ C}$
 $m = 1.7 \times 10^{-27} \text{ kg}$
 $k = 9.0 \times 10^9 \frac{\text{N m}^2}{\text{C}^2}$

$Q = 82q$
 $r = 10^{-15} \text{ m}$

we obtain $v \approx 1.5 \times 10^8 \text{ m/s}$

PROBLEM 3



$Q = Q_A + Q_B$

- a) the potential of both spheres is constant, because the charge distribution is static now. If there was a place with lower potential, the charge would flow there freely. It doesn't $\Rightarrow V = \text{const}$.

b) The potentials of the spheres are equal $V_A = k \frac{Q_A}{A} \quad V_B = k \frac{Q_B}{B} \Rightarrow \frac{Q_A}{Q_B} = \frac{A}{B}$

The ratio of electric field strengths is

$$\frac{E_A}{E_B} = \frac{k \frac{Q_A}{A^2}}{k \frac{Q_B}{B^2}} = \frac{Q_A}{Q_B} \frac{B^2}{A^2} = \frac{B}{A}$$

The electric field is stronger near the smaller sphere.