


**PROBLEM SET 3**
**DUPLICATE:** Friday, September 30, 2011

**READING ASSIGNMENT:** Steven Weinberg, *The First Three Minutes*, Chapter 3.

**FIRST QUIZ:** The first of three quizzes for the term will be given on Thursday, October 6, 2011.

**PROBLEM 1: A CYLINDRICAL UNIVERSE (10 points)**
*The following problem originated on Quiz 2 of 1994, where it counted 30 points.*

The lecture notes showed a construction of a Newtonian model of the universe that was based on a uniform, expanding, sphere of matter. In this problem we will construct a model of a cylindrical universe, one which is expanding in the  $x$  and  $y$  directions but which has no motion in the  $z$  direction. Instead of a sphere, we will describe an infinitely long cylinder of radius  $R_{\text{max},i}$ , with an axis coinciding with the  $z$ -axis of the coordinate system:



We will use cylindrical coordinates, so

$$r = \sqrt{x^2 + y^2}$$

$$\vec{r} = x\hat{i} + y\hat{j}; \quad \hat{r} = \frac{\vec{r}}{r},$$

and

where  $\hat{i}$ ,  $\hat{j}$ , and  $\hat{k}$  are the usual unit vectors along the  $x$ ,  $y$ , and  $z$  axes. We will assume that at the initial time  $t_i$ , the initial density of the cylinder is  $\rho_i$ , and the initial velocity of a particle at position  $\vec{r}$  is given by the Hubble relation

$$\vec{v}_i = H_i \vec{r}.$$

a) By using Gauss' law of gravity, it is possible to show that the gravitational acceleration at any point is given by

$$\vec{g} = -\frac{A\mu}{r} \hat{r},$$

where  $A$  is a constant and  $\mu$  is the total mass per length contained within the radius  $r$ . Evaluate the constant  $A$ .

b) As in the lecture notes, we let  $r(r_i, t)$  denote the trajectory of a particle that starts at radius  $r_i$  at the initial time  $t_i$ . Find an expression for  $\dot{r}(r_i, t)$ , expressing the result in terms of  $r$ ,  $r_i$ ,  $\rho_i$ , and any relevant constants. (Here an overdot denotes a time derivative.)

c) Defining

$$u(r_i, t) \equiv \frac{r(r_i, t)}{r_i},$$

show that  $u(r_i, t)$  is in fact independent of  $r_i$ . This implies that the cylinder will undergo uniform expansion, just as the sphere did in the case discussed in the lecture notes. As before, we define the scale factor  $a(t) \equiv u(r_i, t)$ .

d) Express the mass density  $\rho(t)$  in terms of the initial mass density  $\rho_i$  and the scale factor  $a(t)$ . Use this expression to obtain an expression for  $\ddot{a}$  in terms of  $a$ ,  $\rho$ , and any relevant constants.

e) Find an expression for a conserved quantity of the form

$$E = \frac{1}{2}\dot{a}^2 + V(a).$$

What is  $V(a)$ ? Will this universe expand forever, or will it collapse?

**PROBLEM 2: A FLAT UNIVERSE WITH UNUSUAL TIME EVOLUTION (5 points)**

Consider a *flat* universe which is filled with some peculiar form of matter, so that the Robertson–Walker scale factor behaves as

$$a(t) = bt^{3/4},$$

where  $b$  is a constant.

(a) For this universe, find the value of the Hubble expansion rate  $H(t)$ .

(b) What is the mass density of the universe,  $\rho(t)$ ? (In answering this question, you will need to know that the equation for  $\dot{a}/a$ , Eq. (3.30) in Lecture Notes 3, holds for all forms of matter, while the equation for  $\ddot{a}$ , Eq. (3.23), requires modification if the matter has a significant pressure. Eq. (3.23) is therefore not applicable to this problem.)

**PROBLEM 3: ENERGY AND THE FRIEDMANN EQUATION** (10 points)

The Friedmann equation,

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi}{3}G\rho - \frac{kc^2}{a^2}, \quad (1)$$

was derived in Lecture Notes 3 as a first integral of the equations of motion. The equation was first derived in a different form,

$$E = \frac{1}{2}\dot{a}^2 - \frac{4\pi G\rho_i}{3}a = \text{constant}, \quad (2)$$

where  $k = -2E/c^2$ . In this form the equation looks more like a conservation of energy relation, although the constant  $E$  does not have the dimensions of energy. There are two ways, however, in which the quantity  $E$  can be connected to the conservation of energy. It is related to the total energy of a test particle that moves with the Hubble expansion, and it is also related to the total energy of the entire expanding sphere of radius  $R_{\text{max}}$ , which was discussed in Lecture Notes 3 as a method of deriving the Friedmann equations. In this problem you will derive these relations.

First, to see the relation with the energy of a test particle moving with the Hubble expansion, define a physical energy  $E_{\text{phys}}$  by

$$E_{\text{phys}} \equiv m\dot{r}_i^2 E, \quad (3)$$

where  $m$  is the mass of the test particle and  $r_i$  is its initial radius. Note that the gravitational force on this particle is given by

$$\vec{F} = -\frac{GmM(r_i)}{r^2}\hat{r} = -\vec{\nabla}V_{\text{eff}}(r), \quad (4)$$

where  $M(r_i)$  is the total mass initially contained within a radius  $r_i$  of the origin,  $r$  is the present distance of the test particle from the origin, and the “effective” potential energy  $V_{\text{eff}}(r)$  is given by

$$V_{\text{eff}}(r) = -\frac{GmM(r_i)}{r}. \quad (5)$$

The motivation for calling this quantity the “effective” potential energy will be explained below.

(a) Show that  $E_{\text{phys}}$  is equal to the “effective” energy of the test particle, defined by

$$E_{\text{eff}} = \frac{1}{2}m\dot{v}^2 + V_{\text{eff}}(r). \quad (6)$$

We understand that  $E_{\text{eff}}$  is conserved because it is the energy in an analogue problem in which the test particle moves in the gravitational field of a point particle of mass  $M(r_i)$ , located at the origin, with potential energy function  $V_{\text{eff}}(r)$ . In this analogue problem the force on the test particle is exactly the same as in the real problem, but in the analogue problem the energy of the test particle is conserved.

We call (6) the “effective” energy because it is really the energy of the analogue problem, and not the real problem. The true potential energy  $V(r, t)$  of the test particle is defined to be the amount of work we must supply to move the particle to its present location from some fixed reference point, which we might take to be  $r = \infty$ . We will not bother to write  $V(r, t)$  explicitly, since we will not need it, but we point out that it depends on the time  $t$  and on  $R_{\text{max}}$ , and when differentiated gives the correct gravitational force at any radius. By contrast,  $V_{\text{eff}}(r)$  gives the correct force only at the radius of the test particle,  $r = a(t)r_i$ . The true potential energy function  $V(r, t)$  gives no conservation law, since it is explicitly time-dependent, which is why the quantity  $V_{\text{eff}}(r)$  is useful.

To relate  $E$  to the total energy of the expanding sphere, we need to integrate over the sphere to determine its total energy. These integrals are most easily carried out by dividing the sphere into shells of radius  $r$ , and thickness  $dr$ , so that each shell has a volume

$$dV = 4\pi r^2 dr. \quad (7)$$

(b) Show that the total kinetic energy  $K$  of the sphere is given by

$$K = c_K MR_{\text{max},i}^2 \left\{ \frac{1}{2}\dot{a}^2(t) \right\}, \quad (8)$$

where  $c_K$  is a numerical constant,  $M$  is the total mass of the sphere, and  $R_{\text{max},i}$  is the initial radius of the sphere. Evaluate the numerical constant  $c_K$ .

(c) Show that the total potential energy of the sphere can similarly be written as

$$U = c_U MR_{\text{max},i}^2 \left\{ -\frac{4\pi}{3}G\frac{\rho_i}{a} \right\}. \quad (9)$$

(*Suggestion:* calculate the total energy needed to assemble the sphere by bringing in one shell of mass at a time from infinity.) Show that  $c_U = c_K$ , so that the total energy of the sphere is given by

$$E_{\text{total}} = c_K MR_{\text{max},i}^2 E. \quad (10)$$

**Total points for Problem Set 3: 25.**