

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
Physics Department

Physics 8.286: The Early Universe  
Prof. Alan Guth

Quiz Date: May 9, 2002  
Post Date: May 1, 2004

### QUIZ 3 SOLUTIONS


#### PROBLEM 1: SHORT ANSWERS (40 points)

- (a) The correct answers were the  and the  The neutrino was first hypothesized by Wolfgang Pauli in 1932 (in order to explain the kinematics of beta decay), and first detected in the 1950s. After the positron was discovered in 1932, the antiproton was thought likely to exist, and the Bevatron in Berkeley was built to look for antiprotons. It made the first detection in the 1950s.
- (b) The correct answers were  The others were incorrect for the following reasons:
- (i) the earliest prediction of the CMB temperature, by Alpher and Herman in 1948, was 5 degrees, not 0.1 degrees.
  - (iii) Weinberg quotes his experimental colleagues as saying that the 3° K radiation could have been observed “long before 1965, probably in the mid-1950s and perhaps even in the mid-1940s.” To Weinberg, however, the historically interesting question is not when the radiation could have been observed, but why radio astronomers did not know that they ought to try.
  - (iv) Weinberg argues that physicists at the time did not pay attention to either the steady state model or the big bang model, as indicated by the sentence in item (v) which is a direct quote from the book: “It was extraordinarily difficult for physicists to take seriously *any* theory of the early universe”.
- (c)
- (i)  The rotation curves do not drop outside the luminous part of spiral galaxies but approach an approximately constant velocity. This indicates that there is a dark halo of matter well beyond the luminous part of the galaxy.
  - (ii)  The Doppler spreading is much *larger* than would be expected in the absence of dark matter, not smaller as was stated.
  - (iii)  The intensity and spectrum of the X-rays along a particular line of sight are used to estimate the density and temperature of the gas along that line of sight, and this information is then used with the equations of hydrostatic equilibrium to build a model of the cluster. See, for example, Claude Canizares’ article in *Astrophysics and Space Science* **267**

(1-4): 251-260, 1999, “Dark Matter in Clusters of Galaxies”. For grading, however, all students were given 2 points for this question no matter what answer was given, on the grounds that the the Rowan-Robinson textbook was not very clear on this point.

(iv) False. Most of the mass of the galaxy is in the dark matter, not stars.

(d)



(e) Supersymmetry relates bosons and fermions.

The physical distinction between these is that

- fermions obey the Pauli exclusion principle and bosons do not.

Alternatively, one could say that

- bosons are described in quantum theory by wave functions that are symmetric under the interchange of two identical particles, while fermions are described by wave functions that are antisymmetric under the interchange of two identical particles.

As a third alternative, one could say that

- bosons have integer spin (i.e., 0, 1, 2, ...) and fermions half-integer spin (i.e., 1/2, 3/2, ...).

(These three alternatives are all closely related. The second alternative is really a more detailed version of the first. The third alternative, concerning the spin, has no obvious relation to the first two, but nonetheless in relativistic quantum field theory there is a “spin-statistics” theorem which implies that

any integer-spin particle must be described by a symmetric wave function, and any half-integer-spin particle must be described by an antisymmetric one.)

- (f) Grand unified theories unify the strong, weak, and electromagnetic interactions. Only gravity is excluded.

— *Problem and solution written by Jamie Portsmouth, with some editing by AHG.*

**PROBLEM 2: NEUTRON-PROTON RATIO AND BIG-BANG NUCLEOSYNTHESIS** (20 points)


- (a) In thermal equilibrium, the ratio of neutrons to protons is given by a Boltzmann factor,

$$\boxed{\frac{n_n}{n_p} = e^{-\Delta m c^2 / kT} ,} \quad (1)$$

where  $\Delta m = (m_n - m_p)$ . For  $\Delta m c^2 = 1.293 \times 10^6$  eV,  $k = 8.617 \times 10^{-5}$  eV/K, and  $T = 5 \times 10^{10}$  K, this gives

$$\frac{n_n}{n_p} = \exp \left\{ -1.293 \times 10^6 / (8.617 \times 10^{-5} \times 5 \times 10^{10}) \right\} = \boxed{0.741 .} \quad (2)$$

*Caveat (for stat mech experts):* The above formula would be a precise consequence of statistical mechanics if the neutron and proton were two possible energy levels of the same system. In this case one would describe the system using the *canonical ensemble*, which implies that the probability of the system existing in any specific state  $i$  is proportional to  $\exp(-E_i/kT)$ , where  $E_i$  is the energy of the state. However, the neutron and proton are not really different energy levels of the same system, because the conversion between neutrons and protons involves other particles as well; a sample conversion reaction would be



where  $\nu_e$  is the electron neutrino, and  $e^-$  is the electron. This means that if the universe contained a very large density of electron neutrinos, then  $n$ - $\nu_e$  collisions would occur more frequently, and the reaction would be driven in the forward direction. Thus, a large density of electron neutrinos would lead to a lower ratio of neutrons to protons than the Boltzmann factor given above. Similarly, if the universe contained a large density of electrons, then the reaction would be driven in the reverse direction, and the ratio of neutrons to protons would be higher than the Boltzmann factor. A complete statistical mechanical treatment of this situation would use the *grand canonical ensemble*, which describes systems in which the number of particles of a given type can change by chemical reactions. In this formalism the density of each type of particle is related to a quantity called the *chemical potential*  $\mu$ , where in general the relationship is given by

$$n = g \int \frac{d^3p}{(2\pi\hbar)^3} \frac{1}{e^{(E-\mu)/(kT)} \pm 1} , \quad (3)$$

where

$$E = \sqrt{|\vec{p}|^2 c^2 + m^2 c^4} , \quad (4)$$

the + sign holds for Fermi particles, the – sign holds for Bose particles, and the factor  $g$  has the same meaning as in Lecture Notes 7. It is sometimes more convenient to use  $E$  as the variable of integration, in which case the formula above can be written as

$$n = \frac{g}{2\pi^2(\hbar c)^3} \int_m^\infty \frac{(E^2 - m^2)^{1/2}}{e^{(E-\mu)/(kT)} \pm 1} E \, dE . \quad (5)$$

For the nonrelativistic case, when  $kT \ll mc^2$ , one can neglect the term  $\pm 1$  in Eq. (3), and then the integral can be carried out exactly, yielding

$$n = g \left[ \frac{mkT}{2\pi\hbar^2} \right]^{3/2} e^{(\mu - mc^2)/(kT)} . \quad (6)$$

The ratio of neutrons to protons is then given by

$$\frac{n_n}{n_p} = e^{-(\Delta m c^2 + \mu_\nu - \mu_e)/kT} , \quad (7)$$

where  $\mu_\nu$  and  $\mu_e$  represent the chemical potentials for electron neutrinos and electrons, respectively, and we have ignored a factor  $(m_n/m_p)^{3/2} \approx 1$ . In the early universe, however, the standard theories imply that the chemical potentials for electrons and neutrinos were both negligible, so Eq. (1) above is extremely accurate.

- (b) A larger  $\Delta m$  would mean that the Boltzmann factor described in the previous answer would be smaller, so that there would be fewer neutrons at any given temperature. Fewer neutrons implies less helium, since essentially all the neutrons that exist when the temperature falls enough for deuterium to become stable become bound into helium.
- (c) There are at least four effects that occur when the electron mass/energy is taken as 1 KeV instead of 0.511 MeV:
  - (i) For the real mass/energy of 0.511 MeV the electron-positron pairs freeze out before nucleosynthesis, but a mass/energy of 1 KeV would mean that electron-positron pairs would behave as massless particles throughout the nucleosynthesis process. Just like adding an extra species of neutrino, this additional massless particle would mean that the expansion rate would be larger, since for a flat universe,


$$H^2 = \frac{8\pi}{3} G\rho ,$$

and

$$\rho = \frac{u}{c^2} = g \frac{\pi^2 (kT)^4}{30 \hbar^3 c^5} .$$


Faster expansion means that the weak interactions “freeze out” earlier, since the freeze-out point is the time at which the interactions can no longer maintain equilibrium as the universe expands. An earlier freeze-out means a higher temperature of freeze-out and hence more neutrons at the time of freeze-out. In addition, the faster expansion rate means faster cooling, which means less time before the temperature of nucleosynthesis is reached, and therefore less time for neutrons to decay. Thus, faster expansion means more neutrons. Since essentially all the neutrons present when the deuterium bottleneck breaks are collected into helium, this implies more helium.

- (ii) The most important reactions that keep protons and neutrons in thermal equilibrium all involve electrons and positrons:



If the electron-positron mass/energy were smaller, then the rates of all of these reactions would be enhanced. The reactions in which an  $e^+$  or  $e^-$  appears in the initial state will be enhanced by the presence of more  $e^+$ 's and  $e^-$ 's, and the reactions in which they appear in the final state will be enhanced because a lighter final state is easier to produce. The enhanced rate for these reactions will keep neutrons and protons in thermal equilibrium longer, and hence to lower temperatures, and this would decrease the final abundance of neutrons. Thus this effect will go in the opposite direction as effect (i), leading to the production of less helium.

- (iii) If the electron mass is decreased, then the neutron decay



becomes more exothermic, so it will happen more quickly. Thus more neutrons can decay, leading to less helium.

- (iv) As mentioned in (i), lowering the mass/energy of electron-positron pairs to 1 KeV would mean that their freeze-out would not occur until after nucleosynthesis is over. In the real case, however, with  $m_e c^2 = 0.511$  MeV, the electron-positron pairs start to freeze out at  $t \approx 10$  sec. The energy released by this freeze-out heats the photons, protons, and neutrons, and this extra heat delays the time when the universe cools enough to break the deuterium bottleneck so that helium production can proceed. The delay

allows more time for the neutrons to decay, resulting in less helium. Since the freeze-out that occurs for  $m_e c^2 = 0.511$  MeV results in less helium, the absence of this freeze-out if  $m_e c^2 = 1$  KeV would result in more helium.

Since the effects point in different directions, there is no easy way to know what the net effect will be. I (AHG) tried carrying out a full numerical integration, using the equations from P.J.E. Peebles, "Primordial helium abundance and the primordial fireball II," *Astrophysical Journal* **146**, 542-552 (1966). I found that the net effect of changing  $m_e c^2$  to 1 KeV was to produce less helium. Apparently effects (ii) and (iii) above are the most significant. Of course I did not expect students to figure this out during the exam, although some students recalled this fact from the solutions in the Review Problems.

(d) Part (a) asked for the ratio of neutrons to protons, so its answer is

$$A = \frac{n_{\text{neutron}}}{n_{\text{proton}}} .$$

The fraction of the baryonic mass in neutrons is then

$$\frac{n_{\text{neutron}}}{n_B} = \frac{n_{\text{neutron}}}{n_{\text{neutron}} + n_{\text{proton}}} = \frac{\frac{n_{\text{neutron}}}{n_{\text{proton}}}}{\frac{n_{\text{neutron}}}{n_{\text{proton}}} + 1} = \frac{A}{1 + A} .$$

The fraction of the baryonic mass in helium is twice this number, since after nucleosynthesis essentially all neutrons are in helium, and the mass of each helium nucleus is twice the mass of the neutrons within it. Thus

$$Y = \frac{2A}{1 + A} .$$

This gives  $Y = 0.851$ .

**PROBLEM 3: PRESSURE AND ENERGY DENSITY OF MYSTERIOUS STUFF** (25 points)

(a) If  $u \propto 1/\sqrt{V}$ , then one can write

$$u(V + \Delta V) = u_0 \sqrt{\frac{V}{V + \Delta V}} .$$

(The above expression is proportional to  $1/\sqrt{V + \Delta V}$ , and reduces to  $u = u_0$  when  $\Delta V = 0$ .) Expanding to first order in  $\Delta V$ ,

$$u = \frac{u_0}{\sqrt{1 + \frac{\Delta V}{V}}} = \frac{u_0}{1 + \frac{1}{2} \frac{\Delta V}{V}} = u_0 \left( 1 - \frac{1}{2} \frac{\Delta V}{V} \right) .$$

The total energy is the energy density times the volume, so

$$U = u(V + \Delta V) = u_0 \left( 1 - \frac{1}{2} \frac{\Delta V}{V} \right) V \left( 1 + \frac{\Delta V}{V} \right) = U_0 \left( 1 + \frac{1}{2} \frac{\Delta V}{V} \right) ,$$

where  $U_0 = u_0 V$  . Then

$$\Delta U = \frac{1}{2} \frac{\Delta V}{V} U_0 .$$

(b) The work done by the agent must be the negative of the work done by the gas, which is  $p \Delta V$ . So

$$\Delta W = -p \Delta V .$$

(c) The agent must supply the full change in energy, so

$$\Delta W = \Delta U = \frac{1}{2} \frac{\Delta V}{V} U_0 .$$

Combining this with the expression for  $\Delta W$  from part (b), one sees immediately that

$$p = -\frac{1}{2} \frac{U_0}{V} = -\frac{1}{2} u_0 .$$



**PROBLEM 4: AGE OF A UNIVERSE WITH MYSTERIOUS STUFF**  
(15 points)

- (a) The critical density  $\rho_c$  is defined as that density for which  $k = 0$ , where the Friedmann equation from the front of the exam implies that

$$H^2 = \frac{8\pi}{3}G\rho - \frac{kc^2}{R^2} .$$

Thus the critical density today is given by

$$\rho_c = \frac{3H_0^2}{8\pi G} .$$

The mass density today of any species  $X$  is then related to  $\Omega_{X,0}$  by

$$\rho_{X,0} = \rho_c \Omega_{X,0} = \frac{3H_0^2 \Omega_{X,0}}{8\pi G} .$$

The total mass density today is then expressed in terms of its four components as

$$\rho_0 = \frac{3H_0^2}{8\pi G} [\Omega_{m,0} + \Omega_{r,0} + \Omega_{v,0} + \Omega_{\text{ms},0}] .$$

But we also know how each of these contributions to the mass density scales with  $x(t)$ :  $\rho_m \propto 1/x^3$ ,  $\rho_r \propto 1/x^4$ ,  $\rho_v \propto 1$ , and  $\rho_{\text{ms}} \propto 1/\sqrt{V} \propto 1/x^{3/2}$ . Inserting these factors,

$$\rho(t) = \frac{3H_0^2}{8\pi G} \left[ \frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{\text{ms},0}}{x^{3/2}} \right] .$$

- (b) The Friedmann equation then becomes

$$\left( \frac{\dot{x}}{x} \right)^2 = \frac{8\pi G}{3} \frac{3H_0^2}{8\pi G} \left[ \frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{\text{ms},0}}{x^{3/2}} \right] - \frac{kc^2}{R^2} .$$

Defining

$$H_0^2 \Omega_{k,0} = -\frac{kc^2}{R^2(t_0)} ,$$

so

$$-\frac{kc^2}{R^2(t)} = -\frac{kc^2}{R^2(t_0)} \frac{1}{x^2} = \frac{H_0^2 \Omega_{k,0}}{x^2} ,$$

and then the Friedmann equation becomes

$$\left( \frac{\dot{x}}{x} \right)^2 = H_0^2 \left[ \frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{\text{ms},0}}{x^{3/2}} + \frac{\Omega_{k,0}}{x^2} \right] .$$

Applying this equation today, when  $\dot{x}/x = H_0$ , one finds that

$$\Omega_{k,0} = 1 - \Omega_{m,0} - \Omega_{r,0} - \Omega_{v,0} - \Omega_{ms,0} .$$

Rearranging the equation for  $(\dot{x}/x)^2$  above,

$$H_0 dt = \frac{dx}{x \sqrt{\frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{ms,0}}{x^{3/2}} + \frac{\Omega_{k,0}}{x^2}}} .$$

The age of the universe is found by integrating over the full range of  $x$ , which starts from 0 when the universe is born, and is equal to 1 today. So

$$t_0 = \frac{1}{H_0} \int_0^1 \frac{dx}{x \sqrt{\frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{ms,0}}{x^{3/2}} + \frac{\Omega_{k,0}}{x^2}}} .$$

*Extra Credit for Super-Sharpies (no partial credit):*

Since  $\Omega_{\text{tot}} < 1$ , we use the Robertson-Walker open universe form

$$ds^2 = -c^2 d\tau^2 = -c^2 dt^2 + R^2(t) \left\{ \frac{dr^2}{1+r^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right\} ,$$

where I have started with the general form from the front of the exam, and replaced  $k$  by  $-1$ . To discuss the radial transmission of light rays it is useful to define a new radial coordinate

$$r = \sinh \psi ,$$

so

$$dr = \cosh \psi d\psi = \sqrt{1+r^2} d\psi ,$$

where I used the identity that  $\cosh^2 \psi = 1 + \sinh^2 \psi$ . The metric can then be rewritten as

$$ds^2 = -c^2 d\tau^2 = -c^2 dt^2 + R^2(t) \{ d\psi^2 + \sinh^2 \psi (d\theta^2 + \sin^2 \theta d\phi^2) \} .$$

Light rays then travel with  $d\tau^2 = 0$ , so

$$\frac{d\psi}{dt} = \frac{c}{R(t)} .$$

If a light ray leaves the object at time  $t_e$  and arrives at Earth today, then it will travel an interval of  $\psi$  given by

$$\psi = \int_{t_e}^{t_0} \frac{c}{R(t')} dt' .$$

We will need to know  $\psi$ , but we don't know either  $t_e$  or  $R(t)$ . So we need to manipulate the right-hand side of the above equation to express it in terms of things that we do know. Changing integration variables from  $t'$  to  $x$ , where  $x = R(t')/R(t_0)$ , one finds  $dx = \dot{x} dt'$ , and then

$$\psi = \int_{x_e}^1 \frac{c}{R(t_0)} \frac{1}{x} \frac{dx}{\dot{x}} .$$

Using  $H = \dot{x}/x$ ,

$$\psi = \frac{c}{R(t_0)} \int_{x_e}^1 \frac{dx}{x^2 H} .$$

Now use the formula for  $H = \dot{x}/x$  from part (b), so

$$\psi = \frac{c}{R(t_0)H_0} \int_{x_e}^1 \frac{dx}{x^2 \sqrt{\frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{ms,0}}{x^{3/2}} + \frac{\Omega_{k,0}}{x^2}}} .$$

Here

$$x_e = \frac{R(t_e)}{R(t_0)} = \frac{1}{1+z} ,$$

and the coefficient in front of the integral can be evaluated using the Friedman equation for  $k = -1$ :

$$H_0^2 = \frac{8\pi}{3}G\rho_0 + \frac{c^2}{R^2(t_0)} = H_0^2\Omega_0 + \frac{c^2}{R^2(t_0)} ,$$

so


$$\frac{c^2}{R^2(t_0)H_0^2} = 1 - \Omega_0 = \Omega_{k,0} .$$

Finally, then, the expression for  $\psi$  can be written

$$\psi = \sqrt{\Omega_{k,0}} \int_{x_e}^1 \frac{dx}{x^2 \sqrt{\frac{\Omega_{m,0}}{x^3} + \frac{\Omega_{r,0}}{x^4} + \Omega_{v,0} + \frac{\Omega_{ms,0}}{x^{3/2}} + \frac{\Omega_{k,0}}{x^2}}} ,$$

where  $x_e$  is given by the boxed equation above.

Once we know  $\psi$ , the rest is straightforward. We draw a picture in comoving coordinates of the light rays leaving the object and arriving at Earth:



In this picture  $\Delta\theta$  is the angular size that would be measured. Using the  $d\theta^2$  part of the metric,

$$ds^2 = R^2(t) \sinh^2 \psi d\theta^2 ,$$

we can relate  $w$ , the physical size of the object at the time of emission, to  $\Delta\theta$ :

$$w = R(t_e) \sinh \psi \Delta\theta .$$

To evaluate  $R(t_e)$  we can use

$$R(t_e) = x_e R(t_0) = \frac{x_e c}{H_0 \sqrt{\Omega_{k,0}}} .$$

Finally, then,

$$\Delta\theta = \frac{w H_0 \sqrt{\Omega_{k,0}}}{x_e c \sinh \psi} ,$$

where  $\psi$  is given by the boxed equation above.