

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
Physics Department

Physics 8.286: The Early Universe  
Prof. Alan Guth

December 21, 2013

**QUIZ 3 SOLUTIONS**

**Quiz Date: December 5, 2013**

**PROBLEM 1: DID YOU DO THE READING?** (35 points)

- (a) (5 points) Ryden summarizes the results of the COBE satellite experiment for the measurements of the cosmic microwave background (CMB) in the form of three important results. The first was that, in any particular direction of the sky, the spectrum of the CMB is very close to that of an ideal blackbody. The FIRAS instrument on the COBE satellite could have detected deviations from the blackbody spectrum as small as  $\Delta\epsilon/\epsilon \approx 10^{-n}$ , where  $n$  is an integer. To within  $\pm 1$ , what is  $n$ ?

*Answer:*  $n = 4$

- (b) (5 points) The second result was the measurement of a dipole distortion of the CMB spectrum; that is, the radiation is slightly blueshifted to higher temperatures in one direction, and slightly redshifted to lower temperatures in the opposite direction. To what physical effect was this dipole distortion attributed?

*Answer:* The large dipole in the CMB is attributed to the motion of the satellite relative to the frame in which the CMB is very nearly isotropic. (The entire Local Group is moving relative to this frame at a speed of about  $0.002c$ .)

- (c) (5 points) The third result concerned the measurement of temperature fluctuations after the dipole feature mentioned above was subtracted out. Defining

$$\frac{\delta T}{T}(\theta, \phi) \equiv \frac{T(\theta, \phi) - \langle T \rangle}{\langle T \rangle},$$

where  $\langle T \rangle = 2.725$  K, the average value of  $T$ , they found a root mean square fluctuation,

$$\left\langle \left( \frac{\delta T}{T} \right)^2 \right\rangle^{1/2},$$

equal to some number. To within an order of magnitude, what was that number?

*Answer:*

$$\left\langle \left( \frac{\delta T}{T} \right)^2 \right\rangle^{1/2} = 1.1 \times 10^{-5}.$$

- (d) (5 points) Which of the following describes the Sachs-Wolfe effect?
- (i) Photons from fluid which had a velocity toward us along the line of sight appear redder because of the Doppler effect.
  - (ii) Photons from fluid which had a velocity toward us along the line of sight appear bluer because of the Doppler effect.
  - (iii) Photons from overdense regions at the surface of last scattering appear redder because they must climb out of the gravitational potential well.
  - (iv) Photons from overdense regions at the surface of last scattering appear bluer because they must climb out of the gravitational potential well.
  - (v) Photons traveling toward us from the surface of last scattering appear redder because of absorption in the intergalactic medium.
  - (vi) Photons traveling toward us from the surface of last scattering appear bluer because of absorption in the intergalactic medium.

- (e) (5 points) The flatness problem refers to the extreme fine-tuning that is needed in  $\Omega$  at early times, in order for it to be as close to 1 today as we observe. Starting with the assumption that  $\Omega$  today is equal to 1 within about 1%, one concludes that at one second after the big bang,

$$|\Omega - 1|_{t=1 \text{ sec}} < 10^{-m} ,$$

where  $m$  is an integer. To within  $\pm 3$ , what is  $m$ ?

*Answer:*  $m = 18$ . (See the derivation in Lecture Notes 8.)

- (f) (5 points) The total energy density of the present universe consists mainly of baryonic matter, dark matter, and dark energy. Give the percentages of each, according to the best fit obtained from the Planck 2013 data. You will get full credit if the first (baryonic matter) is accurate to  $\pm 2\%$ , and the other two are accurate to within  $\pm 5\%$ .

*Answer:* Baryonic matter: 5%. Dark matter: 26.5%. Dark energy: 68.5%. The Planck 2013 numbers were given in Lecture Notes 7. To the requested accuracy, however, numbers such as Ryden's Benchmark Model would also be satisfactory.

- (g) (5 points) Within the conventional hot big bang cosmology (without inflation), it is difficult to understand how the temperature of the CMB can be correlated at angular separations that are so large that the points on the surface of last scattering was separated from each other by more than a horizon distance. Approximately what angle, in degrees, corresponds to a separation on the surface last scattering of one horizon length? You will get full credit if your answer is right to within a factor of 2.

*Answer:* Ryden gives  $1^\circ$  as the angle subtended by the Hubble length on the surface of last scattering. For a matter-dominated universe, which would be a good model for our universe, the horizon length is twice the Hubble length. Any number from  $1^\circ$  to  $5^\circ$  was considered acceptable.

**PROBLEM 2: FREEZE-OUT OF MUONS** (25 points)

See the solutions to Problem Set 7, Problem 2 (2013).

**PROBLEM 3: THE EVENT HORIZON FOR OUR UNIVERSE** (25 points)

- (a) In a spherical pulse each light ray is moving radially outward, so  $d\theta = d\phi = 0$ . A light ray travels along a null trajectory, meaning that  $ds^2 = 0$ , so we have

$$ds^2 = -c^2 dt^2 + a^2(t) dr^2 = 0 . \quad (3.1)$$

from which it follows that

$$\frac{dr}{dt} = \pm \frac{c}{a(t)} . \quad (3.2)$$

We are interested in a radial pulse that starts at  $r = 0$  at time  $t = t_0$ , so the limiting value of  $r$  is given by

$$r_{\max} = \int_{t_0}^{\infty} \frac{c}{a(t)} dt . \quad (3.3)$$

- (b) Changing variables of integration to

$$x = \frac{a(t)}{a(t_0)} , \quad (3.4)$$

the integral becomes

$$r_{\max} = \int_1^{\infty} \frac{c}{a(t)} \frac{dt}{dx} dx = \frac{c}{a(t_0)} \int_1^{\infty} \frac{1}{x} \frac{dt}{dx} dx , \quad (3.5)$$

where we used the fact that  $t = t_0$  corresponds to  $x = a(t_0)/a(t_0) = 1$ . As given to us on the formula sheet, the first-order Friedmann equation can be written as

$$x \frac{dx}{dt} = H_0 \sqrt{\Omega_{m,0} x + \Omega_{\text{rad},0} + \Omega_{\text{vac},0} x^4 + \Omega_{k,0} x^2} . \quad (3.6)$$

Using this substitution,

$$r_{\max} = \frac{c}{a(t_0)H_0} \int_1^{\infty} \frac{dx}{\sqrt{\Omega_{m,0} x + \Omega_{\text{rad},0} + \Omega_{\text{vac},0} x^4}} , \quad (3.7)$$

where we have used  $\Omega_{k,0} = 0$ , since the universe is taken to be flat.

- (c) To find the value of the redshift for the light that we are presently receiving from coordinate distance  $r_{\max}$ , we can begin by noticing that the time of emission  $t_e$  can be determined by the equation which implies that the coordinate distance traveled by a light pulse between times  $t_e$  and  $t_0$  must equal  $r_{\max}$ . Using Eq. (3.2) for the coordinate velocity of light, this equation reads

$$\int_{t_e}^{t_0} \frac{c}{a(t)} dt = r_{\max} . \quad (3.8)$$

The “half-credit” answer to the quiz problem would include the above equation, followed by the statement that the redshift  $z_{\text{eh}}$  can be determined from

$$z = \frac{a(t_0)}{a(t_e)} - 1 . \quad (3.9)$$

The “full-credit” answer is obtained by changing the variable of integration as in part (b), so Eq. (3.8) becomes

$$\begin{aligned} r_{\max} &= \int_{x_e}^1 \frac{c}{a(t)} \frac{dt}{dx} dx \\ &= \frac{c}{a(t_0)} \int_{x_e}^1 \frac{1}{x} \frac{dt}{dx} dx , \end{aligned} \quad (3.10)$$

where  $x_e$  is the value of  $x$  corresponding to  $t = t_e$ . Then using Eq. (3.6) with  $\Omega_{k,0} = 0$ , we find

$$r_{\max} = \frac{c}{a(t_0)H_0} \int_{x_e}^1 \frac{dx}{\sqrt{\Omega_{m,0}x + \Omega_{\text{rad},0} + \Omega_{\text{vac},0}x^4}} . \quad (3.11)$$

To complete the answer in this language, we use

$$z = \frac{1}{x_e} - 1 . \quad (3.12)$$

Eqs. (3.11) and (3.12) constitute a full answer to the question, but one could go further and replace  $r_{\max}$  using Eq. (3.7), finding

$$\begin{aligned} &\int_1^{\infty} \frac{dx}{\sqrt{\Omega_{m,0}x + \Omega_{\text{rad},0} + \Omega_{\text{vac},0}x^4}} \\ &= \int_{x_e}^1 \frac{dx}{\sqrt{\Omega_{m,0}x + \Omega_{\text{rad},0} + \Omega_{\text{vac},0}x^4}} . \end{aligned} \quad (3.13)$$

In this form the answer depends only on the values of  $\Omega_{X,0}$ .

You were of course not asked to evaluate this formula numerically, but you might be interested in knowing that the Planck 2013 values  $\Omega_{m,0} = 0.315$ ,  $\Omega_{\text{vac},0} = 0.685$ , and  $\Omega_{\text{rad},0} = 9.2 \times 10^{-5}$  lead to  $z_{\text{eh}} = 1.87$ . Thus, no event that is happening now (i.e., at the same value of the cosmic time) in a galaxy at redshift larger than 1.87 will ever be visible to us or our descendants, even in principle.

**PROBLEM 4: BEHAVIOR OF  $\Omega$  IN A UNIVERSE DOMINATED BY MYSTERIOUS STUFF** (15 points)

(a) From the Friedmann equation,

$$H^2 = \frac{8\pi}{3}G\rho - \frac{kc^2}{a^2} ,$$

and the definition of  $\rho_c$ ,

$$H^2 = \frac{8\pi}{3}G\rho_c ,$$

we have

$$\frac{8\pi}{3}G\rho - \frac{kc^2}{a^2} = \frac{8\pi}{3}G\rho_c ,$$

which can be rearranged to give

$$\rho - \rho_c = \frac{3kc^2}{8\pi Ga^2} .$$

Dividing both sides by  $\rho$  we find

$$\frac{\rho - \rho_c}{\rho} = \frac{\Omega - 1}{\Omega} = \frac{3kc^2}{8\pi Ga^2\rho} .$$

(The middle expression above is obtained from the expression on the left by multiplying both the numerator and the denominator by  $1/\rho_c$ .) Now multiply both the numerator and denominator of the expression on the right by  $T^2$ , recognizing that  $aT$  is approximately constant. We then have

$$\frac{\Omega - 1}{\Omega} = A \frac{T^2}{\rho} ,$$

where

$$A = \frac{3kc^2}{8\pi G(aT)^2} .$$

- (b) Combining the above equation with our knowledge that  $T \propto 1/a$  and  $\rho \propto 1/a^5$ , we have immediately that

$$\frac{\Omega - 1}{\Omega} \propto \frac{1/a^2}{1/a^5} \propto a^3 .$$

To determine how  $a$  depends on time, we can solve the Friedmann equation for a flat universe dominated by mysterious stuff:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{\text{const}}{a^5} \implies \frac{da}{dt} = \frac{\sqrt{\text{const}}}{a^{3/2}} .$$

Thus

$$a^{3/2} da = \sqrt{\text{const}} dt$$

and then

$$\frac{2}{5} a^{5/2} = \sqrt{\text{const}} t ,$$

where the constant of integration is set to zero by using the convention that  $a = 0$  when  $t = 0$ . Thus

$$a \propto t^{2/5} ,$$

and then

$$\boxed{\frac{\Omega - 1}{\Omega} \propto a^3 \propto t^{6/5} .}$$