8.286 Lecture 12
October 22, 2013

NON-EUCLIDEAN SPACES: OPEN UNIVERSES AND THE SPACETIME METRIC

Summary of Lecture 11: Surface of a Sphere



$$x^2 + y^2 + z^2 = R^2 \ .$$

Alan Guth
Massachusetts Institute of Technology

Polar Coordinates:

$$x = R \sin\theta \cos\phi$$
$$y = R \sin\theta \sin\phi$$
$$z = R \cos\theta \ ,$$



Varying θ:

$$ds = R\,d\theta$$



Varying ϕ:

$$ds = R \sin\theta \, d\phi$$



Varying θ and ϕ

Varying θ: $\quad ds = R\,d\theta$

Varying ϕ: $\quad ds = R\sin\theta\,d\phi$

$$ds^2 = R^2\left(d\theta^2 + \sin^2\theta\,d\phi^2\right)$$

Review of Lecture 11: A Closed Three-Dimensional Space

$$x^2 + y^2 + z^2 + w^2 = R^2$$

$$\begin{aligned} x &= R \sin\psi \sin\theta \cos\phi \\ y &= R \sin\psi \sin\theta \sin\phi \\ z &= R \sin\psi \cos\theta \\ w &= R \cos\psi \ , \end{aligned}$$

$$ds = R\, d\psi$$



Metric for the Closed 3D Space

Varying ψ: $\quad ds = R\,d\psi$

Varying θ or ϕ: $\quad ds^2 = R^2 \sin^2\psi (d\theta^2 + \sin^2\theta\, d\phi^2)$

If the variations are orthogonal to each other, then

$$ds^2 = R^2 \left[d\psi^2 + \sin^2\psi \left(d\theta^2 + \sin^2\theta\, d\phi^2 \right) \right]$$

Proof of Orthogonality of Variations

Let $d\vec{R}_\psi$ = displacement of point when ψ is changed to $\psi + d\psi$.

Let $d\vec{R}_\theta$ = displacement of point when θ is changed to $\theta + d\theta$.

- ☆ $d\vec{R}_\theta$ has no w-component $\quad\Longrightarrow\quad d\vec{R}_\psi \cdot d\vec{R}_\theta = d\vec{R}_\psi^{(3)} \cdot d\vec{R}_\theta^{(3)}$, where (3) denotes the projection into the x-y-z subspace.
- ☆ $d\vec{R}_\psi^{(3)}$ is radial; $d\vec{R}_\theta^{(3)}$ is tangential

$$\Longrightarrow \quad d\vec{R}_\psi^{(3)} \cdot d\vec{R}_\theta^{(3)} = 0$$

Review of Lecture 11: Implications of General Relativity

- $ds^2 = R^2\left[d\psi^2 + \sin^2\psi\left(d\theta^2 + \sin^2\theta\, d\phi^2\right)\right]$, where R is radius of curvature.
- According to GR, matter causes space to curve.
- R cannot be arbitrary. Instead, $R^2(t) = \dfrac{a^2(t)}{k}$.
- Finally,

$$\boxed{\mathrm{d}s^2 = a^2(t)\left\{\frac{\mathrm{d}r^2}{1-kr^2} + r^2\left(\mathrm{d}\theta^2 + \sin^2\theta\,\mathrm{d}\phi^2\right)\right\}\ ,}$$

where $r = \dfrac{\sin\psi}{\sqrt{k}}$. Called the Robertson-Walker metric.