


8.286 Lecture 10
October 17, 2016

INTRODUCTION TO NON-EUCLIDEAN SPACES

Announcements

- ★ Quiz 1: Average = 77.2% — very good! There is a histogram on the website, which shows number-to-letter conversions. The number-to-letter conversions will be different for each quiz and for the homework, but your letter grades should be a reliable indicator of where you stand.
- ★ Grading questions: we aim to grade fairly, but we make mistakes. By all means feel free to ask me if you have any questions about the grading of your quizzes. The solutions are posted.
- ★ Problem Set 4 is due today (Monday), 4:00 pm.
- ★ Problem Set 5 is due Friday, 4:00 pm.




curved space ?

curves with respect to what ?

curved spacetime ?

non-Euclidean geometry



Slide created by Mustafa Amin

Note on image: I have added the "Dali" clock and Fermi satellite images to the original image created for the Gravity Probe B collaboration

-2-

5

Slide created by Mustafa Amin

Euclid's *Postulates*



1. A straight line segment can be drawn joining any two points.
2. Any straight line segment can be extended indefinitely in a straight line.
3. Given a straight line segment, a circle can be drawn having the segment as radius and one endpoint as center.
4. All right angles are congruent.
5. If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines if produced indefinitely meet on that side on which the angles are less than two right angles


Corrected 10/10/13

-3-

7

Slide created by Mustafa Amin

5th Postulate




5. If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines if produced indefinitely meet on that side on which the angles are less than two right angles

Corrected 10/10/13

-4-

Equivalent Statements of the 5th Postulate




(a) "If a straight line intersects one of two parallels (i.e., lines which do not intersect however far they are extended), it will intersect the other also."

MIT Alan Guth
Massachusetts Institute of Technology
8.286 Lecture 10, October 17, 2016

-5-

Equivalent Statements of the 5th Postulate




(b) "There is one and only one line that passes through any given point and is parallel to a given line."

MIT Alan Guth
Massachusetts Institute of Technology
8.286 Lecture 10, October 17, 2016

-6-

Equivalent Statements of the 5th Postulate




(c) "Given any figure there exists a figure, similar to it, of any size." (Two polygons are similar if their corresponding angles are equal, and their corresponding sides are proportional.)

MIT Alan Guth
Massachusetts Institute of Technology
8.286 Lecture 10, October 17, 2016


-7-

Equivalent Statements of the 5th Postulate



(d) "There is a triangle in which the sum of the three angles is equal to two right angles (i.e., 180°)."

Giovanni Gerolamo Saccheri (1667-1733)



In 1733, Saccheri, a Jesuit priest, published *Euclides ab omni naevo vindicatus* (*Euclid Freed of Every Flaw*).

The book was a study of what geometry would be like if the 5th postulate were false.

He hoped to find an inconsistency, but failed.

Carl Friedrich Gauss (1777-1855)




German mathematician and physicist.

Born as the son of a poor working-class parents. His mother was illiterate and never even recorded the date of his birth.

His students included Richard Dedekind, Bernhard Riemann, Peter Gustav Lejeune Dirichlet, Gustav Kirchhoff, and August Ferdinand Möbius.


~1750-1850



- infinite
- constant negative curvature



~~5th Postulate~~



Slide created by Mustafa Amin

Slide created by Mustafa Amin

GBL geometry with Klein



1. constant negative curvature
2. infinite
3. ~~5th postulate~~



(x_1, y_1)

$$x^2 + y^2 < 1$$

(x_2, y_2)

$$d(1, 2) = a \cosh^{-1} \left[\frac{1 - x_1 x_2 - y_1 y_2}{\sqrt{1 - x_1^2 - y_1^2} \sqrt{1 - x_2^2 - y_2^2}} \right]$$


Note: no global embedding in 3D Euclidean space possible

-12-

12

Slide created by Mustafa Amin

Geometry (after Klein)



co-ordinates

(x_1, y_1)

(x_2, y_2)

distance function


$$d[(x_1, y_1), (x_2, y_2)]$$

-13-

13

Slide created by Mustafa Amin

tiny distances



$$ds^2 = dx^2 + dy^2$$


$$ds^2 = g_{xx}(x, y)dx^2 + 2g_{xy}(x, y)dxdy + g_{yy}(x, y)dy^2$$

-14-

14

Slide created by Mustafa Amin

quadratic form




$$ds^2 = g_{xx}(x, y)dx^2 + 2g_{xy}(x, y)dxdy + g_{yy}(x, y)dy^2$$

-15-

15

Slide created by Mustafa Amin

locally Euclidean




$$ds^2 = dx'^2 + dy'^2$$

$$ds^2 = g_{xx}(x, y)dx^2 + 2g_{xy}(x, y)dxdy + g_{yy}(x, y)dy^2$$

$$g_{xx}g_{yy} - g_{xy}^2 > 0$$

-16-


Non-Euclidean Geometry: The Surface of a Sphere



$$x^2 + y^2 + z^2 = R^2$$


-17-

Polar Coordinates:



-18-

Varying θ :



-19-

