

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
Physics Department

**8.323: Relativistic Quantum Field Theory I**

Prof. Alan Guth

March 13, 2008

**PROBLEM SET 5**

**DUE DATE:** Thursday, March 20, 2008, at 5:00 p.m. in the 8.323 homework box, at the intersection of the 3rd floor of building 8 and the 4th floor of building 16.

**REFERENCES:** *Lecture Notes #4: Dirac Delta Function as a Distribution*, on the website. Peskin and Schroeder, Sec. 2.4; Optional reference: *Quantum Field Theory*, by Lowell Brown, section 1.7 (coherent states).

**Problem 1: Subtleties of delta functions** (10 points)

- (a) Consider  $g_1(t)$  and  $g_2(t)$  defined by

$$g_1(t) \equiv f(t) \delta(t - a) \quad (1.1)$$

and

$$g_2(t) \equiv f(a) \delta(t - a) , \quad (1.2)$$

where  $f(t)$  is an arbitrary smooth function,  $a$  is a constant, and  $\delta$  denotes the Dirac delta function. By “smooth,” I mean that  $f(t)$  is continuous, and differentiable as many times as might be necessary for any issue that arises. Note that  $g_1(t)$  and  $g_2(t)$  are distributions, defined more explicitly as

$$T_{g_i}[\varphi] \equiv \int_{-\infty}^{\infty} dt g_i(t) \varphi(t) , \quad (1.3)$$

where  $\varphi(t)$  is a test function. By evaluating the functionals shown in Eq. (1.3) for an arbitrary allowed test function, find out if  $g_1(t)$  and  $g_2(t)$  are equal to each other as distributions.

- (b) Using a prime to indicate that a function is differentiated with respect to its argument, consider the distributions  $h_1(t)$  and  $h_2(t)$  defined by

$$h_1(t) \equiv f(t) \delta'(t - a) \quad (1.4)$$

and

$$h_2(t) \equiv f(a) \delta'(t - a) , \quad (1.5)$$

where again  $f(t)$  is an arbitrary smooth function and  $\delta$  denotes the Dirac delta function. The derivative of a distribution was defined in Lecture Notes 4. By

evaluating these two distributions for an arbitrary allowed test function  $\varphi(t)$ , find out if they are equal to each other as distributions.

- (c) Now consider the derivatives of the distributions defined above in Eqs. (1.1) and (1.2):

$$g_1'(t) = f'(t) \delta(t-a) + f(t) \delta'(t-a) \quad (1.6)$$

and

$$g_2'(t) = f(a) \delta'(t-a) . \quad (1.7)$$

**IF** you concluded that  $g_1(t)$  and  $g_2(t)$  are equal, then you should certainly expect that their derivatives should be equal, even if they do not appear to be identical. Find out if  $g_1'(t)$  and  $g_2'(t)$  are equal.

- (d) Let the function  $\theta(t)$  be defined in the standard way,

$$\theta(t) = \begin{cases} 1 & \text{if } t > 0 \\ 0 & \text{otherwise,} \end{cases} \quad (1.8)$$

and consider the corresponding distribution

$$T_\theta[\varphi] \equiv \int_{-\infty}^{\infty} dt \theta(t) \varphi(t) . \quad (1.9)$$

Define the derivative of a  $\theta$ -function as a distribution, so

$$\int_{-\infty}^{\infty} dt \theta'(t) \varphi(t) \equiv T'_\theta[\varphi] \equiv -T_\theta \left[ \frac{d\varphi}{dt} \right] . \quad (1.10)$$

Show that

$$\theta'(t) = \delta(t) . \quad (1.11)$$

**[Warning:** The product of two distributions cannot be defined in general, nor can the square of a distribution. As a function, it is clear from the definition (1.8) that

$$\theta^n(t) = \theta(t) , \quad (1.12)$$

where  $n$  denotes any positive integer. Any function that is piecewise continuous and bounded by a power can be promoted to a corresponding distribution, so  $T_\theta[\varphi]$  and  $T_{\theta^n}[\varphi]$  are both well-defined distributions and are equal to each other. One might conjecture that

$$\frac{d}{dt} \theta(t) = \delta(t) , \quad (1.13a)$$

$$\frac{d}{dt} \theta^2(t) \stackrel{?}{=} 2 \theta(t) \delta(t) , \quad (1.13b)$$

and

$$\frac{d}{dt} \theta^3(t) \stackrel{?}{=} 3 \theta^2(t) \delta(t) . \quad (1.13c)$$

Eq. (1.13a) is identical to Eq. (1.11) and is correct, but the right-hand sides of Eqs. (1.13b) and (1.13c) are ill-defined. Since  $\theta^n(t) = \theta(t)$ , the left-hand sides of Eqs. (1.13a) – (1.13c) are all well-defined and are equal to each other. It is hard to imagine any consistent definition that would make the right-hand sides equal, so the standard approach is to consider them undefined.]

**Problem 2:**  $(\square_x + m^2) D_F(x - y) = -i\delta^{(4)}(x - y)$  (10 points)

The Feynman propagator for a free scalar field  $\phi(x)$  is defined by

$$D_F(x - y) = \theta(x^0 - y^0) \langle 0 | \phi(x) \phi(y) | 0 \rangle + \theta(y^0 - x^0) \langle 0 | \phi(y) \phi(x) | 0 \rangle ,$$

where  $|0\rangle$  denotes the vacuum state. Use the canonical commutation relations to show that

$$(\square_x + m^2) D_F(x - y) = -i\delta^{(4)}(x - y) ,$$

where

$$\square_x \equiv \eta^{\mu\nu} \frac{\partial}{\partial x^\mu} \frac{\partial}{\partial x^\nu} .$$

[*Suggestion:* First calculate

$$\frac{\partial}{\partial x^0} D_F(x - y) ,$$

and then differentiate this expression again to find

$$\frac{\partial^2}{\partial (x^0)^2} D_F(x - y) .$$

Then add in the other terms to find the final answer.]

**Problem 3: Coherent states** (15 points)

In lecture we solved the problem of a quantized scalar field  $\phi(x)$  interacting with a fixed classical source  $j(x)$ ,

$$(\square + m^2)\phi(x) = j(x) .$$

We found that the in and out operators are related by

$$\phi_{\text{out}}(x) = S^{-1} \phi_{\text{in}}(x) S$$

$$a_{\text{out}}(\vec{p}) = S^{-1} a_{\text{in}}(\vec{p}) S ,$$

where  $S$  can be written

$$S = e^{-\frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_p} |\tilde{j}(p)|^2} e^F e^G ,$$

where

$$F = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \tilde{j}(p) a_{\text{in}}^\dagger(\vec{p})$$

$$G = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \tilde{j}(-p) a_{\text{in}}(\vec{p}) ,$$

and

$$\tilde{j}(p) \equiv \int d^4 y e^{ip \cdot y} j(y) .$$

(a) Show that  $S$  can also be written as

$$S = e^{-\frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_p} |\tilde{j}(p)|^2} e^{F'} e^{G'} ,$$

where

$$F' = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \tilde{j}(p) a_{\text{out}}^\dagger(\vec{p})$$

$$G' = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \tilde{j}(-p) a_{\text{out}}(\vec{p}) ,$$

and hence that

$$|0_{\text{in}}\rangle = e^{-\frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_p} |\tilde{j}(p)|^2} e^{F'} |0_{\text{out}}\rangle .$$

Note that the right-hand-side of the above equation gives a useful description of the final state, since the out operators have a straightforward interpretation at late times. States of this form — exponentials of creation operators acting on the vacuum — are called coherent states.

(b) To study further the properties of coherent states, it is useful to consider a single harmonic oscillator,

$$H = \frac{p^2}{2m} + \frac{\omega^2 m}{2} q^2 ,$$

which is simplified by the canonical transformation

$$p = \sqrt{m\omega} \bar{p} , \quad q = \frac{1}{\sqrt{m\omega}} \bar{q} .$$

Dropping the overbars, the creation and annihilation operators are then given by

$$a^\dagger = \frac{1}{\sqrt{2}}(q - ip)$$

$$a = \frac{1}{\sqrt{2}}(q + ip) .$$

A coherent state  $|z\rangle$  can be defined by

$$|z\rangle \equiv e^{za^\dagger} |0\rangle .$$

Show that  $|z\rangle$  is an eigenstate of the annihilation operator, and find its eigenvalue.

- (c) Show that  $\langle z_2 | z_1 \rangle = e^{z_2^* z_1}$ . (If you look at Lowell Brown's book, note that I am defining  $\langle z|$  to be the bra vector that corresponds to the ket  $|z\rangle$ , so my  $\langle z|$  is equal to Brown's  $\langle z^*|.)$

- (d) Find

$$\langle q \rangle_z \equiv \frac{\langle z | q | z \rangle}{\langle z | z \rangle}$$

and

$$\langle p \rangle_z \equiv \frac{\langle z | p | z \rangle}{\langle z | z \rangle} .$$

- (e) Compute the standard deviations of  $q$  and  $p$ ,

$$\Delta q^2 = \langle (q - \langle q \rangle)^2 \rangle$$

$$\Delta p^2 = \langle (p - \langle p \rangle)^2 \rangle$$

and show that  $|z\rangle$  is a minimal-uncertainty state, in the sense that

$$\Delta q \Delta p = \frac{1}{2} .$$