

8.962 Lecture 10
March 12, 2018

MATHEMATICAL CONCEPTS and DEFINITIONS

Basic Notation

- \cup $A \cup B$ denotes the union of sets A and B
- \cap $A \cap B$ denotes the intersection of the sets A and B
- \subset $A \subset B$ denotes that A is a subset of B .
(May or may not mean proper subset.)
- $-$ $B - A$ denotes the complement in B of the set A
- \in $p \in A$ denotes that p is an element of A
- $\{|\}$ $\{p \in A | Q\}$ denotes the set consisting of those elements p of the set A which satisfy condition Q
- \times Cartesian product; $A \times B$ is the set $\{(a, b) | a \in A \text{ and } b \in B\}$
- \emptyset the empty set

- \mathbb{R} the set of real numbers
- \mathbb{R}^n the set of n -tuples of real numbers
- \mathbb{C} the set of complex numbers
- \mathbb{C}^n the set of n -tuples of complex numbers
- $\colon\rightarrow$ $f : A \rightarrow B$ denotes that f is a map from the set A to the set B
- $f \circ g$ denotes the composition of maps $g : A \rightarrow B$ and $f : B \rightarrow C$, i.e. ., for $p \in A$ we have
- $$(f \circ g)(p) = f[g(p)]$$
- [] $f[A]$ denotes the image of the set A under the map f , i.e., the set $\{f(x) | x \in A\}$

- C^n the set of n -times continuously differentiable functions.
Note that C^0 means simply continuous, while C^1 means that the first derivative exists and is continuous.
- C^∞ the set of infinitely continuously differentiable (i.e., smooth) functions
- \exists there exists; i.e., for all $u \in \mathbb{R}$, $\exists v \mid v + u = 0$
- \forall for all; i.e., $\forall u \in \mathbb{R}$, $\exists v \mid v + u = 0$

Properties of Maps

- ★ If f is a function $f : M \rightarrow N$, M is called the **domain** of f , and N is called its **codomain**.
- ★ The set of points in N that M gets mapped into is called the **image** of f .
- ★ For any subset $U \subset N$, the set of elements of M that get mapped to U is called the **preimage** of U under f , or $f^{-1}(U)$.
- ★ A map $f : M \rightarrow N$ is called **one-to-one** (or **injective**) if each element of N has at most one element of M mapped into it.
- ★ A map $f : M \rightarrow N$ is called **onto** (or **surjective**) if each element of N has at least one element of M mapped into it.

★ A map that is both one-to-one and onto is known as **invertible** (or **bijective**). In this case we can define the inverse map $f^{-1} : N \rightarrow M$ by $(f^{-1} \circ f)(x) = x$, for any $x \in M$.

Continuity

- ★ (Weierstrass definition): For functions $f : D \rightarrow \mathbb{R}$, where $D \subset \mathbb{R}$, $f(x)$ is continuous at x_0 if and only if for every $\epsilon > 0$ there exists a $\delta > 0$ such that $|x - x_0| < \delta \implies |f(x) - f(x_0)| < \epsilon$.
- ★ (General topological definition): If open sets have been defined, then a function $f : X \rightarrow Y$ is continuous if and only if the preimage $f^{-1}(V)$, where V is an open subset of Y (which could be the whole set), is always an open subset of X .
- ★ For the usual definition of open sets on \mathbb{R} , the two definitions are equivalent.

- ★ If $f : D \rightarrow \mathbb{R}^n$, where $D \subset \mathbb{R}^m$, then the definition of continuity is a natural generalization of the $\mathbb{R} \rightarrow \mathbb{R}$ definition. f can be described as a collection of functions

$$f^i(x^1, x^2, \dots, x^m) ,$$

where $i = 1, \dots, n$. f is C^p if each f^i is at least C^p in each of the variables (x^1, x^2, \dots, x^m) .

- ★ Suppose that M and N are topological spaces (i.e., spaces on which open sets have been defined). Then if $f : M \rightarrow N$ is continuous, one-to-one, and onto, and its inverse is continuous, then f is called a **homeomorphism**, and the spaces M and N are said to be **homeomorphic**. As far as topology is concerned, M and N are then identical. (See Wald. Carroll never uses the word “homeomorphic”.)

- ★ Suppose that M and N are manifolds (to be defined shortly). Then if $f : M \rightarrow N$ is C^∞ , one-to-one, and onto, and its inverse is C^∞ , then f is called a **diffeomorphism**, and the spaces M and N are said to be **diffeomorphic**. As far as manifold properties are concerned, M and N are then identical.*
- ★ An **open ball** is the set of all points x in \mathbb{R}^n such that $|x - y| < r$ for some fixed $y \in \mathbb{R}^n$ and $r \in \mathbb{R}$, where $|x - y|^2 = \sum_i (x^i - y^i)^2$. Note that $|x - y|$ must be less than r . The ball does not include its boundary.
- ★ An **open set** in \mathbb{R}^n is a set constructed from an arbitrary (maybe infinite) union of open balls. Equivalently, a set $V \subset \mathbb{R}^n$ is open if, for any $y \in V$, there is an open ball centered at y that is completely inside V .

*This entry has been corrected from the version shown in lecture, which mistakenly omitted the requirement that f and f^{-1} must be C^∞ .

MANIFOLDS!

★ If M is a set, a **chart** or **coordinate system** on M is a one-to-one map $\phi : U \rightarrow \mathbb{R}^n$ such that the image $\phi(U)$ of the map is an open subset of \mathbb{R}^n . We do not assume a topology on M , so U is said to be open in M if $\phi(U)$ is open in \mathbb{R}^n .




FIGURE 2.13 A coordinate chart covering an open subset U of M .

(From Sean Carroll, *Spacetime and Geometry*.)

★ A C^∞ **atlas** of charts is a collection of charts $\{(U_\alpha, \phi_\alpha)\}$ that satisfies two conditions:

- 1) The U_α cover M , so that any point in M is contained in at least one chart U_α .
- 2) The charts smoothly sew together. Whenever two charts overlap, the map from one coordinate system to the other must be C^∞ . In symbols, whenever $U_\alpha \cap U_\beta \neq \emptyset$, the map $(\phi_\alpha \circ \phi_\beta^{-1})$ takes points in $\phi_\beta(U_\alpha \cap U_\beta) \subset \mathbb{R}^n$ onto the open set $\phi_\alpha(U_\alpha \cap U_\beta) \subset \mathbb{R}^n$. All such maps must be C^∞ where they are defined.




FIGURE 2.14 Overlapping coordinate charts.
(From Sean Carroll, *Spacetime and Geometry*.)

A C^∞ n -dimensional **manifold** (or n -manifold for short) is simply a set M along with a maximal atlas, one that contains every possible compatible chart.