

Proof of a graph-theoretic lemma

December 12, 2007

Question: Let G be a simple undirected d -regular graph, and let H be a subset of the vertices of G . Consider a random walk starting from a vertex s in H . At each step, the walk stays at the current vertex with probability $\frac{1}{2}$ and it moves to each neighbor of the current vertex with probability $\frac{1}{2d}$. If the starting vertex s is chosen uniformly at random from H , what is the expected probability that the walk leaves H immediately and takes more than ℓ steps to return to a vertex in H ?

Answer: At most $\frac{|G|}{|H|} \frac{1}{\ell}$.

I think the lemma and its proof is interesting for a couple of reasons. First, the upper-bound holds unconditionally for *any* simple undirected d -regular graph and any $\ell > 0$. Second, the proof makes elegant use of the stationary distribution of the walk, while in the question, there is no mention of it.

Here is a rough sketch of the proof. As mentioned above, we look at the stationary distribution of the random walk on G . By the way the walk is defined, the stationary distribution assigns probability $\frac{1}{|G|}$ to every vertex. Now, we ask the following question: for a given u and v in H and for a given $t > \ell$, what is the probability under the stationary distribution of being on a vertex that lies on a path of length¹ t between u and v that passes only through vertices not in H ? This probability equals t times the probability of entering such a path of length t from u to v (after the walk has totally mixed). The latter probability is the probability of being at u , that is $1/|G|$, times the probability of a walk outside H of length t that starts at u and ends at v . Summing over all u and v in H and all $t > \ell$, we find that the probability of walks staying outside H and longer than ℓ between some two vertices in H is at most $|G|/\ell$ times the probability under the stationary distribution of being on some vertex that lies on a path of at least ℓ between two vertices in H . The latter probability, we argue, is at most 1, and that proves our claim. Details follow.

Proof. For convenience, let us suppose that we add d self-loops to each vertex in G . Then, a random walk on G consists of uniformly choosing one of the $2d$ edges incident to the current vertex at each step. Thus, there are exactly $(2d)^t$ distinct paths of length t starting from a particular vertex.

To model the types of walks that we need to analyze, let us define a new Markov chain M . The states of M consist of $H \cup \{(h_{u,v,t}, i) \mid h_{u,v,t} \text{ is a path in } G \text{ of length } t \geq 2 \text{ between vertices } u \text{ and } v \text{ in } H \text{ in which all the vertices other than } u \text{ and } v \text{ are in } G - H, 1 \leq i < t\}$. Let P be the transition matrix for Markov chain M . P is defined as follows:

- If $u \in H$, then $P_{u,u} = \frac{1}{2}$
- If $u, v \in H$, $(u, v) \in E(G)$ and $u \neq v$, then $P_{u,v} = \frac{1}{2d}$
- If $h_{u,v,t}$ is a path between two vertices u and v in H of length t that only passes through vertices not in H , then $P_{u,(h_{u,v,t},1)} = \frac{1}{(2d)^t}$, $P_{(h_{u,v,t},i-1),(h_{u,v,t},i)} = 1$ for $1 < i < t$ and $P_{(h_{u,v,t},t-1),v} = 1$.
- All unspecified transition probabilities are 0

These probabilities are defined so that for any two vertices u and v in H , the probability of taking a walk in M from u to v is exactly the same as the probability of taking a walk in G from u to v .

Let π_M be the stationary distribution of M . From the above observation, it follows that for all $u \in H$, $\pi_M(u) = \frac{1}{|G|}$, the same probability that a random walk in G assigns to u . Also, note that for a given u and v in H :

$$\sum_{h_{u,v,t}} \pi_M((h_{u,v,t}, 1)) = \pi_M(u) p_{u,v,t} = \frac{p_{u,v,t}}{|G|} \quad (1)$$

where $p_{u,v,t}$ is the probability of a random walk in G of length t that starts from u and ends at v without passing through any other vertex in H . Also, for each of the paths $h_{u,v,t}$, it is true that $\pi_M((h_{u,v,t}, 1)) = \pi_M((h_{u,v,t}, i))$ for all $1 \leq i < t$ because of how the transition probabilities of M are defined. Therefore:

$$\frac{p_{u,v,t}}{|G|} = \sum_{h_{u,v,t}} \pi_M((h_{u,v,t}, 1)) = \frac{1}{t-1} \sum_{1 \leq i < t} \sum_{h_{u,v,t}} \pi_M((h_{u,v,t}, i)) \quad (2)$$

Summing over all $t > \ell$,

$$\sum_{t > \ell} p_{u,v,t} = |G| \sum_{t > \ell} \frac{1}{t-1} \sum_{1 \leq i < t} \sum_{h_{u,v,t}} \pi_M((h_{u,v,t}, i)) \leq \frac{|G|}{\ell} \sum_{1 \leq i < t} \sum_{h_{u,v,t}} \pi_M((h_{u,v,t}, i)) \quad (3)$$

¹By the length of a walk, we mean the number of steps taken, not the number of edges in G that are followed.

Finally, summing over all u and v in H :

$$\sum_{u,v \in H} \sum_{t > \ell} p_{u,v,t} \leq \frac{|G|}{\ell} \sum_{u,v \in H} \sum_{1 \leq i < t} \pi_M((h_{u,v,t}, i)) \leq \frac{|G|}{\ell} \cdot 1 \quad (4)$$

The last inequality holds because all the $(h_{u,v,t}, i)$ states in the sum are disjoint.

Therefore, if u is chosen uniformly at random from H as the starting vertex of a random walk in G , the expected probability that it takes more than ℓ steps to return to a vertex in H is at most $\frac{\sum_{u \in H} \sum_{v \in H, t > \ell} p_{u,v,t}}{|H|} \leq \frac{|G|}{|H|} \frac{1}{\ell}$.

□

This lemma and its proof is taken from Lemma 4.2 in [GR99].

References

- [GR99] Oded Goldreich and Dana Ron. A sublinear bipartiteness tester for bounded degree graphs. *Combinatorica*, 19(3):335–373, 1999.