

The Cognitive and Neural Development of Face Recognition in Humans

Elinor McKone¹, Kate Crookes¹ & Nancy Kanwisher²

- 1 School of Psychology, Australian National University, Canberra, ACT 0200, Australia
- 2 McGovern Institute for Brain Research and Department of Brain & Cognitive Science, MIT

Author contacts: elinor.mckone@anu.edu.au; ngk@mit.edu

Revised Final Version after comments 24th June 2008

Abstract

Conventional wisdom has long held that face recognition develops very slowly throughout infancy, childhood, and adolescence, with perceptual experience as the primary engine of this development. However, striking new findings from just the last few years have overturned much of this traditional view by demonstrating genetic influences on the face recognition system as well as impressive face discrimination abilities present in newborns and in monkeys who were reared without ever seeing a face. Nevertheless, experience does play a role, for example in narrowing the range of facial subtypes for which discrimination is possible and perhaps also increasing discrimination abilities within that range. Here we first describe the cognitive and neural characteristics of the adult system for face recognition, and then we chart the development of this system over infancy and childhood. This review identifies a fascinating new puzzle to be targeted in future research: all qualitative aspects of adult face recognition measured behaviorally are present very early in development (by 4 yrs of age; all that have been tested are also present in infancy) – yet fMRI and ERP evidence shows very late maturity of face-selective neural responses (with the fusiform face area increasing substantially in volume between age 7 and adulthood).

1. Introduction

One of the most impressive skills of the human visual system is our ability to identify a specific individual from a brief glance at their face, thus distinguishing that individual from hundreds of other people we know, despite the wide variations in the appearance of each face as it changes in viewpoint, lighting, emotional expression, and hairstyle. Though many mysteries remain, important insights have been gleaned over the last two decades about the cognitive and neural mechanisms that enable us to recognize faces. Here we address an even more difficult and fundamental question: how does the machinery of face recognition get wired up during development in the first place?

Our review of the available evidence supports a view of the development of face recognition dramatically different from that suggested by the first studies in the field. Twenty years ago, the standard theory was that core aspects of the ability to discriminate faces were not present until 10 years of age, and their emergence and eventual maturity were determined primarily by experience (Carey & Diamond, 1977; Carey, Diamond et al., 1980). This position has been overturned by recent findings demonstrating striking abilities even in neonates, and by mounting evidence of genetic contributions.

We organize our review by age group. Throughout, we ask how the available data address the following fundamental theoretical questions: (a) what are the inherited genetic contributions to the specification of the adult system for processing facial identity information; (b) what is derived from experience; and (c) how exactly do genes and/or experience work separately or together across the course of development to produce the adult system?

2. The Perception of Face Identity in Adulthood

We begin with a characterization of the end state of development: the cognitive and neural basis of the perception of facial identity in adults. Note that this is a major topic in its own right, with much internal theoretical debate. However, to facilitate our present interest in the *developmental* course of face recognition, we focus on empirical phenomena, especially those that (a) are well-established in adults, and (b) have subsequently been tested in development.

a. Core Behavioral Properties of Face Identity Perception in Adult Humans.

Basic properties of face identification in adults are as follows. Identification is more accurate when faces are upright than when they are inverted (i.e., upside down) on both memory and perceptual tasks, and the inversion decrement is substantially larger for faces than nonface objects (the *disproportionate inversion effect*; Yin, 1969; also (Robbins & Mckone, 2007). Generalization from a single image of a novel face in one viewpoint to an image in another is relatively poor, albeit better from the three-quarter view to front or profile views than between the more distinct profile and front views (the *three-quarter view advantage*; Logie, Baddeley et al., 1987). For familiar faces, performance in memory tasks relies more strongly on inner face regions than on

external regions that include hair; for unfamiliar faces, the pattern is reversed (*inner vs. outer features effects*; (Ellis, Sheperd et al., 1979). Finally, identification of own-race faces is better than identification of other-race faces (the *other-race effect*; (Meissner & Brigham, 2001). Note that the first two properties (i.e., the disproportionate inversion effect and the three-quarter view advantage) derive directly from perceptual processing, but the last two are known to derive at least partly from deliberate task strategies (e.g., reliance on hair for novel faces if distinctive hair is present, (Duchaine & Weidenfeld, 2003) or social and attentional factors (other-race effect, (Bernstein, Young et al., 2007).

Additional experimental findings can be grouped under the heading of phenomena that have motivated the concept of *holistic/configural* processing. Holistic/configural processing is defined (e.g., Tanaka & Farah, 1993; (Maurer, Lewis et al., 2005) as (a) a strong integration at the perceptual level of information from all regions of the face (so that altering one region leads to changes in the percept of other regions), which (b) codes the exact spacing between face features (and more controversially exact feature shape as well; (Yovel & Duchaine, 2006) and, (c) is strongly sensitive to face inversion. Relevant phenomena are as follows. Subjects find it harder to identify one half of a combination face (e.g., top half of George Bush with bottom half of Tony Blair) if the inconsistent other half-face is aligned with the target half rather than misaligned (the *composite effect*; (Young, Hellawell et al., 1987). Subjects are also better able to distinguish which of two face parts (e.g., two noses) appeared in a previously-shown face when these are presented in the context of the whole face than when presented in isolation (the *part-whole effect*; (Tanaka & Farah, 1993); part-choice is also better in the original whole than in a version of the whole face with an alteration in spacing between non-target features (the *part-in-spacing-altered-whole effect*; (Tanaka & Sengco, 1997), a finding consistent with other evidence of strong *sensitivity to spacing changes* (eg., distance between eyes) in upright faces (e.g., (Rhodes, Brake et al., 1993; Mckone, Aitkin et al., 2005). When an upright and inverted version of a face is superimposed in transparency, the upright face is perceived more strongly (*perceptual bias to upright*, (Martini, Mckone et al., 2006). All these holistic effects are specific to upright faces: they are not found for inverted or scrambled faces (Young, Hellawell et al., 1987; Tanaka & Sengco, 1997; Robbins & Mckone, 2003; Martini, Mckone et al., 2006), and are weak or absent for objects, including objects-of-expertise (for review see (Mckone, Kanwisher et al., 2007; Robbins & Mckone, 2007).

Finally, other behavioral phenomena have been taken to indicate coding within a perceptual 'face-space', defined as a multidimensional space in which each individual face is coded as a point by its value on underlying dimensions describing different aspects of facial structure, and for which the 'average' face lies at the centre of the space (Valentine, 1991). These phenomena include: *distinctiveness effects*, in which performance is better for distinctive faces than typical faces on old-new recognition tasks, but the pattern is reversed on face-nonface classification tasks (Valentine & Bruce, 1986) and *adaptation aftereffects*, in which, for example, adaptation to expanded faces make a physically normal face appear contracted (Webster & Maclin, 1999) and adaptation to 'anti-Bill' (the physical opposite of Bill in face space) makes the average face appear like Bill (Leopold, O' Toole et al., 2001).

b. Neurophysiology and fMRI in Adult Monkeys.

Adult monkeys show cortical mechanisms specialized for face perception. Strongly face-selective responses from single neurons (“face cells”) are well established in the temporal lobes of macaques (Desimone, Albright et al., 1984; Foldiak, Xiao et al., 2004), and face-selective cortical regions have been reported in macaques using fMRI (Tsao, Freiwald et al., 2003; Pinski, 2005). Tsao, Freiwald, Tootell, and Livingstone (2006) demonstrated direct correspondence between face-selective fMRI patches and face selectivity of single cells within those patches. Note that the role of “face cells” in supporting the behavioural phenomena described in the previous section is mostly unexplored, with the exceptions that a preponderance of face-selective cells are tuned to upright (Perrett, Mistlin et al., 1988) and that their tuning to facial distortions from the ‘average face’ is consistent with a face-space coding of facial identity (Leopold, Bondar et al., 2006). In development, only basic face-selectivity has been studied.

c. fMRI: Cortical Loci of Face Identity Processing in Adult Humans

Brain imaging in humans reveals three face-selective cortical regions (Figure 1), of which the “fusiform face area” or FFA (Kanwisher, McDermott et al., 1997) is the main one investigated in children. This region, which can be found in essentially every normal adult in a short “localizer” scan (Saxe, Brett et al., 2006), responds more strongly to faces than to letterstrings and textures (Puce, Allison et al., 1996), flowers (McCarthy, Luby et al., 1997), and indeed all other nonface stimuli that have been tested to date, including mixed everyday objects, houses, hands (Kanwisher, McDermott et al., 1997), and objects of expertise (Kanwisher & Yovel, in press).

fMR-adaptation studies show that neural populations in the FFA can discriminate face identity (Rotshtein, Henson et al., 2005), but not facial expression (Winston, Vuilleumier et al., 2003). The FFA is involved in individual discrimination of upright but not inverted faces (Yovel & Kanwisher, 2005; Mazard, Schiltz et al., 2006), and its inversion effect (i.e., higher response to upright than inverted faces) correlates with the behavioral inversion effect (Yovel & Kanwisher, 2005). The FFA also demonstrates holistic processing, specifically a composite effect (Schiltz & Rossion, 2006).

d. Electrophysiological Signatures in Human Adults

A negative-going ERP response peaking about 170 ms after stimulus onset over posterior temporal sites (N170) has been widely replicated to be face-selective (Halgren, Raji et al., 2000; Liu, Harris et al., 2002). This peak is delayed by 10 ms, and is larger in amplitude, for inverted faces relative to upright faces (Bentin, Allison et al., 1996). The N170 also shows identity discrimination (lower response for repeated compared to unrepeated faces), when the faces are upright but not inverted (Jacques & Rossion, 2006; Jacques, D'arripe et al., 2007). An important point relevant to the interpretation of developmental studies is that

the neural source of the N170 is unknown even in adults, and the sources of suggested equivalent components in children and infants could possibly be different again.

3. Data from adult subjects relevant to the roles of experience and genetics

Before considering what developmental studies tell us about the roles of experience and genetics in face recognition, we describe several findings from adults that also bear directly upon these issues.

Clearly, *experience in isolation* can influence face perception. Adults continue to learn new faces throughout life, and this improves perceptual discrimination of these faces: matching the correct face photograph to a degraded security-camera video image is more accurate if the face is familiar than if it is unfamiliar (Burton, Wilson et al., 1999) also see (Bruce, Henderson et al., 2001). Temporary aftereffects from adaptation to distorted faces (e.g., (Webster & Maclin, 1999) also indicate purely experience-based changes in the tuning of perceptual representations of faces. Training effects on ability to discriminate trained and novel faces has also been demonstrated in an adult prosopagnosic (Degutis, Bentin et al., 2007). Interestingly, however, there is no evidence that experience alone produces any fundamental *qualitative* change in face processing either neurally or cognitively: for example, holistic processing, 'face-space' effects, and FFA activation all occur strongly for both familiar faces and unfamiliar faces (Young, Hellawell et al., 1987; Kanwisher, Mc Dermott et al., 1997; Webster & Maclin, 1999; Le Grand, Mondloch et al., 2004; Carbon, Strobach et al., 2007).

Studies of human adults provide two sources of evidence for genetic contributions. Inability to recognize faces in the absence of any known brain injury ('developmental prosopagnosia') often runs in families (Duchaine, Germine et al., 2007; Grueter, Grueter et al., 2007; Kennerknecht, Pluempfe et al., 2008). And, in normal adults, fMRI shows greater similarity in the pattern of activation across the ventral visual stream for monozygotic compared to dizygotic twins, but only for stimulus classes for which an evolutionary origin of the observed selective cortical regions could reasonably be proposed: faces, and places, but not written words or chairs (Polk, Park et al., 2007).

In summary, results from adults tell us that experience can fine-tune face recognition without changing its qualitative properties, and that genes explain some of the variation behaviorally and neurally. Importantly adult studies do not tell us at what developmental stage genes have their influence. In particular, they do not necessarily demonstrate that a face system is *present at birth*. Some genetically pre-determined processes are present at birth (e.g., sucking reflex), but others affect maturational processes later in childhood or adolescence (e.g., puberty).

4. Development: Infancy

In exploring genetic and experience-based contributions to face recognition via infancy studies, several interrelated questions are relevant. First, which abilities, if any, are present *at birth*? Visual abilities present in neonates (or in monkeys deprived of all face input) cannot be derived from experience and so

provide the only method of revealing *genetic influences in isolation* from any visual learning. Second, if babies are born with a face representation, is its purpose merely to draw attention to faces (cf. CONSPEC in (Morton & Johnson, 1991) or to support individuation? Third, how broadly tuned is any such representation: broad enough to cover any primate face, specific to own-species faces, or perhaps even to own-race faces? Finally, which, if any, of the types of effects of experience in early infancy that are found in other perceptual and cognitive domains occur for faces: Improvements with increasing experience? Perceptual narrowing (i.e., destruction of earlier ability)? Critical periods? Studies of these topics published within the last few years have dramatically altered our understanding of infant face recognition.

In a classic result, newborns (median age 9 *minutes*) track an upright 'paddle face' (Figure 2a) further than versions in which the position of the internal blobs is scrambled or inverted (Goren, Sarty et al., 1975; Johnson, Dziurawiec et al., 1991). Although it has been suggested this preference could arise from general visual biases (eg., for stimuli with more elements in the upper visual field; (Simion, Macchi Cassia et al., 2003), preference only for the normal contrast polarity of a (Caucasian) face (Farroni et al., 2005) argues for a level of specificity to face-like structure. Thus, humans are born with some type of innate preference that, at the very least, attracts infants' attention to faces. Note the innate representation supporting face preference may be different from that supporting face individuation in adults (Johnson, 2005); indeed, a finding that neonates track faces in the temporal but not nasal visual field (Simion et al., 1998) suggests a subcortical rather than cortical origin.

Our concern in the present chapter is primarily with the development of face *individuation* ability. This can be measured in infants by looking time measures that assess preference and dishabituation-to-perceived-novelty. A classic finding is that neonates can discriminate their mother from other similar-looking women when less than 4 days old (Pascalis, De Schonen et al., 1995; Bushnell, 2001) although mother recognition in the first 24 hrs may be partially dependent on prenatal familiarity with her voice, (Sai, 2005). More recent data demonstrate even more striking abilities. Three-month olds can recognize the identity of *novel* individuals, with *similar-looking* faces (same sex, age, race), *without hair*, and *across view changes* (Pascalis, De Haan et al., 1998; Kelly, Quinn et al., 2007). Indeed, it has very recently been discovered that *newborns* (<3 days) can perform this task (Turati, Bulf et al., 2008) (see Figure 2b). The newborns moreover discriminated only front to 3/4 view changes and not 3/4 to profile, in a pattern somewhat (although not precisely) similar to the 3/4 view advantage seen in adults. Finally, newborns demonstrate an inversion effect on discrimination, with babies 1-3 days old discriminating same-view faces without hair upright but not inverted (Turati, Macchi Cassia et al., 2006).

The newborn discrimination findings strongly suggest that a face representation, tuned to upright and able to support individual-level representation, is *present at birth*. It seems unlikely that 3 'days' of experience with faces – in fact, a maximum of perhaps 12 hrs of visual experience of any kind (newborns sleep 16 hrs per day plus have their eyes shut during breastfeeding and crying) – would be sufficient for a purely learning-based system to support the level of fine discrimination ability observed.

Even more compelling, however, is a recent behavioral study in monkeys (Sugita, 2008). Japanese macaques were raised by human caregivers wearing masks, giving them no exposure to faces, but otherwise normal visual experience in a complex environment. On their *very first* experience with faces (aged 6-24 months), the monkeys showed a preference to look at static photographs of faces over photographs of objects equally novel in their visual environment (e.g., cars, houses), and discriminated very subtle differences between individual faces (Fig 2c) in a habituation paradigm.

A variety of other infant findings also either directly argue that a representational capacity for differentiating individual face structures is present at birth, or at least do not reject this conclusion. Newborns (<1 week) prefer faces rated by adults as attractive over unattractive faces, when they are upright but not inverted (Slater, Quinn et al., 2000). Regarding holistic processing, Sugita's (2008) monkeys discriminated spacing changes (Figure 2c) with almost no prior experience of faces (they had been exposed to faces only during the short face-preference task), and five-month-old humans discriminate spacing changes small enough to fall within the normal physical range, upright but not inverted (Hayden, Bhatt et al., 2007); also babies 6-8 months old show a composite-like effect where the combination of the inner features of one old face with the outer features of another old face is treated as a new individual, upright but not inverted (Cohen & Cashon, 2001). At 3 months (although not 1 month), human infants falsely recognise the average of four studied faces as 'old', a phenomenon also shown by adults (De Haan, Johnson et al., 2001). Importantly, there are no major behavioral properties of face recognition present in adults that are known *not* to be present in infants; where we have not mentioned properties (e.g., adaptation aftereffects), this is because no relevant data exist, not because infants have been tested and failed to show effects.

Findings of *perceptual narrowing* indicate that a representational capacity for faces that is present at birth (a) can initially be applied to a wide range of faces but that (b) this range gets restricted during the first several months of life to include only the kinds of faces (i.e., species or race) that have been seen in this period. Perceptual narrowing is best known from the domain of language (eg. (Kuhl, Tsao et al., 2003). Infants are born with the ability to discriminate phoneme boundaries from all possible languages in the world (e.g., English and Japanese), but over the first 6-12 months of life lose the ability to discriminate phonemes from non-experienced languages (Japanese for a child from a monolingual English-speaking family), and even extensive exposure as an adult is usually insufficient to regain native-speaker levels of discrimination and reproduction. For faces, five studies have reported and explored properties of perceptual narrowing. In humans, Pascalis, de Haan and Nelson (2002) showed that 6 month old infants could discriminate both human and monkey faces, while 9-month-olds and adults could discriminate only human faces. Kelly et al. (2007) reported that Caucasian babies from the north of England, with high exposure to Caucasians but essentially no exposure to African or Asian faces, could recognize individuals (across view change) from all three races at 3 months of age. At 6 months, Caucasian babies could no longer individuate African faces; at 9 months they had additionally lost the ability to individuate Asians. The Sugita (2008) study described earlier reported that, on first exposure to faces, the monkeys could not only discriminate individual monkey faces (other macaques),

but could also make extremely fine discriminations amongst human faces (Figure 2c). Following 1 month of exposure to a single face type (either human or monkey, involving live interaction for least 2 hrs per day), Sugita's monkeys lost the ability to discriminate individuals of the non-experienced species. Re-learning was also difficult: monkeys initially exposed only to humans failed to discriminate monkey faces even after subsequently sharing a cage with 10 other monkeys for 11 months. (Note, however, that there is some evidence of flexibility in humans into middle childhood: Korean children adopted to Caucasian Francophone countries at age 3-9 yrs showed, as adults, better recognition memory for Caucasian faces than Korean faces; (Sangrigoli, 2005). During human infancy, perceptual narrowing can be avoided by deliberate exposure to face types that the infant would not naturally see, with regular exposure to monkey faces beginning at 6 months leading to retained ability to discriminate monkey faces at 9 months (Pascalis, Scott et al., 2005). Perceptual narrowing for faces also has an interesting possible link with narrowing for language. Lewkowicz and Ghazanfar (2006) reported that human infants could make cross-modality matches of a monkey vocalization to a picture of a monkey face making that particular sound at 4 and 6 months, but that this ability was lost at 8 and 10 months.

Importantly, the perceptual narrowing effects for faces described above indicate only a *destructive* effect of experience across infancy (i.e., *loss* of initial ability with other-species and other-races). In the domain of language, loss of phonetic discrimination ability within nonexperienced languages has been shown to co-occur with an improvement of phonetic discriminability within the experienced language (Kuhl et al., 2006). Thus, perceptual narrowing for faces might similarly include enhanced ability to discriminate experienced face subtypes: that is, discrimination for own-species own-race faces might start crude and *improve* with practice. Potentially consistent with this prediction, Humphreys and Johnson (2006) showed the physical difference between faces required to produce novelty preference was smaller in 7 month olds than 4 month olds, indicating that the older babies could either make finer perceptual discriminations, or keep these in memory longer across the 1-5 item test delay. Neural systems present at birth are often associated with a *critical* (or *sensitive*) *period* (Sengpiel, 2007), requiring environmental input of the appropriate stimulus type within a specified period after birth to avoid being taken over for other purposes. In a classic example, cats are born with cells tuned to all line orientations, but if raised in an environment containing only vertical lines they lose horizontal-responsive cells and a corresponding lack of behavioral sensitivity to horizontal lines. For faces, Le Grand and colleagues report evidence consistent with a critical period for one important aspect of face perception, holistic processing. Congenital cataract patients, specifically people born with dense cataracts disrupting all pattern vision who had the cataracts removed at 2-28 months of age, were tested at ages ranging between 9 years and adulthood. Despite their many years of post-cataract exposure to faces, patients who had had early bilateral cataracts showed no composite effect for faces (Le Grand, Mondloch et al., 2004). Also, patients who had had right-eye-only or bilateral cataracts – which produce a deficit of input to the right hemisphere due to the wiring of the infant visual system – showed a later deficit in processing spacing information in faces, while patients who had had left-eye-only cataracts did not

(Le Grand, Mondloch et al., 2003), a pattern consistent with the normal role of the right hemisphere in holistic processing (Rossion, Dricot et al., 2000). Interestingly, there does not appear to be a critical period for the ability to discriminate faces per se. Anecdotally, the Canadian cataract patients are not functionally prosopagnosic (Daphne Maurer, pers comm), for example reporting even being able to recognize other-race students when teaching English in Korea (Rachel Robbins, pers comm). Formal testing shows good ability to match novel faces (without view change) both in these patients (Geldart, Mondloch et al., 2002) and in an Indian woman whose congenital cataracts were not removed until 12 years of age (Ostrovsky, Andalman et al., 2006). Also, lack of visual experience with faces for the first 6-24 months in Sugita's (2008) monkeys did not destroy discrimination ability. The reason why a requirement for early visual input exists for holistic processing but not face discrimination remains to be resolved. One possibly relevant observation is that holistic processing could perhaps have a particular role in cross-view recognition (Mckone, 2008), and the Canadian cataract patients have a specific problem with recognition of once-seen faces across view changes (Geldart, Mondloch et al., 2002; note the Indian patient and Sugita's monkeys were tested on same-view faces only).

The behavioral findings reviewed above – demonstrating abilities present at birth, perceptual narrowing and critical periods – are all consistent with a genetically determined “innate” contribution to infant face recognition. In particular, they argue for an innate contribution to face *individuation*.

Neurally, face individuation in adults is associated with cortical rather than subcortical function. What is the evidence regarding *cortical* face-processing function in infants? There are few available studies, and none in neonates. Results do, however, demonstrate face-selectivity and inversion effects. In infant macaques, Rodman et al (Rodman, Scalaidhe et al., 1993) found that the response magnitude of single units in inferotemporal cortex was lower overall than in adults, but selectivity for form including face selectivity was present at the youngest ages tested, within 2 months of birth. In humans, a PET study of 2.5-month-olds is somewhat suggestive of face-selective activation in the fusiform gyrus (and other cortical regions), although the infants were not neurologically normal, the statistical threshold was extremely lenient ($p < .05$ uncorrected), and the contrast (faces versus blinking diodes) confounds selectivity for faces with responses to visual shape information (Tzourio-Mazoyer, De Schonen et al., 2002). Using ERPs, human 3-month-olds exhibit an “N290” component that has larger amplitude for human compared to monkey faces in the right hemisphere only (Halit, De Haan et al., 2003) although adult N170 shows the opposite pattern. At 12 months of age, this N290 was higher in amplitude for inverted than upright faces, only for human, not monkey faces (like the adult N170). Although the same study reported that this sensitivity to inversion was not found in 3-month-olds, another analysis of the same data using a different method (Johnson, Griffin et al., 2005) did claim to find such inversion sensitivity. Further, other ERP components (the P400 and the P1) do show inversion effects at 3 months, the youngest age tested (Halit, De Haan et al., 2003). Similarly, near infrared spectroscopy (NIRs) responses in 5-8-month-old infants are stronger for upright than inverted faces over the right hemisphere only (Otsuka, Nakato et al., 2007); note the cortical source of this effect was most likely the STS). Overall, the available neural evidence from infants is consistent with the existence of

cortical machinery for processing faces within a few months after birth, and there is no evidence to suggest this is not present earlier.

Taking all findings together, we conclude that infants are born with a rich capacity to represent the structure of upright faces which supports face discrimination, rather than merely drawing attention to faces. Results further show that this representation interacts with experience during infancy in particular ways. A probable critical period suggests holistic processing is 'experience-expectant' (i.e., early environmental input is required for its maintenance). Perceptual narrowing shows early experience restricts the range of faces that can be accommodated: that is, an initial representation of faces is sufficiently broadly tuned to support individuation of all face types including those of other primates, and experience with one subtype of face (own-species, own-race) *removes* this initial ability with other face types (other species, other races), at the same time that it possibly improves perceptual tuning for faces of the experienced subtype. Regarding neural origin of face discrimination in infants, there is evidence of relevant cortical representation by mid-infancy, but no data are available regarding whether the discrimination ability present *at birth* is supported by cortical as opposed to subcortical representations.

5. Development: Four year olds to adults

In understanding the interaction of genetic inheritance and learning, investigation of the developmental trajectory of face processing in childhood through adulthood can also be informative. When no change is found in a given behavioral or neural measure of face perception in this period, that argues against extended maturation or learning as necessary for the construction of the adult system. If instead protracted development is observed, this could reflect learning (as often assumed), though crucially it could also reflect biological maturation (Carey, Diamond & Woods, 1980), or an interaction of genetic and experiential factors.

a. Behavioral Measures of Face Identity Perception

For children 4-5 years and older, it is possible, with care, to adapt adult behavioral paradigms directly, and thus to compare child performance with adult performance on exactly the same tasks. For each phenomenon established in adults, two empirical questions are of interest. First, is there some age below which children simply do not show that phenomenon at all? (i.e., is there *qualitative* change with age?). Second, regarding any phenomena that are observed, when are full maturity levels reached? (i.e., is there *quantitative* change with age).

We consider qualitative change first. Early behavioral research appeared to suggest that core perceptual processes involved in face identification did not emerge at all until quite late in development (e.g., 10 years for holistic processing, (Carey & Diamond, 1977; Carey, Diamond et al., 1980). Unfortunately, researchers in the face neuroscience literature (e.g., (Gathers, Bhatt et al., 2004; Aylward, Park et al., 2005; Golarai, Ghahemani et al., 2007;

Scherf, Behrmann et al., 2007) commonly emphasize only these few early findings, which give an inaccurate representation of the current state of knowledge. In fact, research in the last 15 years has clearly established that *all* standard adult face recognition effects are present in young children. (Indeed, Section 3 showed all phenomena tested – including inversion effects – were present in infancy.)

In child-age studies using adult tasks, every key adult property of face recognition investigated has been obtained at the youngest age tested. With respect to holistic processing, these results include the inversion effect on short- and long-term recognition memory (3 y.o. Sangrigoli & de Schonen, 2004; 4 y.o. Carey, 1981; 5-6 y.o. Brace, Hole et al., 2001; 7 y.o. Flin, 1985; the composite effect (4 y.o. De Heering, Houthuys et al., 2007; 6 y.o. Carey & Diamond, 1994; 6 y.o. Mondloch, Pathman, et al., 2007), the part-whole effect for upright but not inverted faces (4 y.o. Pellicano & Rhodes, 2003; 6 y.o. (Tanaka, Kay et al., 1998), the part-in-spacing-changed-whole effect for upright but not inverted faces (4 y.o. (Pellicano, Rhodes et al., 2006) sensitivity to exact spacing between facial features (4 y.o. McKone & Boyer, 2006; 4 y.o. (Pellicano, Rhodes et al., 2006), the perceptual bias to upright in superimposed faces (8 y.o. (Donnelly, Hadwin et al., 2003) and the internal-over-external features advantage for familiar face identification (5-6 y.o. (Wilson, Blades et al., 2007). Regarding face space coding, results include distinctiveness effects on perception at 4 years (Mckone & Boyer, 2006) and on memory at 6-7 yrs (Gilchrist & Mckone, 2003), an other-race disadvantage on recognition memory at 3 years (Sangrigoli & De Schonen, 2004) and a recent conference report of adaptation aftereffects in 4-5 yr olds (Jeffrey & Rhodes, 2008). Where early studies did not show effects, this has generally been established to have arisen from methodological problems, the most common one being floor effects on the task in young children (e.g., see Carey et al., 1980 vs. Carey, 1981; or Johnston & Ellis, 1995 vs. (Gilchrist & Mckone, 2003). Another case of note is the early suggestion that children could not perform face identification *at all* in the presence of distracting paraphernalia (Carey & Diamond, 1977); this finding was overturned (Lundy, Jackson et al., 2001), by simply making the faces larger. (Also note that even adults are sometimes strongly distracted by paraphernalia, (Simons & Levin, 1998). In summary, it is clear that there is no qualitative change in face perception beyond 4-5 years of age; quite possibly, there is none beyond infancy.

The question of whether quantitative change occurs is more difficult to answer. Certainly, performance on just about any experimental task involving faces improves very substantially across childhood and well into adolescence (see Fig 3a,b). The crucial issue is how much of this development reflects development in *face perception* (e.g., in holistic processing, or in the fine tuning of face-space), and how much reflects development in other general cognitive factors that are known to improve substantially across this age range and would affect task performance whatever the stimuli (e.g., explicit memory ability, ability to concentrate on the task or instruction). A common bias of face researchers is to assume, given data showing increasing memory for faces with age (e.g., Figure 3a), that it is face perception that is changing, and that the task type – explicit memory – is irrelevant; yet, an implicit memory researcher looking at the same set of data would likely conclude ‘explicit memory’ is developing and presume the particular stimulus type – faces – is irrelevant.

Various attempts have been made to overcome the limitations of simply tracking age-related improvement in raw performance. To our minds, however, none of these are methodologically satisfactory, and none produce a clear conclusion regarding whether face perception per se does, or does not, improve between early childhood and adulthood. One approach is to compare *two* conditions across development, for example asking whether the size of the difference between upright and inverted (or typical and distinctive, etc) changes with age (e.g., (Carey & Diamond, 1980; Johnston & Ellis, 1995). The results of almost all such studies, however, are confounded with overall 'baseline' changes across age groups, such that (a) when room to show effects is potentially compressed by approaching floor in young children, but is not restricted (i.e., no ceiling effect) in adults, results seem to suggest quantitative *increases* in the effect of interest with age (Fig 3a), but that (b) when room to show effects is restricted by approaching ceiling in adults, but is not restricted in young children (i.e., no floor effects on accuracy, or alternatively use of a reaction time measure), results seem to show quantitative *decreases* with age (Fig 3b). Taking seriously the results of the first type of study as showing quantitative development in face perception (as is commonly done), requires also taking seriously the results of the second type of study— apparently leading to the conclusion that face perception gets consistently *worse* between early childhood and adulthood! A further requirement for valid comparison of rates of development for two stimulus types is that performance be equated for the two types in one or other end-point age group. This is commonly not done. As one example, the Mondloch et al. (2002) finding that sensitivity to feature changes reaches adult levels earlier than spacing changes can be attributed (Mckone & Boyer, 2006) simply to the fact that the features changes were easier in adults (that is, performance on an easier stimulus set reaches adult levels before performance on a more difficult stimulus set). Another general issue in studies comparing faces versus objects (e.g., in rate of development, Golarai et al., 2007; or size of inversion effects, (Carey & Diamond, 1977; Teunisse & De Gelder, 2003; Aylward, Park et al., 2005), is that, in addition to producing very mixed results, the object classes tested to date (houses, scenes, sculptures, shoes) have not been well matched to faces on basic parameters, such as not sharing a first-order configuration (houses, scenes), or not being natural objects (sculptures, shoes).

Overall, we conclude that current behavioral evidence demonstrates qualitatively adult-like processing of faces in young children, but does not resolve whether processing is quantitatively mature. We note, however, that at least some evidence suggests a conclusion likely to be surprising to many readers, namely that even *quantitative* maturity might be reached by early childhood. The three studies that appear to have the most suitable methodology – in which baselines were matched across age groups (Carey, 1981; Gilchrist & Mckone, 2003), or restriction of range problems were otherwise avoided (Mondloch, Pathman et al., 2007)– all indicate no change in holistic processing (inversion effect, Carey, 1981; composite effect, Mondloch, Pathman et al., 2007; spacing sensitivity, Gilchrist & McKone, 2003; or distinctiveness effects (Gilchrist & Mckone, 2003) between early childhood (4-6 years) and adulthood (Fig 3c).

b. Neural Measures of Face Identity Processing (FFA and N170)

As with behavioral studies, we discuss results of neuroimaging and ERP studies in children with respect to two questions: qualitative development, and quantitative development.

Three studies have used fMRI to scan children age 5 to adult on face and object tasks, enabling these studies to track the existence and size of face-selective regions of cortex. (A fourth study will not be discussed here because it used such liberal criteria to define “FFAs” that the regions so identified were clearly not face-selective even in adults; see Figure 1 d-f in that study, (Gathers, Bhatt et al., 2004). Considering *qualitative* effects, evidence of a face-selective FFA has been found in most children at the youngest ages tested. Although no FFA was revealed in young children by group analyses (in which all subjects are aligned in a common space; 5-8 y.o., Scherf, Behrmann et al., 2007; 8-10 y.o., (Aylward, Park et al., 2005), in the two studies reporting individual-subject analyses, Scherf et al found an FFA in 80% of the children in 5-8 year olds (albeit at a very liberal statistical threshold), and Golarai et al (2007) found an FFA in 85% of children in their 7-11 year old group (using a more standard statistical threshold). One study (Passarotti, Smith et al., 2007) also reported an inversion effect (higher response to inverted than upright faces) in the region of the right (but not left) FFA in children 8-11 years of age (and an effect in the opposite direction in adults). Regarding ERPs, young children (like infants) show both face-selective responses and inversion effects upon these (see Figures 5 and 6; Taylor, Batty et al., 2004). These fMRI and ERP findings in children add to the infant data to confirm that at least some form of face-specific neural machinery is established early.

Quantitatively, neural machinery involved in face perception demonstrates substantial changes in face-selective neural responses continuing late into development. In all three fMRI studies, the FFA increases markedly in volume between childhood and adulthood (Aylward, Park et al., 2005; Golarai, Ghahemani et al., 2007; Scherf, Behrmann et al., 2007), even though total brain volume does not change substantially after age 5. These studies clearly show that the rFFA is still changing late in life, certainly after age 7 and in some studies much later.

Comparing fMRI data across children and adults is fraught with potential pitfalls. Children move more in the scanner, and are less able to maintain attention on a task. These or other differences between children and adults could in principle explain the change in volume of the rFFA. However, notably, control areas identified in the same scanning sessions do not change with age. For example object-responsive regions and the scene-selective “parahippocampal place area” in the right hemisphere or rPPA (Epstein & Kanwisher, 1998) did not change in volume from childhood to adulthood (Golarai, Ghahemani et al., 2007; Scherf, Behrmann et al., 2007) although somewhat surprisingly Golarai et al found that the lPPA did increase in volume with age. These findings are reassuring that the changes in the rFFA with age are not due to across-the-board changes in the ability to extract good functional data from young children.

Golarai et al (2007) asked how changes in the rFFA relate to changes in behavioral face recognition over development. Right FFA size was correlated (separately in children and adolescents but not in adults) with face recognition memory, but not with place or object memory. Conversely, lPPA size was correlated (in all age groups independently) with place memory but not with

object or face memory. This double dissociation of behavioral correlations clearly associates the rFFA with changes in face recognition measured behaviorally.

ERP findings are consistent with the evidence from fMRI that the cortical regions involved in face recognition continue to change well into the teenage years. Face-related ERPs show gradual changes in scalp distribution, latency, and amplitude into the mid-teen years (Figure 5 and 6). Both the early P1 component and the later N170 component show gradual decreases in latency from age 4 to adulthood. Regarding neural inversion effects, late developmental changes are found with both fMRI and ERP (see Figure 5), including a reversal of the direction of the inversion effect between children and adults in both methods (Taylor, Batty et al., 2004; Passarotti, Smith et al., 2007). Future research might best approach this question by measuring not just mean responses to upright versus inverted faces, but instead using identity-specific adaptation to ask when the better discrimination of upright than inverted faces seen in adulthood (Yovel & Kanwisher, 2005; Mazard, Schiltz et al., 2006).

c. Comparing Development for Behavioral and Neural Measures

Taking the findings from the 4-to-adult range together with the infant literature, we can draw the following conclusions. First, the results regarding *qualitatively* adult-like face processing appear to agree well across behavioral and neural measures; that is, just as all behavioral face recognition effects have been obtained in the youngest age groups tested, face-selective neural machinery as revealed by fMRI, ERPs, NIRs and single-cell recording has also been found in the youngest children and infants tested. Nonetheless, fMRI data are not available for children younger than 5-8 (pooled together), and the ERP studies in infants and children often go in opposite directions from adults. For example, and the inversion effect on the N170 switches polarity between childhood and adulthood, as shown in Figure 6, despite maintaining the same polarity in behavior.

Second, the evidence for quantitative development is less clear. It may be that the improvements with age on behavioral tasks do reflect ongoing development of face perception itself and, if so, this could agree neatly with the increasing size of the FFA. As we have noted, however, findings such as those shown in Figures 3b and 3c suggest that behavioral face perception may be fully mature early, and that ongoing behavioral improvements with age reflect changes in other, more general, cognitive factors. This view would produce an apparent discrepancy – behavioral maturity arising well before maturity of relevant cortical regions – that would need to be resolved. If this is the case, two ideas might worth exploring. It may be that the measured size of the FFA in children is affected by top-down strategic processing which (for some unknown reason) affects faces and not objects. Another possibility is that the FFA might play some role in the long-term storage of individual faces (e.g., it shows repetition priming, (Pourtois, Schwartz et al., 2005; Williams, Berberovic et al., 2007) and that the increased size of the FFA could arise simply because people continue to learn faces across life; this idea would have to propose that the number of new faces learned is much greater than the number of new objects.

6. Conclusion

For decades, conventional wisdom has held that face recognition arises very slowly in development, and that experience is the primary engine of this development. The new evidence reviewed here refutes this hypothesis. Impressive face recognition abilities are present within a few days of birth, and are present in monkeys who have never seen faces before. Some form of inherited genetic influence is also indicated by Polk's imaging study of twins, and by the fact that developmental prosopagnosia can run in families. Qualitatively, behavioral findings indicate establishment of *all* adult-like face recognition effects by 4 years at the latest, and in infancy wherever tested; the striking breadth of this evidence is summarised in Figure 7. The available evidence also indicates early initial establishment of face-selective neural machinery at the cortical level; again see Figure 7. It is not, however, that experience plays no role in development. Perceptual narrowing of the range of facial subtypes for which discrimination is possible reveals a destructive role for experience. Further, there is a requirement for early-infancy input (consistent with a critical period) for the development of holistic face processing but (mysteriously) not face discrimination.

Three major questions remain for future research. First, it will be critical to determine whether face perception *per se* improves quantitatively after age 4, or whether instead improvement in performance after this age reflects improvement in domain-general mechanisms. Second, if face perception itself does improve quantitatively after age 4, what role does experience play in this improvement? A final critical challenge will be to understand the relationship between cognitive and neural development, especially the substantial increase in the size of the FFA.

Acknowledgements

We thank Mark Johnson for useful comments and Bettiann McKay for help with the manuscript. Preparation of this chapter was supported by NEI grant 13455 and a grant from the Ellison Foundation to NK and by grant DP0450636 from the Australian Research Council to EM.

Figure Legends

Figure 1. Cortical regions selectively engaged in face perception and the development of one of these regions (the FFA) from childhood to adulthood. A. Adults: Face selective activation (faces > objects, $p < .0001$) on an inflated brain of one adult subject, shown from lateral and ventral views of the right and left hemispheres. Three face-selective regions are shown: the FFA in the fusiform gyrus along the ventral part of the brain, the OFA in the lateral occipital area and the fSTS in the posterior region of the superior temporal sulcus. For studies of face *identification* (rather than expression, etc), the FFA and OFA are of greatest interest.

Figure 2. Face perception without experience. (A) Newborn humans (<1 hr old) track the 'paddle face' on the left further than the scrambled version (Morton & Johnson, 1991); (B) Newborn humans (<3 days) look longer at the novel than habituated face, indicating recognition of face identity even across view change (Turati, Bulf et al., 2008); (C) Japanese macaques raised with no exposure to faces can, on first testing, discriminate very subtle differences between individual monkey faces (including differences both in shape and in spacing of internal features), and can also do this for human faces (Sugita, 2008).

Figure 3. Behavioral face recognition effects in the preschooler to adult age range. A basic finding is of overall improvement with age – higher accuracy or lower reaction time; note that in C, the left and middle plots show studies where the researchers deliberately removed this trend by using smaller learning set sizes in younger children. Our major point is that apparent developmental trends in the strength of core effects (size of inversion effect, size of composite effect, ability to represent recently-seen faces in implicit memory, etc) depend on whether, and how, room to show effects is potentially restricted.

Figure 4. Developmental data from Golarai et al (Golarai, Ghahemani et al., 2007): Mean volume across subjects in each age group of individually-defined left (A) and right (B) FFA, (C) anatomically-defined right mid-fusiform gyrus, (D) functionally-defined right LOC, and functionally-defined face-selective right STS (E) and right place-selective PPA (F). Red bars indicated values in subsets of subjects matched for BOLD-related confounds.

Figure 5. ERPs from right posterior temporal scalp locations in response to face stimuli, separately for each age group, from Taylor et al (Taylor, Batty et al., 2004).

Figure 6. Mean N170 latency (left) and amplitude (right) for upright and inverted faces as a function of age, from Taylor et al (Taylor, Batty et al., 2004).

Figure 7. For each property of face processing, we indicate for each age group whether that property is *qualitatively* present (☺), debatable (?), not present (X), or not yet tested (grey). Deprived = monkeys deprived of face input from birth. Note: All references can be found in text except: Inversion effect on spacing

sensitivity aged 6years-adult (Mondloch, Le Grand et al., 2002), Adaptation
aftereffect aged 9 years – adult (Pellicano, Jeffrey et al., 2007).

Bibliography

- Aylward, E. H., Park, J. E., Field, K. M., Parsons, A. C., Richards, T. L., Cramer, S. C., & Meltzoff, A. N. (2005). Brain activation during face perception: Evidence of a developmental change. *Journal of Cognitive Neuroscience*, 17(2), 308-319.
- Bentin, S., Allison, T., Puce, A., Perez, E., & Mc Carthy, G. (1996). Electrophysiological studies of face perception in humans. *Journal of Cognitive Neuroscience*, 8(6), 551-565.
- Bernstein, M. J., Young, S. G., & Hugenberg, K. (2007). The cross-category effect: Mere social categorization is sufficient to elicit an own-group bias in face recognition. *Psychological Science*, 8, 706-712.
- Bruce, V., Henderson, Z., Newman, C., & Burton, M. A. (2001). Matching identities of familiar and unfamiliar faces caught on CCTV images. *Journal of Experimental Psychology: Applied*, 7(3), 207-218.
- Burton, A. M., Wilson, S., Cowan, M., & Bruce, V. (1999). Face recognition in poor-quality video: Evidence from security surveillance. *Psychological Science*, 10(3), 243-248.
- Bushnell, I. W. R. (2001). Mother's face recognition in newborn infants: Learning and memory. *Infant and Child Development*, 10, 67-74.
- Carbon, C. C., Strobach, T., Langton, S., Harsányi, G., Leder, H., & Kovács, G. (2007). Adaptation effects of highly familiar faces: immediate and long lasting. *Memory and Cognition*, 35(8), 1966-1976.
- Carey, S. (1981). The development of face perception. In G. Davies, H. D. Ellis & J. Shepherd (Eds.), *Perceiving and remembering faces* (pp. 9-38). New York: Academic Press Inc.
- Carey, S., & Diamond, R. (1977). From piecemeal to configurational representation of faces. *Science*, 195, 312-314.
- Carey, S., & Diamond, R. (1980). Maturation determination of the development course of face encoding. In D. Caplan (Ed.), *Biological studies of mental processes*. Cambridge, MA: MIT press.

- Carey, S., Diamond, R., & Woods, B. (1980). Development of face recognition - A maturational component? *Developmental Psychology*, 16(4), 257-269.
- Cohen, L. B., & Cashon, C. H. (2001). Do 7-month-old infants process independent features of facial configurations? *Infant and Child Development*, 10, 83-92.
- de Haan, M., Johnson, M. H., Maurer, D., & Perrett, D. I. (2001). Recognition of individual faces and average face prototypes by 1- and 3-month-old infants. *Cognitive Development*, 16, 659-678.
- Degutis, J. M., Bentin, S., Robertson, L. C., & D'Esposito, M. (2007). Functional plasticity in ventral temporal cortex following cognitive rehabilitation of a congenital prosopagnosic. *Journal of Cognitive Neuroscience*, 19(11), 1790-1802.
- Desimone, R., Albright, T. D., Gross, C. G., & Bruce, C. (1984). Stimulus-selective properties of inferior temporal neurons in the macaque. *Journal of Neuroscience*, 4(8), 2051-2062.
- Donnelly, N., Hadwin, J. A., Cave, K., & Stevenage, S. (2003). Perceptual dominance of oriented faces mirrors the distribution of orientation tuning in inferotemporal neurons. *Cognitive Brain Research*, 17, 771-780.
- Duchaine, B., Germine, L., & Nakayama, K. (2007). Family resemblance: Ten family members with prosopagnosia and within-class object agnosia. *Cognitive Neuropsychology*, 24(4), 419-430.
- Duchaine, B. C., & Weidenfeld, A. (2003). An evaluation of two commonly used tests of unfamiliar face recognition. *Neuropsychologia*, 41(6), 713-720.
- Ellis, H. D., Sheperd, J. W., & Davies, G. M. (1979). Identification of familiar and unfamiliar faces from internal and external features: Some implications for theories of face recognition. *Perception*, 8(4), 431-439.
- Epstein, R., & Kanwisher, N. (1998). A cortical representation of the local visual environment. *Nature*, 392(6676), 598-601.
- Foldiak, P., Xiao, D., Keysers, C., Edwards, R., & Perrett, D. I. (2004). Rapid serial visual presentation for the determination of neural selectivity in area STSa. *Progress In Brain Research*, 144, 107-116.

- Gathers, A. D., Bhatt, R. S., Corbly, C. R., Farley, A. B., & Joseph, J. E. (2004). Developmental shifts in cortical loci for face and object recognition. *NeuroReport*, 15(10), 1549-1553.
- Geldart, S., Mondloch, C. J., Maurer, D., de Schonen, S., & Brent, H. P. (2002). The effect of early visual deprivation on the development of face processing. *Developmental Science*, 5(4), 490-501.
- Gilchrist, A., & McKone, E. (2003). Early maturity of face processing in children: Local and relational distinctiveness effects in 7-year-olds. *Visual Cognition*, 10(7), 769-793.
- Golarai, G., Ghashemi, D. G., Whitfield-Gabrieli, S., Reiss, A., Eberhardt, J. L., Gabrieli, J. D., & Grill-Spector, K. (2007). Differential development of high-level visual cortex correlates with category-specific recognition memory. *Nature Neuroscience*, 10, 512-522.
- Goren, C. C., Sarty, M., & Wu, P. Y. K. (1975). Visual following and pattern discrimination of face-like stimuli by newborn infants. *Pediatrics*, 56(4), 544-549.
- Grueter, M., Grueter, T., Bell, V., Horst, J., Laskowski, W., Sperling, K., Halligan, P. W., Ellis, H. D., & Kennerknecht, I. (2007). Hereditary prosopagnosia: the first case series. *Cortex*, 43(6), 734-749.
- Halgren, E., Raij, T., Marinkovic, K., Jousmaki, V., & Hari, R. (2000). Cognitive response profile of the human fusiform face area as determined by MEG. *Cerebral Cortex* 10(1), 69-81.
- Halit, H., de Haan, M., & Johnson, M. H. (2003). Cortical specialisation for face processing: face-sensitive event-related potential components in 3- and 12-month-old infants. *Neuroimage*, 19(3), 1180-1193.
- Hayden, A., Bhatt, R. S., Reed, A., Corbly, C. R., & Joseph, J. E. (2007). The development of expert face processing: Are infants sensitive to normal differences in second-order relational information? *Journal of Experimental Child Psychology*, 97, 85-98.
- Jacques, C., d'Arripe, O., & Rossion, B. (2007). The time course of the inversion effect during individual face discrimination. *Journal of Vision*, 7(8), 1-9.

- Jacques, C., & Rossion, B. (2006). The speed of individual face categorization. *Psychological Science*, 17(6), 485-492.
- Jeffrey, L., & Rhodes, G. (2008). Aftereffects reveal enhanced face-coding plasticity in young children, *Vision Sciences Society*). Naples, Florida.
- Johnson, M. H., Dziurawiec, S., Ellis, H., & Morton, J. (1991). Newborns' preferential tracking of face-like stimuli and its subsequent decline. *Cognition*, 40(1-2), 1-19.
- Johnson, M. H., Griffin, R., Csibra, G., Halit, H., Farroni, T., de Haan, M., Tucker, L. A., Baron-Cohen, S., & Richards, J. (2005). The emergence of the social brain network: evidence from typical and atypical development. *Development and Psychopathology*, 17(3), 599-619.
- Johnston, R. A., & Ellis, H. D. (1995). Age effects in the processing of typical and distinctive faces. *The Quarterly Journal of Experimental Psychology*, 48A(2), 447-465.
- Kanwisher, N., Mc Dermott, J., & Chun, M. M. (1997). The fusiform face area: A module in human extrastriate cortex specialized for face perception. *Journal of Neuroscience*, 17(11), 4302-4311.
- Kanwisher, N., & Yovel, G. (in press). Cortical Specialization for Face Perception in Humans. In J. T. Cacioppo & G. G. Berntson (Eds.), *Handbook of Neuroscience for the Behavioral Sciences*). Hoboken, NJ: J. Wiley and Sons.
- Kanwisher, N. G., McDermott, J., & Chun, M. M. (1997). The fusiform face area: A module in human extrastriate cortex specialized for face perception. *Journal of Neuroscience*, 17(11), 4302-4311.
- Kelly, D. J., Quinn, P. C., Slater, A., Lee, K., Ge, L., & Pascalis, O. (2007). The other-race effect develops during infancy: Evidence of perceptual narrowing. *Psychological Science*, 18(12), 1084-1089.
- Kennerknecht, I., Pluempke, N., & Welling, B. (2008). Congenital prosopagnosia--a common hereditary cognitive dysfunction in humans. *Frontiers in Bioscience*, 1(13), 3150-3158.

- Kuhl, P. K., Tsao, F.-M., & Liu, H.-M. (2003). Foreign-language experience in infancy: Effects of short-term exposure and social interaction on phonetic learning. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 9096-9101.
- Le Grand, R., Mondloch, C. J., Maurer, D., & Brent, H. P. (2003). Expert face processing requires visual input to the right hemisphere during infancy. *Nature Neuroscience*, 6(10), 1108-1112.
- Le Grand, R., Mondloch, C. J., Maurer, D., & Brent, H. P. (2004). Impairment in holistic face processing following early visual deprivation. *Psychological Science*, 15(11), 762-768.
- Leopold, D. A., Bondar, I. V., & Giese, M. A. (2006). Norm-based face encoding by single neurons in the monkey inferotemporal cortex. *Nature*, 442(7102), 572-575.
- Leopold, D. A., O' Toole, A. J., Vetter, T., & Blanz, V. (2001). Prototype-referenced shape encoding revealed by high-level aftereffects. *Nature Neuroscience*, 4(1), 89-94.
- Liu, J., Harris, A., & Kanwisher, N. (2002). Stages of processing in face perception: an MEG study. *Nature Neuroscience*, 5(9), 910-916.
- Logie, R. H., Baddeley, A. D., & Woodhead, M. M. (1987). Face recognition, pose and ecological validity. *Applied Cognitive Psychology*, 1, 53-69.
- Lundy, B. L., Jackson, J. W., & Haaf, R. A. (2001). Stimulus properties, attentional limitations, and young children's face recognition. *Perceptual and Motor Skills*, 92, 919-929.
- Martini, P., McKone, E., & Nakayama, K. (2006). Orientation tuning of human face processing estimated by contrast matching on transparency displays. *Vision Research*, 46(13), 2102-2109.
- Maurer, D., Lewis, T. L., & Mondloch, C. J. (2005). Missing sights: consequences for visual cognitive development. *Trends in Cognitive Science*, 9(3), 144-151.

- Mazard, A., Schiltz, C., & Rossion, B. (2006). Recovery from adaptation to facial identity is larger for upright than inverted faces in the human occipito-temporal cortex. *Neuropsychologia*, 44(6), 912-922.
- McCarthy, G., Luby, M., Gore, J., & Goldman-Rakic, P. (1997). Infrequent events transiently activate human prefrontal and parietal cortex as measured by functional MRI. *Journal of Neurophysiology*, 77(3), 1630-1634.
- McKone, E. (2008). Configural processing and face viewpoint. *Journal of Experimental psychology. Human perception and performance*, 34(2), 310-327.
- McKone, E., Aitkin, A., & Edwards, M. (2005). Categorical and coordinate relations in faces, or Fechner's law and face space instead? *Journal of Experimental Psychology: Human Perception and Performance*, 31(6), 1181-1198.
- McKone, E., & Boyer, B. L. (2006). Sensitivity of 4-year-olds to featural and second-order relational changes in face distinctiveness. *Journal of Experimental Child Psychology*, 94(2), 134-162.
- McKone, E., Kanwisher, N., & Duchaine, B. C. (2007). Can generic expertise explain special processing for faces? *Trends in Cognitive Sciences*, 11(1), 8-15.
- Meissner, C. A., & Brigham, J. C. (2001). Thirty years of investigating the own-race bias in memory for faces: A meta-analytic review. *Psychology, Public Policy, & Law*, 7(1), 3-35.
- Mondloch, C. J., Le Grand, R., & Maurer, D. (2002). Configural face processing develops more slowly than featural face processing. *Perception*, 31, 553-566.
- Mondloch, C. J., Pathman, T., Maurer, D., Le Grand, R., & de Schonen, S. (2007). The composite face effect in six-year-old children: Evidence of adult-like holistic face processing. *Visual Cognition*, 15(5), 564-577.
- Morton, J., & Johnson, M. H. (1991). CONSPEC and CONLERN: A two-process theory of infant face recognition. *Psychological Review*, 98(2), 164-181.
- Ostrovsky, Y., Andalman, A., & Sinha, P. (2006). Vision following extended congenital blindness. *Psychological Science*, 17(12), 1009-1014.

- Otsuka, Y., Nakato, E., Kanazawa, S., Yamaguchi, M. K., Watanabe, S., & Kakigi, R. (2007). Neural activation to upright and inverted faces in infants measured by near infrared spectroscopy. *NeuroImage*, 34, 399-406.
- Pascalis, O., de Haan, M., Nelson, C. A., & de Schonen, S. (1998). Long-term recognition memory for faces assessed by visual paired comparison in 3- and 6-month-old infants. *Journal of Experimental Psychology: Learning, Memory and Cognition*, 24(1), 249-260.
- Pascalis, O., de Schonen, S., Morton, J., Deruelle, C., & Fabre-Grenet, M. (1995). Mother's face recognition by neonates: A replication and an extension. *Infant Behavior and Development*, 18, 79-85.
- Pascalis, O., Scott, L. S., Kelly, D. J., Shannon, R. W., Nicholson, E., Coleman, M., & Nelson, C. A. (2005). Plasticity of face processing in infancy. *Proceedings of the National Academy of Sciences of the United States of America*, 102(14), 5297-5300.
- Passarotti, A. M., Smith, J., DeLano, M., & Huang, J. (2007). Developmental differences in the neural bases of the face inversion effect show progressive tuning of face-selective regions to the upright orientation. *Neuroimage*, 34(4), 1708-1722.
- Pellicano, E., Jeffrey, L., Burr, D., & Rhodes, G. (2007). Abnormal adaptive face-coding mechanisms in children with autism spectrum disorder. *Current Biology*, 17(17), 1508-1542.
- Pellicano, E., Rhodes, G., & Peters, M. (2006). Are preschoolers sensitive to configural information in faces? *Developmental Science*, 9(3), 270-277.
- Perrett, D. I., Mistlin, A. J., Chitty, A. J., Smith, P. A., Potter, D. D., Broennimann, R., & Harries, M. (1988). Specialized face processing and hemispheric asymmetry in man and monkey: evidence from single unit and reaction time studies. *Behavioural Brain Research*, 29(3), 245-258.
- Pinski, M. A., DeSimone, K., Moore, T., Gross, C.G., Kastner, S. (2005). Representations of faces and body parts in macaque temporal cortex: a functional MRI study. *Proceedings of the National Academy Science, U.S.A.*, 102(19), 6996-7001.

- Polk, T. A., Park, J., Smith, M. R., & Park, D. C. (2007). Nature versus nurture in ventral visual cortex: a functional magnetic resonance imaging study of twins. *Journal of Neuroscience*, 27(51), 13921-13925.
- Pourtois, G., Schwartz, S., Seghier, M. L., Lazeyras, F., & Vuilleumier, P. (2005). Portraits or people? Distinct representations of face identity in the human visual cortex. *Journal of Cognitive Neuroscience*, 17(7), 1043-1057.
- Puce, A., Allison, T., Asgari, M., Gore, J. C., & McCarthy, G. (1996). Differential sensitivity of human visual cortex to faces, letterstrings, and textures: A functional magnetic resonance imaging study. *Journal of Neuroscience*, 16(16), 5205-5215.
- Rhodes, G., Brake, S., & Atkinson, A. P. (1993). What's lost in inverted faces? *Cognition*, 47, 25-57.
- Robbins, R., & McKone, E. (2003). Can holistic processing be learned for inverted faces? *Cognition*, 88, 79-107.
- Robbins, R., & McKone, E. (2007). No face-like processing for objects-of-expertise in three behavioural tasks. *Cognition*, 103(1), 34-79.
- Rodman, H. R., Scalaidhe, S. P., & Gross, C. G. (1993). Response properties of neurons in temporal cortical visual areas of infant monkeys. *Journal of Neurophysiology*, 70(3), 1115-1136.
- Rossion, B., Dricot, L., A., D., Bodart, J., Crommelinck, M., de Gelder, B., & Zoontjes, R. (2000). Hemispheric asymmetries for whole-based and part-based face processing in the human fusiform gyrus. *Journal of Cognitive Neuroscience*, 14(5), 793-802.
- Rotshtein, P., Henson, R. N., Treves, A., Driver, J., & Dolan, R. J. (2005). Morphing Marilyn into Maggie dissociates physical and identity face representations in the brain. *Nature Neuroscience*, 8(1), 107-113.
- Sai, F. Z. (2005). The role of the mother's voice in developing mother's face preference: Evidence for intermodal perception at birth. *Infant and Child Development*, 14, 29-50.

- Sangrigoli, S., & de Schonen, S. (2004). Effect of visual experience on face processing: a developmental study of inversion and non-native effects. *Developmental Science*, 7(1), 74-87.
- Sangrigoli, S., Pallier, C., Argenti, A.M., Ventureyra, V.A., de Schonen, S. (2005). Reversibility of the other-race effect in face recognition during childhood. *Psychological Sciences*, 16(6), 440-444.
- Saxe, R., Brett, M., & Kanwisher, N. (2006). Divide and conquer: a defense of functional localizers. *Neuroimage*, 30(4), 1088-1096.
- Scherf, K. S., Behrmann, M., Humphrey, K., & Luna, B. (2007). Visual category-selectivity for faces, places and objects emerges along different developmental trajectories. *Developmental Science*, 10(4), F15-F30.
- Schiltz, C., & Rossion, B. (2006). Faces are represented holistically in the human occipito-temporal cortex. *NeuroImage*, 32, 1385-1394.
- Sengpiel, F. (2007). The critical period, . *Current Biology*, 17, R742-R743.
- Simion, F., Macchi Cassia, V., Turati, C., & Valenza, E. (2003). Non-specific perceptual biases at the origins of face processing. In O. Pascalis & A. Slater (Eds.), *The development of face processing in infancy and early childhood: Current perspectives* (pp. 13-25). New York: Nova Science Publisher, Inc.
- Simons, D., J., & Levin, D. T. (1998). Failure to detect changes to people during real-world interaction. *Psychonomic Bulletin and Review*, 5(4), 644-649.
- Slater, A., Quinn, P. C., Hayes, R., & Brown, E. (2000). The role of facial orientation in newborn infants' preference for attractive faces. *Developmental Science*, 3(2), 181-185.
- Sugita, Y. (2008). Face perception in monkeys reared with no exposure to faces. *Proceedings of the National Academy of Sciences of the United States of America*, 105(1), 394-398.
- Tanaka, J., Kay, J. B., Grinnell, E., Stansfield, B., & Szechter, L. (1998). Face recognition in young children: When the whole is greater than the sum of its parts. *Visual Cognition*, 5, 479-496.

- Tanaka, J. W., & Farah, M. (1993). Parts and wholes in face recognition. *The Quarterly Journal of Experimental Psychology*, 46A(2), 225-245.
- Tanaka, J. W., & Sengco, J. A. (1997). Features and their configuration in face recognition. *Memory and Cognition*, 25(5), 583-592.
- Taylor, M. J., Batty, M., & Itier, R. J. (2004). The faces of development: a review of early face processing over childhood. *Journal of Cognitive Neuroscience*, 16(8), 1426-1442.
- Teunisse, J. P., & de Gelder, B. (2003). Face processing in adolescents with autistic disorder: the inversion and composite effects. *Brain and Cognition*, 52(3), 285-294.
- Tsao, D. Y., Freiwald, W. A., Knutsen, T. A., Mandeville, J. B., & Tootell, R. B. (2003). Faces and objects in macaque cerebral cortex. *Nature Neuroscience*, 6(9), 989-995.
- Turati, C., Bulf, H., & Simion, F. (2008). Newborns' face recognition over changes in viewpoint. *Cognition*, 106, 1300-1321.
- Turati, C., Macchi Cassia, V., Simion, F., & Leo, I. (2006). Newborns' face recognition: Role of inner and outer facial features. *Child Development*, 77(2), 297-311.
- Tzourio-Mazoyer, N., de Schonen, S., Crivello, F., Reutter, B., Aujard, Y., & Mazoyer, B. (2002). Neural correlates of woman face processing by 2-month-old infants. *NeuroImage*, 15, 454-461.
- Valentine, T. (1991). A unified account of the effects of distinctiveness, inversion, and race in face recognition. *The Quarterly Journal of Experimental Psychology*, 43A(2), 161-204.
- Valentine, T., & Bruce, V. (1986). The effects of distinctiveness in recognising and classifying faces. *Perception*, 15, 525-535.
- Webster, M. A., & MacLin, O. H. (1999). Figural aftereffects in the perception of faces. *Psychonomic Bulletin and Review*, 6(4), 647-653.

- Williams, M. A., Berberovic, N., & Mattingley, J. B. (2007). Abnormal fMRI adaptation to unfamiliar faces in a case of developmental prosopagnosia. *Current Biology*, 17(14), 1259-1264.
- Wilson, R. R., Blades, M., & Pascalis, O. (2007). What do children look at in an adult face with which they are personally familiar? *British Journal of Development Psychology*, 25, 375-382.
- Winston, J. S., Vuilleumier, P., & Dolan, R. J. (2003). Effects of low-spatial frequency components of fearful faces on fusiform cortex activity. *Current Biology*, 13(20), 1824-1829.
- Young, A. W., Hellawell, D., & Hay, D. C. (1987). Configurational information in face perception. *Perception*, 16, 747-759.
- Yovel, G., & Duchaine, B. (2006). Specialized face perception mechanisms extract both part and spacing information: evidence from developmental prosopagnosia. *Journal of Cognitive Neuroscience*, 18(4), 580-593.
- Yovel, G., & Kanwisher, N. (2005). The neural basis of the behavioral face-inversion effect. *Current Biology*, 15(24), 2256-2262.

Figures




Figure 1.






Figure 2

A. Restriction of range in young children: face effects increase with age



B. Restriction of range in adults: face effects decrease with age



C. No range restrictions: face effects are stable with age






Figure 3.


A) Left FFA




B) Right FFA




C) Right mid-fusiform gyrus



D) Right LOC E)



E) Face-selective right STS



E) Right PPA




Figure 4.

Figure 5.




Figure 6.




Figure 7.

	Newborns / deprived	< = 3 months	Later infancy	4 years	5 years	6 years	7 years	8 years	9 years	10 years	11 years	Adults
Behavioral Properties												
Ability to discriminate individual faces	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺
Inversion effect on discrimination (looking time or recognition memory)	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺
Composite-like effect, upright not inverted			☺									☺
Composite effect				☺	☺	☺	☺	☺	☺	☺	☺	☺
Part-whole effect, upright not inverted				☺	☺	☺	☺	☺	☺	☺	☺	☺
Part-in-spacing-altered-whole effect, upright not inverted				☺	☺	☺	☺	☺	☺	☺	☺	☺
Sensitivity to spacing changes	☺		☺	☺	☺	☺	☺	☺	☺	☺	☺	☺
Inversion effect on spacing sensitivity			☺			☺	☺	☺	☺	☺	☺	☺
Perceptual bias to upright in superimposed faces							☺	☺	☺			☺
Distinctiveness effects				☺	☺	☺	☺	☺	☺	☺	☺	☺
Adaptation aftereffects				☺	☺				☺	☺	☺	☺
Attractiveness preference, upright not inverted	☺											☺
Neural Properties												
Face-selective cells, macaques		☺										☺
Face-selective ERPs		☺		☺	☺	☺	☺	☺	☺	☺	☺	☺
FFA present				☺	☺	☺	☺	☺	☺	☺	☺	☺
Some type of inversion effect on neural response		?	☺	☺	☺	☺	☺	☺	☺	☺	☺	☺
Perceptual Narrowing												
Looking time discrimination of other race/species faces	☺	☺	X									X

Note: All references can be found in text except: Inversion effect on spacing sensitivity aged 6years-adult (Mondloch, Le Grand & Maurer, 2002); Adaptation aftereffect aged 9 years – adult (Pellicano, Jeffery, Burr & Rhodes, 2007).

