

**24.111: Philosophy of Quantum Mechanics, Spring 2016**  
**Homework 3 Solutions**

1. Read the excerpt on the readings page from Feynmann's Lecture on Physics, on the two-slit experiment. This experiment illustrates the "mystery of superposition" in much the same way the two-path experiment (discussed in class) does. The goal of this question is for you to see how close the parallel is. Consider these three hypotheses:
  - (1) Every electron (in the two-slit experiment) either passes through the top slit or passes through the bottom slit (but not both).
  - (2) Some electrons pass through both slits.
  - (3) Some electrons pass through neither slit.

For each of these hypotheses, provide arguments (reasons to believe) that it is false. (Note! Do not spend lots of time looking in the Feynman for the arguments — though his description of the experiment is definitely relevant. It's not clear that Feynman is trying to provide the arguments I am asking for. It's better to think about the analogous arguments with the two-path experiment.)

Solution: (i) If electrons always passed through exactly one of the slits, the outcome distribution should look like the 'sum' of the distributions you get with one or the other slit closed. But it does not. (ii) Could say: if you put detectors in both slits, you never register it going through both. I also accepted the answer that electron impacts on the screen always look like points, not 'spread out'; if the electron somehow went through both slits, there should be a time we it is 'spread out' in space. (iii) If you block both slits, you don't get any results.

2. Suppose an electron in state  $|0 \uparrow\rangle$ , followed by an electron in state  $|180 \downarrow\rangle$ , will pass through magnets oriented at  $180^\circ$ . Will they be deflected in the same direction? Different directions? Or are you unable to make any prediction with certainty? Justify your answer.

Solution: (The  $|180 \downarrow\rangle$  electron will definitely go down; you can use the statistical

algorithm to show that  $|0 \uparrow\rangle$  will too; or you can say that, since it is equal to  $-|180 \downarrow\rangle$ , it will too. But it was not enough to just say:  $|0 \uparrow\rangle = -|180 \downarrow\rangle$  so they'll definitely go in the same direction. For they will not definitely go in the same direction, if the SG magnets are at 90 degrees.)

3. In the following questions we have two electrons, and we represent their joint spin state with vectors in the tensor product space  $\mathbb{R}^2 \otimes \mathbb{R}^2$ . So in product vectors  $|x\rangle |y\rangle$  the “vector on the left”  $|x\rangle$  represents the state of electron 1, the vector on the right, the state of electron 2.

Show, in painstakingly pedantic detail, that the product vector  $|0 \uparrow\rangle |0 \downarrow\rangle$  is orthogonal to  $|0 \uparrow\rangle |0 \uparrow\rangle$ . (This is not hard, if you look up how the inner product of product vectors is defined in the notes.)

Solution:

$$|0 \uparrow\rangle |0 \downarrow\rangle \cdot |0 \uparrow\rangle |0 \uparrow\rangle = \langle 0 \uparrow | 0 \uparrow\rangle \cdot \langle 0 \downarrow | 0 \uparrow\rangle = 1 \cdot 0 = 0.$$

4. Write the electron-pair state  $\frac{1}{\sqrt{2}} |0 \uparrow\rangle |0 \downarrow\rangle - \frac{1}{\sqrt{2}} |0 \downarrow\rangle |0 \uparrow\rangle$  as a linear combination of the basis vectors  $|90 \uparrow\rangle |90 \uparrow\rangle, |90 \uparrow\rangle |90 \downarrow\rangle, |90 \downarrow\rangle |90 \uparrow\rangle, |90 \downarrow\rangle |90 \downarrow\rangle$ . (Substitute  $|0 \uparrow\rangle = \frac{1}{\sqrt{2}}(|90 \uparrow\rangle - |90 \downarrow\rangle)$ , and a similar expression for  $|0 \downarrow\rangle$ , into the above expression and use the linearity of the tensor product operation. For full credit you must show your work, not just write down the answer.)

Solution:

$$\begin{aligned}
& \frac{1}{\sqrt{2}} |0 \uparrow\rangle |0 \downarrow\rangle \\
&= \frac{1}{\sqrt{2}} \left( \frac{1}{\sqrt{2}} |90 \uparrow\rangle - \frac{1}{\sqrt{2}} |90 \downarrow\rangle \right) |0 \downarrow\rangle \\
&= \frac{1}{2} |90 \uparrow\rangle |0 \downarrow\rangle - \frac{1}{2} |90 \downarrow\rangle |0 \downarrow\rangle \\
&= \frac{1}{2} |90 \uparrow\rangle \left( \frac{1}{\sqrt{2}} |90 \uparrow\rangle + \frac{1}{\sqrt{2}} |90 \downarrow\rangle \right) - \frac{1}{2} |90 \downarrow\rangle \left( \frac{1}{\sqrt{2}} |90 \uparrow\rangle + \frac{1}{\sqrt{2}} |90 \downarrow\rangle \right) \\
&= \frac{1}{2\sqrt{2}} |90 \uparrow\rangle |90 \uparrow\rangle + \frac{1}{2\sqrt{2}} |90 \uparrow\rangle |90 \downarrow\rangle - \frac{1}{2\sqrt{2}} |90 \downarrow\rangle |90 \uparrow\rangle - \frac{1}{2\sqrt{2}} |90 \downarrow\rangle |90 \downarrow\rangle
\end{aligned}$$

For the other term,

$$\frac{1}{\sqrt{2}} |0 \downarrow\rangle |0 \uparrow\rangle,$$

by symmetry, take the solution for the first term and switch the order of the arrows in each term, giving

$$\frac{1}{2\sqrt{2}} |90 \uparrow\rangle |90 \uparrow\rangle + \frac{1}{2\sqrt{2}} |90 \downarrow\rangle |90 \uparrow\rangle - \frac{1}{2\sqrt{2}} |90 \uparrow\rangle |90 \downarrow\rangle - \frac{1}{2\sqrt{2}} |90 \downarrow\rangle |90 \downarrow\rangle.$$

Subtracting the second result from the first, we get

$$\frac{1}{\sqrt{2}} |90 \uparrow\rangle |90 \downarrow\rangle - \frac{1}{\sqrt{2}} |90 \downarrow\rangle |90 \uparrow\rangle.$$

5. Use the statistical algorithm to compute the probability that electron 1 will be deflected up through magnets at 0 degrees, when the state of the pair is  $|90 \uparrow\rangle |62 \downarrow\rangle$ .

Solution: A labeled orthonormal basis for this experiment is:  $|0 \uparrow\rangle |62 \uparrow\rangle, |0 \uparrow\rangle |62 \downarrow\rangle$  (these are labeled 'down' or -1),  $|0 \downarrow\rangle |62 \uparrow\rangle, |0 \downarrow\rangle |62 \downarrow\rangle$  (these are labeled 'up').

To compute the probability you square the inner product of the state vector with each 'up' basis vector, and add the results:  $(|90 \uparrow\rangle |62 \downarrow\rangle \cdot |0 \uparrow\rangle |62 \uparrow\rangle)^2 + (|90 \uparrow\rangle |62 \downarrow\rangle \cdot |0 \uparrow\rangle |62 \downarrow\rangle)^2 = 0 + (\langle 90 \uparrow | 0 \uparrow \rangle)^2 = 1/2$ .

6. Similarly, compute the probability that electron 1 will be deflected up through magnets at 0 degrees, when the state of the pair is  $1/\sqrt{2}|0 \uparrow\rangle|90 \uparrow\rangle + 1/\sqrt{2}|0 \uparrow\rangle|90 \downarrow\rangle$ .

Solution: A labeled orthonormal basis is  $|0 \uparrow\rangle|90 \uparrow\rangle, |0 \uparrow\rangle|90 \downarrow\rangle$  (these are the up vectors),  $|0 \downarrow\rangle|90 \uparrow\rangle, |0 \downarrow\rangle|90 \downarrow\rangle$ . Using the statistical algorithm we get that the probability is 1. (I didn't require you to do it this way: you could also have just 'seen' that the state vector is one where electron 1 is definitely an up at 0 degrees electron.)