

Synthesis of Molybdenum(VI) Neopentylidene Neopentylidyne Complexes


Hosein Tafazolian,^{†,‡} Richard R. Schrock,^{*,†,‡,§} and Peter Müller[†]

[†]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

[‡]Department of Chemistry, University of California at Riverside, Riverside, California 92521, United States

S Supporting Information

ABSTRACT: Mo(C-*t*-Bu)(CH-*t*-Bu)(Cl)(PMe₂Ph)₂ (**1**) was prepared as off-white crystals in 26% yield through addition of 2.5 equiv of Mg(CH₂-*t*-Bu)₂ to Mo(O)[OC(CF₃)₃]₄ in diethyl ether followed by 3 equiv of PMe₂Ph and a workup that includes dichloromethane (the source of Cl). Compound **1** is largely a *syn* isomer initially that equilibrates to give approximately a 1:1 mixture of *syn* and *anti* isomers within 1–2 h. Compound **1** reacts with Li(3,5-dimethylpyrrolide) to give Mo(C-*t*-Bu)(CH-*t*-Bu)(*η*¹-Me₂Pyr)(PMe₂Ph)₂ (**2a**) as a pale yellow solid in 76% yield, and **2a** reacts with Ph₃SiOH to give a mixture of *syn* and *anti* Mo(C-*t*-Bu)(CH-*t*-Bu)(OSiPh₃)(PMe₂Ph)₂ (**3a**) in 84% yield. All three compounds tend to lose PMe₂Ph to give 14e monophosphine complexes with the formulas Mo(C-*t*-Bu)(CH-*t*-Bu)(X)(PMe₂Ph) (X = Cl, Me₂Pyr, or OSiPh₃), none of which could be isolated. X-ray studies show the structures of **1**, **2a**, and **3a** to be analogous with τ values of 0.45, 0.53, and 0.69, respectively.



Imido alkyl complexes such as M(NR')₂(CH₂R)₂ (M = Mo or W; R' = aryl, *tert*-butyl, or adamantyl and CH₂R = neopentyl or neophyl), which can be prepared readily from M(NR')₂(dme)Cl₂ complexes and magnesium or lithium alkyls, yield M(NR')(CHR)X₂ complexes when one of the imido ligands is protonated with 2 equiv of HX (e.g., X = triflate).¹ However, analogous approaches to oxo alkylidene complexes usually are thwarted by the fact that oxo ligands are attacked by the alkylating agent and removed from the metal. Attempted alkylations of Mo oxo complexes more often give rise to low yields of oxo products, or none at all. For example, Osborn reported the synthesis of Mo(O)(CH₂-*t*-Bu)₃Cl and “Mo(O)(CH₂-*t*-Bu)₄” through addition of Mg(CH₂-*t*-Bu)₂ to Mo(O)Cl₄,² but experimental details (including yields) were not provided in either of the succinct reports, and Mo(O)(CH₂-*t*-Bu)₄ had to be reformulated (as noted in footnote 1 in ref 2b, without further details) as Mo(O)(CH₂-*t*-Bu)₃(OCH₂-*t*-Bu). The neopentoxide in Mo(O)(CH₂-*t*-Bu)₃(OCH₂-*t*-Bu) could arise through attack by the alkylating agent on the oxo ligand.³ It also has been reported that addition of 6 equiv of (*t*-BuCH₂)MgCl to MoO₂Cl₂ in diethyl ether gives the best reported yield (~35%) of Mo(C-*t*-Bu)(CH₂-*t*-Bu)₃.⁴ The analogous trimethylsilylmethylidyne complex, Mo(CSiMe₃)(CH₂SiMe₃)₃ (8% yield; liquid at 22 °C), has been prepared similarly with (Me₃SiCH₂)₃Mo≡Mo(CH₂SiMe₃)₃ (25% yield; crystalline) and unstable (Me₃SiCH₂)₃Mo=CHSiMe₃ (proposed) being two other metal-containing products.⁵

One exception to low yield alkylations of Mo oxo complexes is the class of molybdenum or tungsten compounds with the formula MO₂(CH₂R)₂(bipy) (CH₂R can be a variety of alkyls, including neopentyl), which can be prepared through alkylation of MO₂Cl₂(bipy) compounds with Grignard

reagents.⁶ The final M(VI) complexes are formed only after exposure of the crude product to water and air, so it appears likely that at least some, if not most, of the metal is reduced and then reoxidized in the presence of water and air. WO₂(CH₂-*t*-Bu)₂(bipy) is a precursor to W(O)(CH-*t*-Bu)-Cl₂L₂ upon reaction of WO₂(CH₂-*t*-Bu)₂(bipy) with Me₃SiCl, ZnCl₂, and 2 equiv of L (e.g., L = PPhMe₂); W(O)(CH₂-*t*-Bu)₂Cl₂(bipy) is a plausible intermediate.⁷ This approach to the synthesis of Mo(O)(CH-*t*-Bu)Cl₂L₂ complexes failed in our hands so far. A second exception is the reaction between Mo(O)(Cl)₂(*t*-Bu₃P=N)₂ and 2 equiv of LiCH₂SiMe₃ to give Mo(O)(CHSiMe₃)(*t*-Bu₃P=N)₂ in 87% yield;⁸ Mo(O)(CH₂SiMe₃)₂(*t*-Bu₃P=N)₂ is the plausible intermediate in this reaction. Unfortunately, Mo(O)(CHSiMe₃)(*t*-Bu₃P=N)₂ and analogous molybdenum imido alkylidene complexes that contain two *t*-Bu₃SiO⁻ ligands (which are sterically approximately equivalent to *t*-Bu₃P=N⁻ ligands)⁹ are relatively unreactive in olefin metathesis reactions.

We recently prepared molybdenum oxo alkylidene complexes through a controlled addition of 1 equiv of water to molybdenum(VI) OC(CF₃)₃ or OC(CF₃)₂Me benzylidyne complexes.¹⁰ In the hope that more direct routes to these Mo oxo alkylidene complexes could be successful, we turned to alkylation of Mo(O)[OC(CF₃)₃]₄.¹¹ Yellow Mo(O)[OC(CF₃)₃]₄ can be prepared readily from Mo(O)Cl₄ and 4 equiv of NaOC(CF₃)₃. It can be sublimed under a good vacuum (<10⁻² mm) at 60–80 °C. It is poorly soluble in organic solvents (pentane, benzene, toluene) but partially soluble in diethyl ether or dichloromethane.

Received: June 19, 2019

Published: July 24, 2019

We decided to explore alkylation of $\text{Mo}(\text{O})[\text{OC}(\text{CF}_3)_3]_4$ with $\text{Mg}(\text{CH}_2\text{-}t\text{-Bu})_2$ in the presence of PMe_2Ph in order to trap any monometallic products. Alkylation reactions using 0.5, 1.0, 2.0, and 2.5 equiv of $\text{Mg}(\text{CH}_2\text{CMe}_3)_2$ in the presence of PMe_2Ph in diethyl ether were explored initially. Each crude sample was dissolved in C_6D_6 or CD_2Cl_2 and examined by proton and/or ^{31}P NMR. A small multiplet in the region characteristic of a molybdenum alkylidene having two phosphines bound to the metal in proton NMR spectra in CD_2Cl_2 grew slightly over a day, but no other product could be identified. When 2.5 equiv of $\text{Mg}(\text{CH}_2\text{CMe}_3)_2$ were used and CH_2Cl_2 was part of the workup, the alkylidene could be extracted into pentane and crystallized reproducibly from a concentrated pentane solution at -20°C as off-white needles in a yield of 26%. When the crude brown residue was dissolved in C_6D_6 and not exposed to dichloromethane, no alkylidene resonance was observed, even after addition of 2 equiv of PMe_2Ph . However, when a drop of CD_2Cl_2 was added to this sample, the alkylidene resonance grew in with time. The *syn* alkylidene resonance can be observed in the ^1H NMR spectrum 5 h after addition of CD_2Cl_2 and stops increasing after ~ 60 h. There was no further change in the ^1H NMR spectrum.

An X-ray structural study confirms that **1** is the “Ene/Yne” complex, $\text{Mo}(\text{C-}t\text{-Bu})(\text{CH-}t\text{-Bu})(\text{PPhMe}_2)_2\text{Cl}$ (**1**; eq 1 and





Figure 1. Structure of **1** ($\tau = 0.45$).

Figure 1), not an oxo alkylidene complex. The overall geometry is approximately halfway between a TBP and an SP ($\tau^{12} = 0.45$), although **1** (and related derivatives described below) will be drawn as a TBP with L in the apical positions for convenience. The bond lengths and angles (**Table 1**) are not unusual for neopentylidene ligands (1.691(3) Å, 166.80°) or *syn* neopentylidene ligands (1.985(3) Å, 152.57°). The chloride must come from dichloromethane, a phenomenon that has been reported in the literature for certain $\text{Mo}^{13\text{a,b}}$ and $\text{Re}^{13\text{c}}$ compounds. No mechanistic details have been reported for formation of chloride complexes in the presence of

Table 1. Selected Bond Lengths and Angles in **1**, **2a**, and **3a**

	1	2a	3a
Mo1–C1	1.691(3)	1.661(11)	1.764(6)
Mo1–C6	1.985(3)	2.000(9)	1.9214(14)
Mo1–P1	2.5089(3)	2.5092(10)	2.5202(3)
Mo1–P2	2.5196(3)	2.5092(10)	2.5366(4)
Mo1–C1– <i>t</i> -Bu	166.80	172.59	170.61
Mo1–C6– <i>t</i> -Bu	152.57	151.52	152.16
P1–Mo1–P2	160.28	164.97	171.25
C1–Mo1–C6	104.33	102.49	103.51
Mo1–X ^a	2.5197(3)	2.227(3)	2.0748(9)


^aX = Cl (in **1**), N (in **2a**), or O (in **3a**).

dichloromethane, and we also do not want to propose any for forming **1** at this stage.

Compound **1** is related to tantalum bisneopentylidene complexes (e.g., $\text{Ta}(\text{CH-}t\text{-Bu})_2(\text{PMe}_3)_2\text{Cl}$) which contain two different neopentylidene ligands, one of which has a significantly greater agostic interaction of its alkylidene CH_α bond with the metal than does the other.¹⁴ It also is related to $\text{W}(\text{C-}t\text{-Bu})(\text{CH-}t\text{-Bu})(\text{CH}_2\text{-}t\text{-Bu})(\text{PMe}_3)_2$,¹⁵ square pyramidal $\text{W}(\text{C-}t\text{-Bu})(\text{CH-}t\text{-Bu})(\text{CH}_2\text{-}t\text{-Bu})(\text{dmpe})$,¹⁵ and trimethylsilyl relatives.¹⁶ The alkylidene is in the *syn* orientation in the crystal chosen for the X-ray study. Compound **1** is stable in benzene solution for several days at temperatures as high as 80°C . Attempts to remove one of the phosphines from a C_6D_6 solution of **1** through addition of 1 equiv of $\text{B}(\text{C}_6\text{F}_5)_3$ to **1** in 0.5 mL of C_6D_6 (0.04 M) led only to decomposition to unidentified products. We propose that loss of phosphine in the solid state at 1 atm is the reason for a failure to obtain satisfactory elemental analyses for **1** and related derivatives **2a** and **3a** described below.

Proton NMR spectra of **1** usually show two triplet alkylidene resonances at 13.87 ppm (*anti*, $^1J_{\text{CH}} = 137.0$ Hz) and 12.26 ppm (*syn*, $^1J_{\text{CH}} = 100.6$ Hz). Freshly isolated **1** is usually *syn*-rich, occasionally as high as 95% *syn* (see **SI**). Upon recrystallizations of **1**, the *syn/anti* ratio approaches 1:1 and remains so.

Compound **1** reacts with Li(2,5-dimethylpyrrolide) to yield the monopyrrolide complex, **2a** (eq 2). Compound **2a** can be



crystallized from pentane as off-white needles in 76% yield. An X-ray study showed the structure of **2a** (**Figure 2**) to be analogous to that of **1**, with the pyrrolide bound to the metal in an η^1 fashion. The overall geometry and bond lengths and angles in **2a** are similar to those found in **1** (**Table 1**).

In ^1H NMR spectra of pure samples of **2a**, two *syn* alkylidene resonances are found, one triplet (at 11.65 ppm in C_6D_6) for **2a** and one doublet ($\sim 10\%$ of the total at 13.48 ppm in C_6D_6). We ascribe the latter to a monophosphine complex, $\text{Mo}(\text{C-}t\text{-Bu})(\text{CH-}t\text{-Bu})(\text{PPhMe}_2)(\text{Me}_2\text{Py})$ (**2b**). Compounds **2a** and **2b** (plus phosphine) are in equilibrium at room temperature; in a sample of 0.033 M **2a** in C_6D_6 , the K_{eq} was found by ^1H NMR spectroscopy to be 3.7×10^{-4} M at room temperature. The ^{31}P NMR of a sample of **2a** in C_6D_6 at room temperature shows three resonances at 30.4, 7.3, and -46.5





Figure 2. Structure of **2a** ($\tau = 0.53$).

ppm for **2b**, **2a**, and free phosphine, respectively. The magnetization exchange rates between these two species were measured through a series of 1D EXSY experiments at different temperatures (>40 °C). For dissociation of phosphine from **2b**, the ΔH^\ddagger was found to be 121 (± 8) kJ/mol and ΔS^\ddagger was found to be 0.095 (± 0.026) kJ/mol K. For the reverse reaction, ΔH^\ddagger and ΔS^\ddagger were found to be 76 (± 5) kJ/mol and -0.028 (± 0.017) kJ/mol K, respectively. **2b** was also the major product in an NMR-scale reaction in C_6D_6 of **2a** with 1 equiv of $B(C_6F_5)_3$ (see SI).

Compound **2a** reacts cleanly with 1 equiv of triphenylsilanol to generate **3a** (eq 3). An X-ray study shows the structure of **3a**



to be analogous to the structures of **1** and **2a** (Table 1 and Figure 3). An alkyldiene multiplet can be observed in the proton NMR spectra of **3a** at 13.29 ppm (in C_6D_6) with a $^1J_{CH}$ of 100.6 Hz that is characteristic of a *syn* alkyldiene. No




Figure 3. Structure of **3a** ($\tau = 0.69$).

doublet alkyldiene resonance characteristic of a mono-phosphine adduct can be detected by 1H NMR in this case, even at temperatures up to 70 °C. Nevertheless, one of the phosphines is labile enough to be scavenged as a borane adduct upon addition of 1 equiv of $B(C_6F_5)_3$ to **3a** in C_6D_6 . The resulting mono-phosphine adduct shows a characteristic *syn* alkyldiene resonance in the 1H NMR spectrum at 13.32 ppm (d, $J_{HP} = 3.8$ Hz, $J_{CH} = 106.1$ Hz). To date, we have not been able to isolate **3b**.

An exploratory NMR-scale ROMP reaction showed that cyclooctene was polymerized by **3a** (1%) in C_6D_6 upon addition of $B(C_6F_5)_3$ (1.1 equiv). The amount of cyclooctene consumed was 69% in 1 h, 76% in 4 h, and 87% in 24 h.

We will seek higher yield routes to **1** and related “Ene/Yne” complexes, preferably those that do not require removal of oxo ligands from the metal and/or scavenging of chloride from dichloromethane. We are especially interested in 14e “stereogenic at metal” $Mo(CR)(CH-t-Bu)(L)X$ complexes that contain a single donor (L) and anionic X, e.g., **2b** and **3b** and analogs. These are 14e relatives of (largely) imido alkyldiene^{1,17} and oxo^{7,10} alkyldiene complexes that have been explored to date as metathesis initiators. An important question is whether an alkyldiene ligand (CR) can survive unchanged in a sustained metathesis reaction, a possibility that has not been addressed to our knowledge.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.organomet.9b00412.

Detailed NMR data and spectra for all compounds and details of X-ray studies (PDF)

Accession Codes

CCDC 1922920–1922922 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: rrs@mit.edu.

ORCID

Richard R. Schrock: 0000-0001-5827-3552

Author Contributions

H.T. performed all synthetic work while P.M. performed all X-ray structural studies.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for financial support from the National Science Foundation (CHE-1463707) and the University of California at Riverside.

■ REFERENCES

(1) Schrock, R. R. High Oxidation State Multiple Metal-Carbon Bonds. *Chem. Rev.* 2002, 102, 145–180.

- (2) (a) Kress, J. R. M.; Russell, M. J. M.; Wesolek, M. G.; Osborn, J. A. Tungsten(VI) and Molybdenum(VI) Oxo-alkyl Species. Their Role in the Metathesis of Olefins. *J. Chem. Soc., Chem. Commun.* **1980**, 431–432. (b) Kress, J.; Wesolek, M.; Le Ny, J.-P.; Osborn, J. A. Molecular Complexes for Efficient Metathesis of Olefins. The Oxo-ligand as a Catalyst-Cocatalyst Bridge and the Nature of the Active Species. *J. Chem. Soc., Chem. Commun.* **1981**, 0, 1039–1040.
- (3) Aguero, A.; Osborn, J. A. Synthetic Routes to Metal-Alkylidene Complexes. *New J. Chem.* **1988**, 12, 111–118.
- (4) McCullough, L. G.; Schrock, R. R.; Dewan, J. C.; Murdzek, J. S. Preparation of Trialkoxymolybdenum(VI) Alkylidyne Complexes, Their Reactions with Acetylenes, and the X-Ray Structure of $\text{Mo}[\text{C}(\text{CMe}_3)_2][\text{OCH}(\text{CF}_3)_2]_2(\text{C}_5\text{H}_5\text{N})_2$. *J. Am. Chem. Soc.* **1985**, 107, 5987–5998.
- (5) Andersen, R. A.; Chisholm, M. H.; Gibson, J. F.; Reichert, W. W.; Rothwell, I. P.; Wilkinson, G. (Trimethylsilyl)methylidene and (Trimethylsilyl)methylidyne Compounds of Molybdenum and Tungsten: $(\text{Me}_3\text{SiCH}_2)_3\text{M}(\text{CSiMe}_3)$ (M = Mo, W) and $(\text{Me}_3\text{SiCH}_2)_3\text{M} = \text{CHSiMe}_3$. *Inorg. Chem.* **1981**, 20, 3934–3936.
- (6) (a) Schrauzer, G. N.; Hughes, L. A.; Strampach, N.; Robinson, P. R.; Schlemper, E. O. Studies of Molybdenum Compounds. 1. Synthesis and Structure of Dioxo(dimethyl)(2,2'-bipyridyl)-molybdenum(VI), Prototype of a New Class of Organomolybdenum(VI) Compounds. *Organometallics* **1982**, 1, 44–47. (b) Schrauzer, G. N.; Hughes, L. A.; Strampach, N.; Ross, F.; Ross, D.; Schlemper, E. O. Studies of Organomolybdenum Compounds. 2. Synthesis, Structure, and Properties of Dioxidoneopentyl(2,2'-bipyridyl)molybdenum(VI) and of Related Compounds. *Organometallics* **1983**, 2, 481–485. (c) Schrauzer, G. N.; Hughes, L. A.; Therien, M. J.; Schlemper, E. O.; Ross, F.; Ross, D. Studies of Molybdenum Compounds. 4. Synthesis and Structure of Dibenzyl(2,2'-bipyridyl)dioxomolybdenum(VI). *Organometallics* **1983**, 2, 1163–1166. (d) Schrauzer, G. N.; Schlemper, E. O.; Hui, L. N.; Rubin, K.; Zhang, X.; Long, X.; Chin, C. S. Studies of Molybdenum Compounds. 5. Diethyl(2,2'-bipyridyl)-dioxomolybdenum(VI) and Other Higher Dialkyl Derivatives of Dioxomolybdenum(VI). *Organometallics* **1986**, 5, 2452–2456. (e) Schrauzer, G. N.; Zhang, X.; Liu, N.; Schlemper, E. O. Studies of Molybdenum Compounds. 6. Diaryl(2,2'-bipyridyl)-dioxomolybdenum(VI) and Related Compounds. *Organometallics* **1988**, 7, 279–282. (f) Zhang, C.; Zhang, X.; Liu, N. H.; Schrauzer, G. N.; Schlemper, E. O. Diphenyl(2,2'-bipyridyl)dioxomolybdenum(VI) and -tungsten(VI): A Comparative Study. *Organometallics* **1990**, 9, 1307–1311. (g) Zhang, C.; Schlemper, E. O.; Schrauzer, G. N. Synthesis, Structure, and Reactions of 2,2'-Bipyridyl Complexes of Tetramethyloxotungsten(VI) and Dimethyldioxotungsten(VI) and of Related Compounds. *Organometallics* **1990**, 9, 1016–1020.
- (7) (a) Peryshkov, D. V.; Schrock, R. R.; Takase, M. K.; Müller, P.; Hoveyda, A. H. Z-Selective Olefin Metathesis Reactions Promoted by Tungsten Oxo Alkylidene Complexes. *J. Am. Chem. Soc.* **2011**, 133, 20754–20757. (b) Peryshkov, D. V.; Schrock, R. R. Synthesis of Tungsten Oxo Alkylidene Complexes. *Organometallics* **2012**, 31, 7278–7286.
- (8) Varjas, C. J.; Powell, D. R.; Thomson, R. K. Rapid Access to an Oxo-Alkylidene Complex of Mo(VI). *Organometallics* **2015**, 34, 4806–4809.
- (9) Wampler, K. M.; Hock, A. S.; Schrock, R. R. Synthesis of Molybdenum Imido Alkylidene Complexes that Contain Siloxides. *Organometallics* **2007**, 26, 6674–6680.
- (10) (a) Bukhryakov, K. V.; Schrock, R. R.; Hoveyda, A. H.; Tsay, C.; Müller, P. Syntheses of Molybdenum Oxo Alkylidene Complexes Through Addition of Water to an Alkylidyne Complex. *J. Am. Chem. Soc.* **2018**, 140, 2797–2800. (b) Zhai, F.; Bukhryakov, K. V.; Schrock, R. R.; Hoveyda, A. H.; Tsay, C.; Müller, P. Syntheses of Molybdenum Oxo Benzylidene Complexes. *J. Am. Chem. Soc.* **2018**, 140, 13609–13613.
- (11) (a) Johnson, D. A.; Taylor, J. C.; Waugh, A. B. Volatile Perfluoro-*t*-butoxides of Oxo Molybdenum(VI). *Inorg. Nucl. Chem. Lett.* **1979**, 15, 205–206. (b) Johnson, D. A.; Taylor, J. C.; Waugh, A. B. Crystal and Molecular Structure of Tetra(*tert*-Perfluoro Butoxy) Oxo Mo(VI). *J. Inorg. Nucl. Chem.* **1980**, 42, 1271–1275.
- (12) Addison, A. W.; Rao, T. N.; Reedijk, J.; van Rijn, J.; Verschoor, G. C. Synthesis, Structure, and Spectroscopic Properties of Copper(II) Compounds Containing Nitrogen-Sulfur Donor Ligands: the Crystal and Molecular Structure of Aqua[1,7-Bis(N-methylbenzimidazol-2'-yl)-2,6-dithiaheptane]copper(II) Perchlorate. *J. Chem. Soc., Dalton Trans.* **1984**, 1349–1356.
- (13) (a) Borgmann, C.; Limberg, C.; Driess, A. Synthesis, structure and properties of bisallyltris(trimethylphosphine) molybdenum(II) X-ray crystal structures of $[\text{Mo}(\eta^1:\eta^2\text{-C}_3\text{H}_5)_2(\text{PMe}_3)_3]$ and $\text{PMe}_3^+[\text{MoCl}_4(\text{PMe}_3)_2]^-$. *J. Organomet. Chem.* **1997**, 541, 367–375. (b) Fürstner, A.; Mathes, C.; Lehmann, C. W. Mo[N(*t*-Bu)(Ar)]₃ Complexes As Catalyst Precursors: In Situ Activation and Application to Metathesis Reactions of Alkynes and Dienes. *J. Am. Chem. Soc.* **1999**, 121, 9453–9454. (c) Huhmann-Vincent, J.; Scott, B. L.; Kubas, G. J. Rhenium Complexes with Weakly Coordinating Solvent Ligands, *cis*-[Re(PR₃)(CO)₄(L)][BAR_F], L) CH₂Cl₂, Et₂O, NC₅F₅: Decomposition to Chloride-Bridged Dimers in CH₂Cl₂ Solution. *Inorg. Chem.* **1999**, 38, 115–124.
- (14) (a) Fellmann, J. D.; Rupprecht, G. A.; Wood, C. D.; Schrock, R. R. Multiple Metal-Carbon Bonds. 11. Bisneopentylidene Complexes of Niobium and Tantalum. *J. Am. Chem. Soc.* **1978**, 100, 5964–5966. (b) Fellmann, J. D.; Schrock, R. R.; Rupprecht, G. A. Trigonal-Bipyramidal Bis(neopentylidene), Neopentylidene/Ethylene, and Bis(ethylene) Complexes of Tantalum and How They React with Ethylene. A Catalyst for Rapidly Dimerizing Ethylene to 1-Butene. *J. Am. Chem. Soc.* **1981**, 103, 5752–5758.
- (15) (a) Murdzek, J. S.; Schrock, R. R. High Oxidation State Alkylidyne Complexes. *Carbyne Complexes*; VCH Publishers: Weinheim, NY, 1988; p 147. (b) Schrock, R. R. The Discovery and Development of High Oxidation State Alkylidyne Complexes for Alkyne Metathesis. In *Handbook of Metathesis*; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, 2003; pp 173–189. (c) Churchill, M. R.; Youngs, W. J. Crystal Structure and Molecular Geometry of W(=CCMe₃)(=CHCMe₃)(CH₂CMe₃)(dmpe), a Mononuclear Tungsten(VI) Complex with Metal-Alkylidene, Metal-Alkylidene, and Metal-Alkyl Linkages. *Inorg. Chem.* **1979**, 18, 2454–2458.
- (16) (a) Dougan, B. A.; Xue, Z.-L. Reaction of a Tungsten Alkylidyne Complex with a Chelating Diphosphine. α -Hydrogen Migration in the Intermediates and Formation of an Alkyl Alkylidene Alkylidyne Complex. *Organometallics* **2009**, 28, 1295–1302. (b) Morton, L. A.; Chen, S.; Qiu, H.; Xue, Z.-L. Preparation of Tungsten Alkyl Alkylidene Alkylidyne Complexes and Kinetic Studies of Their Formation. *J. Am. Chem. Soc.* **2007**, 129, 7277–7283.
- (17) Schrock, R. R. Recent Advances in High Oxidation State Mo and W Imido Alkylidene Chemistry. *Chem. Rev.* **2009**, 109, 3211–3226.