

**Erratum: Evolution of circular, nonequatorial orbits of Kerr black holes
due to gravitational-wave emission
[Phys. Rev. D **61**, 084004 (2000)]**

Scott A. Hughes

(Received 4 March 2003; published 25 April 2003)

DOI: 10.1103/PhysRevD.67.089901

PACS number(s): 04.30.Db, 04.25.Nx, 95.30.Sf, 99.10.Cd

Equation (A5) should read

$$\langle slm | \cos^2 \theta | sjm \rangle = \frac{1}{3} \delta_{jl} + \frac{2}{3} \sqrt{\frac{2j+1}{2l+1}} \langle j, 2, m, 0 | l, m \rangle \langle j, 2, -s, 0 | l, -s \rangle \equiv c_{j,l,2}^m,$$

$$\langle slm | \cos \theta | sjm \rangle = \sqrt{\frac{2j+1}{2l+1}} \langle j, 1, m, 0 | l, m \rangle \langle j, 1, -s, 0 | l, -s \rangle \equiv c_{j,l,1}^m,$$

$$\langle slm | sjm \rangle = \delta_{jl}. \tag{A5}$$

The factors under the square roots were inverted in the published version of this paper.

I thank Marc Favata for bringing this error to my attention. Note that the code which was used for the work reported in this paper (and in several followup works) had these equations implemented correctly.