

# The Price of Robustness

Dimitris Bertsimas

Sloan School of Management, Massachusetts Institute of Technology, E53-363, Cambridge, Massachusetts 02139, dbertsim@mit.edu

Melvyn Sim

Operations Research Center, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, melvyn@mit.edu

A robust approach to solving linear optimization problems with uncertain data was proposed in the early 1970s and has recently been extensively studied and extended. Under this approach, we are willing to accept a suboptimal solution for the nominal values of the data in order to ensure that the solution remains feasible and near optimal when the data changes. A concern with such an approach is that it might be too conservative. In this paper, we propose an approach that attempts to make this trade-off more attractive; that is, we investigate ways to decrease what we call the price of robustness. In particular, we flexibly adjust the level of conservatism of the robust solutions in terms of probabilistic bounds of constraint violations. An attractive aspect of our method is that the new robust formulation is also a linear optimization problem. Thus we naturally extend our methods to discrete optimization problems in a tractable way. We report numerical results for a portfolio optimization problem, a knapsack problem, and a problem from the Net Lib library.

*Subject classifications:* Programming, stochastic: robust approach for solving LP/MIP with data uncertainties.

*Area of review:* Financial Services.

*History:* Received September 2001; revision received August 2002; accepted December 2002.

## 1. Introduction

The classical paradigm in mathematical programming is to develop a model that assumes that the input data is precisely known and equal to some nominal values. This approach, however, does not take into account the influence of data uncertainties on the quality and feasibility of the model. It is therefore conceivable that as the data take values different than the nominal ones, several constraints may be violated, and the optimal solution found using the nominal data may no longer be optimal or even feasible. This observation raises the natural question of designing solution approaches that are immune to data uncertainty; that is, they are “robust.”

To illustrate the importance of robustness in practical applications, we quote from the case study by Ben-Tal and Nemirovski (2000) on linear optimization problems from the Net Lib library:

In real-world applications of Linear Programming, one cannot ignore the possibility that a small uncertainty in the data can make the usual optimal solution completely meaningless from a practical viewpoint.

Naturally, the need arises to develop models that are immune, as far as possible, to data uncertainty. The first step in this direction was taken by Soyster (1973), who proposes a linear optimization model to construct a solution that is feasible for all data that belong in a convex set. The resulting model produces solutions that are too conservative in the sense that we give up too much of optimality for the nominal problem in order to ensure robustness

(see the comments of Ben-Tal and Nemirovski 2000). Soyster considers the linear optimization problem

$$\begin{aligned} & \text{maximize} && \mathbf{c}'\mathbf{x} \\ & \text{subject to} && \sum_{j=1}^n \mathbf{A}_j x_j \leq \mathbf{b}, \quad \forall \mathbf{A}_j \in K_j, \quad j = 1, \dots, n \\ & && \mathbf{x} \geq \mathbf{0}, \end{aligned}$$

where the uncertainty sets  $K_j$  are convex. Note that the case considered is “columnwise” uncertainty; i.e., the columns  $\mathbf{A}_j$  of the constraint matrix are known to belong to a given convex set  $K_j$ . Soyster (1973) shows that the problem is equivalent to

$$\begin{aligned} & \text{maximize} && \mathbf{c}'\mathbf{x} \\ & \text{subject to} && \sum_{j=1}^n \bar{\mathbf{A}}_j x_j \leq \mathbf{b}, \\ & && \mathbf{x} \geq \mathbf{0}, \end{aligned} \tag{1}$$

where  $\bar{a}_{ij} = \sup_{\mathbf{A}_j \in K_j} (A_{ij})$ .

A significant step forward for developing a theory for robust optimization was taken independently by Ben-Tal and Nemirovski (1998, 1999, 2000), El-Ghaoui and Lebret (1997), and El-Ghaoui et al. (1998). To address the issue of overconservatism, these papers proposed less conservative models by considering uncertain linear problems with ellipsoidal uncertainties, which involve solving the *robust counterparts* of the nominal problem in the form of conic quadratic problems (see Ben-Tal and Nemirovski 1999).

With properly chosen ellipsoids, such a formulation can be used as a reasonable approximation to more complicated uncertainty sets. However, a practical drawback of such an approach is that it leads to nonlinear, although convex, models, which are more demanding computationally than the earlier linear models by Soyster (1973) (see also the discussion in Ben-Tal and Nemirovski 2000).

In this research, we propose a new approach for robust linear optimization that retains the advantages of the linear framework of Soyster (1973). More importantly, our approach offers full control on the degree of conservatism for every constraint. We protect against violation of constraint  $i$  deterministically, when only a prespecified number  $\Gamma_i$  of the coefficients changes; that is, we guarantee that the solution is feasible if less than  $\Gamma_i$  uncertain coefficients change. Moreover, we provide a probabilistic guarantee that even if more than  $\Gamma_i$  change, then the robust solution will be feasible with high probability. In the process we prove a new, to the best of our knowledge, tight bound on sums of symmetrically distributed random variables. In this way, the proposed framework is at least as flexible as the one proposed by Ben-Tal and Nemirovski (1998, 1999, 2000), El-Ghaoui and Lebret (1997), El-Ghaoui et al. (1998), and possibly more. Unlike these approaches, the robust counterparts we propose are linear optimization problems, and thus our approach readily generalizes to discrete optimization problems. To the best of our knowledge, there was no similar work done in the robust discrete optimization domain that involves deterministic and probabilistic guarantees of constraints against violation.

## Structure of the Paper

In §2, we present the different approaches for robust linear optimization from the literature and discuss their merits. In §3, we propose the new approach and show that it can be solved as a linear optimization problem. In §4, we show that the proposed robust LP has attractive probabilistic and deterministic guarantees. Moreover, we perform sensitivity analysis of the degree of protection the proposed method offers. We provide extensions to our basic framework dealing with correlated uncertain data in §5. In §6, we apply the proposed approach to a portfolio problem, a knapsack problem, and a problem from the Net Lib library. Finally, §7 contains some concluding remarks.

## 2. Robust Formulation of Linear Programming Problems

### 2.1. Data Uncertainty in Linear Optimization

We consider the following nominal linear optimization problem:

$$\begin{aligned} & \text{maximize} && \mathbf{c}'\mathbf{x} \\ & \text{subject to} && \mathbf{Ax} \leq \mathbf{b} \\ & && \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}. \end{aligned}$$

In the above formulation, we assume that data uncertainty only affects the elements in matrix  $\mathbf{A}$ . We assume without loss of generality that the objective function  $\mathbf{c}$  is not subject to uncertainty, since we can use the objective maximize  $z$ , add the constraint  $z - \mathbf{c}'\mathbf{x} \leq 0$ , and thus include this constraint into  $\mathbf{Ax} \leq \mathbf{b}$ .

**Model of Data Uncertainty U.** Consider a particular row  $i$  of the matrix  $\mathbf{A}$  and let  $J_i$  represent the set of coefficients in row  $i$  that are subject to uncertainty. Each entry  $a_{ij}$ ,  $j \in J_i$  is modeled as a symmetric and bounded random variable  $\tilde{a}_{ij}$ ,  $j \in J_i$  (see Ben-Tal and Nemirovski 2000) that takes values in  $[a_{ij} - \hat{a}_{ij}, a_{ij} + \hat{a}_{ij}]$ . Associated with the uncertain data  $\tilde{a}_{ij}$ , we define the random variable  $\eta_{ij} = (\tilde{a}_{ij} - a_{ij})/\hat{a}_{ij}$ , which obeys an unknown but symmetric distribution, and takes values in  $[-1, 1]$ .

### 2.2. The Robust Formulation of Soyster

As we have mentioned in the introduction, Soyster (1973) considers columnwise uncertainty. Under the model of data uncertainty U, the robust Formulation (1) is as follows:

$$\begin{aligned} & \text{maximize} && \mathbf{c}'\mathbf{x} \\ & \text{subject to} && \sum_j a_{ij}x_j + \sum_{j \in J_i} \hat{a}_{ij}y_j \leq b_i \quad \forall i \\ & && -y_j \leq x_j \leq y_j \quad \forall j \\ & && \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \\ & && \mathbf{y} \geq \mathbf{0}. \end{aligned} \tag{2}$$

Let  $\mathbf{x}^*$  be the optimal solution of Formulation (2). At optimality, clearly,  $y_j = |x_j^*|$ , and thus

$$\sum_j a_{ij}x_j^* + \sum_{j \in J_i} \hat{a}_{ij}|x_j^*| \leq b_i \quad \forall i.$$

We next show that for every possible realization  $\tilde{a}_{ij}$  of the uncertain data, the solution remains feasible; that is, the solution is “robust.” We have

$$\begin{aligned} \sum_j \tilde{a}_{ij}x_j^* &= \sum_j a_{ij}x_j^* + \sum_{j \in J_i} \eta_{ij}\hat{a}_{ij}x_j^* \\ &\leq \sum_j a_{ij}x_j^* + \sum_{j \in J_i} \hat{a}_{ij}|x_j^*| \leq b_i \quad \forall i. \end{aligned}$$

For every  $i$ th constraint, the term,  $\sum_{j \in J_i} \hat{a}_{ij}|x_j^*|$  gives the necessary “protection” of the constraint by maintaining a gap between  $\sum_j a_{ij}x_j^*$  and  $b_i$ .

### 2.3. The Robust Formulation of Ben-Tal and Nemirovski

Although the Soyster method (1973) admits the highest protection, it is also the most conservative in practice in the sense that the robust solution has an objective function value much worse than the objective function value of the solution of the nominal linear optimization problem. To

address this conservatism, Ben-Tal and Nemirovski (2000) propose the following robust problem:

$$\begin{aligned}
 & \text{maximize} && \mathbf{c}'\mathbf{x} \\
 & \text{subject to} && \sum_j a_{ij}x_j + \sum_{j \in J_i} \hat{a}_{ij}y_{ij} \\
 & && + \Omega_i \sqrt{\sum_{j \in J_i} \hat{a}_{ij}^2 z_{ij}^2} \leq b_i \quad \forall i \\
 & && -y_{ij} \leq x_j - z_{ij} \leq y_{ij} \quad \forall i, j \in J_i \\
 & && \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \\
 & && \mathbf{y} \geq \mathbf{0}.
 \end{aligned} \tag{3}$$

Under the model of data uncertainty  $\mathbf{U}$ , the authors have shown that the probability that the  $i$  constraint is violated is at most  $\exp(-\Omega_i^2/2)$ . The robust Model (3) is less conservative than Model (2), as every feasible solution of the latter problem is a feasible solution to the former problem.

We next examine the sizes of Formulations (2) and (3). We assume that there are  $k$  coefficients of the  $m \times n$  nominal matrix  $\mathbf{A}$  that are subject to uncertainty. Given that the original nominal problem has  $n$  variables and  $m$  constraints (not counting the bound constraints), Model (2) is a linear optimization problem with  $2n$  variables, and  $m + 2n$  constraints. In contrast, Model (3) is a second-order cone problem, with  $n + 2k$  variables and  $m + 2k$  constraints. Since Model (3) is a nonlinear one, it is not particularly attractive for solving robust discrete optimization models.

### 3. The New Robust Approach

In this section, we propose a robust formulation that is linear, is able to withstand parameter uncertainty under the model of data uncertainty  $\mathbf{U}$  without excessively affecting the objective function, and readily extends to discrete optimization problems.

We motivate the formulation as follows. Consider the  $i$ th constraint of the nominal problem  $\mathbf{a}_i' \mathbf{x} \leq b_i$ . Let  $J_i$  be the set of coefficients  $a_{ij}$ ,  $j \in J_i$  that are subject to parameter uncertainty; i.e.,  $\tilde{a}_{ij}$ ,  $j \in J_i$  takes values according to a symmetric distribution with mean equal to the nominal value  $a_{ij}$  in the interval  $[a_{ij} - \hat{a}_{ij}, a_{ij} + \hat{a}_{ij}]$ . For every  $i$ , we introduce a parameter  $\Gamma_i$ , not necessarily integer, that takes values in the interval  $[0, |J_i|]$ . As would become clear below, the role of the parameter  $\Gamma_i$  is to adjust the robustness of the proposed method against the level of conservatism of the solution. Speaking intuitively, it is unlikely that all of the  $a_{ij}$ ,  $j \in J_i$  will change. Our goal is to be protected against all cases that up to  $\lfloor \Gamma_i \rfloor$  of these coefficients are allowed to change, and one coefficient  $a_{it}$  changes by  $(\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{it}$ . In other words, we stipulate that nature will be restricted in its behavior, in that only a subset of the coefficients will change in order to adversely affect the solution. We will develop an approach that has the property that if nature behaves like this, then the robust solution will

be feasible *deterministically*, and moreover, even if more than  $\lfloor \Gamma_i \rfloor$  change, then the robust solution will be feasible *with very high probability*.

We consider the following (still nonlinear) formulation:

$$\begin{aligned}
 & \text{maximize} && \mathbf{c}'\mathbf{x} \\
 & \text{subject to} && \sum_j a_{ij}x_j \\
 & && + \max_{\{S_i \cup \{t_i\} | S_i \subseteq J_i, |S_i| = \lfloor \Gamma_i \rfloor, t_i \in J_i \setminus S_i\}} \left\{ \sum_{j \in S_i} \hat{a}_{ij}y_j + (\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{it_i}y_{t_i} \right\} \\
 & && \leq b_i \quad \forall i \\
 & && -y_j \leq x_j \leq y_j \quad \forall j \\
 & && \mathbf{l} \leq \mathbf{x} \leq \mathbf{u} \\
 & && \mathbf{y} \geq \mathbf{0}.
 \end{aligned} \tag{4}$$

If  $\Gamma_i$  is chosen as an integer, the  $i$ th constraint is protected by  $\beta_i(\mathbf{x}, \Gamma_i) = \max_{\{S_i | S_i \subseteq J_i, |S_i| = \Gamma_i\}} \{\sum_{j \in S_i} \hat{a}_{ij}|x_j|\}$ . Note that when  $\Gamma_i = 0$ ,  $\beta_i(\mathbf{x}, \Gamma_i) = 0$ , the constraints are equivalent to that of the nominal problem. Likewise, if  $\Gamma_i = |J_i|$ , we have Soyster's method. Therefore, by varying  $\Gamma_i \in [0, |J_i|]$ , we have the flexibility of adjusting the robustness of the method against the level of conservatism of the solution.

In order to reformulate Model (4) as a linear optimization model we need the following proposition.

**PROPOSITION 1.** *Given a vector  $\mathbf{x}^*$ , the protection function of the  $i$ th constraint,*

$$\begin{aligned}
 & \beta_i(\mathbf{x}^*, \Gamma_i) \\
 & = \max_{\{S_i \cup \{t_i\} | S_i \subseteq J_i, |S_i| = \lfloor \Gamma_i \rfloor, t_i \in J_i \setminus S_i\}} \left\{ \sum_{j \in S_i} \hat{a}_{ij}|x_j^*| + (\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{it_i}|x_{t_i}^*| \right\},
 \end{aligned} \tag{5}$$

*equals the objective function of the following linear optimization problem:*

$$\begin{aligned}
 & \beta_i(\mathbf{x}^*, \Gamma_i) = \text{maximize} && \sum_{j \in J_i} \hat{a}_{ij}|x_j^*|z_{ij} \\
 & \text{subject to} && \sum_{j \in J_i} z_{ij} \leq \Gamma_i \\
 & && 0 \leq z_{ij} \leq 1 \quad \forall j \in J_i.
 \end{aligned} \tag{6}$$

**PROOF.** Clearly the optimal solution value of Problem (6) consists of  $\lfloor \Gamma_i \rfloor$  variables at 1 and one variable at  $\Gamma_i - \lfloor \Gamma_i \rfloor$ . This is equivalent to the selection of subset  $\{S_i \cup \{t_i\} | S_i \subseteq J_i, |S_i| = \lfloor \Gamma_i \rfloor, t_i \in J_i \setminus S_i\}$  with corresponding cost function  $\sum_{j \in S_i} \hat{a}_{ij}|x_j^*| + (\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{it_i}|x_{t_i}^*|$ .  $\square$

We next reformulate Model (4) as a linear optimization model.

**THEOREM 1.** *Model (4) has an equivalent linear formulation as follows:*

$$\begin{aligned}
& \text{maximize} && \mathbf{c}'\mathbf{x} \\
& \text{subject to} && \sum_j a_{ij}x_j + z_i\Gamma_i + \sum_{j \in J_i} p_{ij} \leq b_i \quad \forall i \\
& && z_i + p_{ij} \geq \hat{a}_{ij}y_j \quad \forall i, j \in J_i \\
& && -y_j \leq x_j \leq y_j \quad \forall j \\
& && l_j \leq x_j \leq u_j \quad \forall j \\
& && p_{ij} \geq 0 \quad \forall i, j \in J_i \\
& && y_j \geq 0 \quad \forall j \\
& && z_i \geq 0 \quad \forall i.
\end{aligned} \tag{7}$$

**PROOF.** We first consider the dual of Problem (6):

$$\begin{aligned}
& \text{minimize} && \sum_{j \in J_i} p_{ij} + \Gamma_i z_i \\
& \text{subject to} && z_i + p_{ij} \geq \hat{a}_{ij}|x_j^*| \quad \forall i, j \in J_i \\
& && p_{ij} \geq 0 \quad \forall j \in J_i \\
& && z_i \geq 0 \quad \forall i.
\end{aligned} \tag{8}$$

By strong duality, since Problem (6) is feasible and bounded for all  $\Gamma_i \in [0, |J_i|]$ , then the dual problem (8) is also feasible and bounded and their objective values coincide. Using Proposition 1, we have that  $\beta_i(\mathbf{x}^*, \Gamma_i)$  is equal to the objective function value of Problem (8). Substituting to Problem (4), we obtain that Problem (4) is equivalent to the linear optimization problem (7).  $\square$

**REMARK.** The robust linear optimization Model (7) has  $n + k + 1$  variables and  $m + k + n$  constraints, where  $k = \sum_i |J_i|$  the number of uncertain data, contrasted with  $n + 2k$  variables and  $m + 2k$  constraints for the nonlinear Formulation (3). In most real-world applications, the matrix  $\mathbf{A}$  is sparse. An attractive characteristic of Formulation (7) is that it preserves the sparsity of the matrix  $\mathbf{A}$ .

#### 4. Probability Bounds of Constraint Violation

It is clear by the construction of the robust formulation that if up to  $\lfloor \Gamma_i \rfloor$  of the  $J_i$  coefficients  $a_{ij}$  change within their bounds, and up to one coefficient  $a_{i_i}$  changes by  $(\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{i_i}$ , then the solution of Problem (7) will remain feasible. In this section, we show that under the Model of Data Uncertainty  $\mathbf{U}$ , the robust solution is feasible with high probability. The parameter  $\Gamma_i$  controls the trade-off between the probability of violation and the effect to the objective function of the nominal problem, which is what we call *the price of robustness*.

In preparation for our main result in this section, we prove the following proposition.

**PROPOSITION 2.** *Let  $\mathbf{x}^*$  be an optimal solution of Problem (7). Let  $S_i^*$  and  $t_i^*$  be the set and the index, respectively, that achieve the maximum for  $\beta_i(\mathbf{x}^*, \Gamma_i)$  in Equation (5). Suppose that the data in matrix  $\mathbf{A}$  are subjected to the model of data uncertainty  $\mathbf{U}$ .*

(a) *The probability that the  $i$ th constraint is violated satisfies:*

$$\Pr\left(\sum_j \tilde{a}_{ij}x_j^* > b_i\right) \leq \Pr\left(\sum_{j \in J_i} \gamma_{ij}\eta_{ij} \geq \Gamma_i\right),$$

where

$$\gamma_{ij} = \begin{cases} 1, & \text{if } j \in S_i^* \\ \frac{\hat{a}_{ij}|x_j^*|}{\hat{a}_{i_{r^*}}|x_{r^*}^*|}, & \text{if } j \in J_i \setminus S_i^* \end{cases}$$

and

$$r^* = \arg \min_{r \in S_i^* \cup \{t_i^*\}} \hat{a}_{i_r}|x_{r^*}^*|.$$

(b) *The quantities  $\gamma_{ij}$  satisfy  $\gamma_{ij} \leq 1$  for all  $j \in J_i \setminus S_i^*$ .*

**PROOF**

(a) Let  $\mathbf{x}^*$ ,  $S_i^*$ , and  $t_i^*$  be the solution of Model (4). Then the probability of violation of the  $i$ th constraint is as follows:

$$\begin{aligned}
& \Pr\left(\sum_j \tilde{a}_{ij}x_j^* > b_i\right) \\
& = \Pr\left(\sum_j a_{ij}x_j^* + \sum_{j \in J_i} \eta_{ij}\hat{a}_{ij}x_j^* > b_i\right) \\
& \leq \Pr\left(\sum_{j \in J_i} \eta_{ij}\hat{a}_{ij}|x_j^*| > \sum_{j \in S_i^*} \hat{a}_{ij}|x_j^*| + (\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{i_{t_i^*}}|x_{t_i^*}^*|\right) \tag{9} \\
& = \Pr\left(\sum_{j \in J_i \setminus S_i^*} \eta_{ij}\hat{a}_{ij}|x_j^*| > \sum_{j \in S_i^*} \hat{a}_{ij}|x_j^*|(1 - \eta_{ij}) \right. \\
& \quad \left. + (\Gamma_i - \lfloor \Gamma_i \rfloor)\hat{a}_{i_{t_i^*}}|x_{t_i^*}^*|\right) \\
& \leq \Pr\left(\sum_{j \in J_i \setminus S_i^*} \eta_{ij}\hat{a}_{ij}|x_j^*| > \hat{a}_{i_{r^*}}|x_{r^*}^*| \right. \\
& \quad \left. \cdot \left(\sum_{j \in S_i^*} (1 - \eta_{ij}) + (\Gamma_i - \lfloor \Gamma_i \rfloor)\right)\right) \tag{10} \\
& = \Pr\left(\sum_{j \in S_i^*} \eta_{ij} + \sum_{j \in J_i \setminus S_i^*} \frac{\hat{a}_{ij}|x_j^*|}{\hat{a}_{i_{r^*}}|x_{r^*}^*|} \eta_{ij} > \Gamma_i\right) \\
& = \Pr\left(\sum_{j \in J_i} \gamma_{ij}\eta_{ij} > \Gamma_i\right) \\
& \leq \Pr\left(\sum_{j \in J_i} \gamma_{ij}\eta_{ij} \geq \Gamma_i\right).
\end{aligned}$$

Inequality (9) follows from Inequality (4), since  $\mathbf{x}^*$  satisfies

$$\sum_j a_{ij}x_j^* + \sum_{j \in S_i^*} \hat{a}_{ij}y_j + (\Gamma_i - \lfloor \Gamma_i \rfloor) \hat{a}_{it_i^*} y_{t_i^*} \leq b_i.$$

Inequality (10) follows from  $1 - \eta_{ij} \geq 0$  and  $r^* = \arg \min_{r \in S_i^* \cup \{t_i^*\}} \hat{a}_{ir} |x_r^*|$ .

(b) Suppose there exist  $l \in J_i \setminus S_i^*$  such that  $\hat{a}_{il} |x_l^*| > \hat{a}_{ir^*} |x_{r^*}^*|$ . If  $l \neq t_i^*$ , then, since  $r^* \in S_i^* \cup \{t_i^*\}$ , we could increase the objective value of  $\sum_{j \in S_i^*} \hat{a}_{ij} |x_j^*| + (\Gamma_i - \lfloor \Gamma_i \rfloor) \hat{a}_{it_i^*} |x_{t_i^*}^*|$  by exchanging  $l$  with  $r^*$  from the set  $S_i^* \cup \{t_i^*\}$ . Likewise, if  $l = t_i^*$ ,  $r^* \in S_i^*$ , we could exchange  $t_i^*$  with  $r^*$  in the set  $S_i^*$  to increase the same objective function. In both cases, we arrive at a contradiction that  $S_i^* \cup \{t_i^*\}$  is an optimum solution to this objective function.  $\square$

We are naturally led to bound the probability

$$\Pr\left(\sum_{j \in J_i} \gamma_{ij} \eta_{ij} \geq \Gamma_i\right).$$

The next result provides a bound that is independent of the solution  $\mathbf{x}^*$ .

**THEOREM 2.** *If  $\eta_{ij}$ ,  $j \in J_i$  are independent and symmetrically distributed random variables in  $[-1, 1]$ , then*

$$\Pr\left(\sum_{j \in J_i} \gamma_{ij} \eta_{ij} \geq \Gamma_i\right) \leq \exp\left(-\frac{\Gamma_i^2}{2|J_i|}\right). \quad (11)$$

**PROOF.** Let  $\theta > 0$ . Then

$$\Pr\left(\sum_{j \in J_i} \gamma_{ij} \eta_{ij} \geq \Gamma_i\right) \leq \frac{E[\exp(\theta \sum_{j \in J_i} \gamma_{ij} \eta_{ij})]}{\exp(\theta \Gamma_i)} \quad (12)$$

$$= \frac{\prod_{j \in J_i} E[\exp(\theta \gamma_{ij} \eta_{ij})]}{\exp(\theta \Gamma_i)} \quad (13)$$

$$= \frac{\prod_{j \in J_i} 2 \int_0^1 \sum_{k=0}^{\infty} ((\theta \gamma_{ij} \eta)^{2k} / (2k)!) dF_{\eta_{ij}}(\eta)}{\exp(\theta \Gamma_i)} \quad (14)$$

$$\leq \frac{\prod_{j \in J_i} \sum_{k=0}^{\infty} ((\theta \gamma_{ij})^{2k} / (2k)!) \leq \frac{\prod_{j \in J_i} \exp(\theta^2 \gamma_{ij}^2 / 2)}{\exp(\theta \Gamma_i)} \leq \exp\left(|J_i| \frac{\theta^2}{2} - \theta \Gamma_i\right). \quad (15)$$

Inequality (12) follows from Markov's inequality, Equations (13) and (14) follow from the independence and symmetric distribution assumption of the random variables  $\eta_{ij}$ . Inequality (15) follows from  $\gamma_{ij} \leq 1$ . Selecting  $\theta = \Gamma_i / |J_i|$ , we obtain (11).  $\square$

**REMARK.** While the bound we established has the attractive feature that is independent of the solution  $\mathbf{x}^*$ , it is not particularly attractive, especially when  $\Gamma_i^2 / (2|J_i|)$  is small. We next derive the best possible bound, i.e., a bound that is achievable. We assume that  $\Gamma_i \geq 1$ .

**THEOREM 3**

(a) *If  $\eta_{ij}$ ,  $j \in J_i$  are independent and symmetrically distributed random variables in  $[-1, 1]$ , then*

$$\Pr\left(\sum_{j \in J_i} \gamma_{ij} \eta_{ij} \geq \Gamma_i\right) \leq B(n, \Gamma_i), \quad (16)$$

where

$$B(n, \Gamma_i) = \frac{1}{2^n} \left\{ (1 - \mu) \sum_{l=\lfloor \nu \rfloor}^n \binom{n}{l} + \mu \sum_{l=\lfloor \nu \rfloor + 1}^n \binom{n}{l} \right\} \\ = \frac{1}{2^n} \left\{ (1 - \mu) \binom{n}{\lfloor \nu \rfloor} + \sum_{l=\lfloor \nu \rfloor + 1}^n \binom{n}{l} \right\}, \quad (17)$$

where  $n = |J_i|$ ,  $\nu = (\Gamma_i + n)/2$ , and  $\mu = \nu - \lfloor \nu \rfloor$ .

(b) *The bound (16) is tight for  $\eta_{ij}$  having a discrete probability distribution:  $\Pr(\eta_{ij} = 1) = 1/2$  and  $\Pr(\eta_{ij} = -1) = 1/2$ ,  $\gamma_{ij} = 1$ , an integral value of  $\Gamma_i \geq 1$ , and  $\Gamma_i + n$  being even.*

(c) *The bound (16) satisfies*

$$B(n, \Gamma_i) \leq (1 - \mu) C(n, \lfloor \nu \rfloor) + \sum_{l=\lfloor \nu \rfloor + 1}^n C(n, l), \quad (18)$$

where

$$C(n, l) = \begin{cases} \frac{1}{2^n}, & \text{if } l = 0 \text{ or } l = n, \\ \frac{1}{\sqrt{2\pi}} \sqrt{\frac{n}{(n-l)l}} \\ \cdot \exp\left(n \log\left(\frac{n}{2(n-l)}\right) + l \log\left(\frac{n-l}{l}\right)\right), & \text{otherwise.} \end{cases} \quad (19)$$

(d) *For  $\Gamma_i = \theta \sqrt{n}$ ,*

$$\lim_{n \rightarrow \infty} B(n, \Gamma_i) = 1 - \Phi(\theta), \quad (20)$$

where


$$\Phi(\theta) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\theta} \exp\left(-\frac{y^2}{2}\right) dy$$

is the cumulative distribution function of a standard normal.

**PROOF.** See appendix (§8).

**REMARKS**

(a) While Bound (16) is the best possible (Theorem 3(b)), it poses computational difficulties in evaluating the sum of combination functions for large  $n$ . For this reason, we have calculated Bound (18), which is simple to compute and, as we will see, also very tight.

**Figure 1.** Comparison of probability bounds for  $n = |J_i| = 10$ .

(b) Equation (20) is a formal asymptotic theorem that applies when  $\Gamma_i = \theta\sqrt{n}$ . We can use the De Moivre-Laplace approximation of the binomial distribution to obtain the approximation

$$B(n, \Gamma_i) \approx 1 - \Phi\left(\frac{\Gamma_i - 1}{\sqrt{n}}\right), \quad (21)$$

that applies even when  $\Gamma_i$  does not scale as  $\theta\sqrt{n}$ .

(c) We next compare the bounds: (11) (Bound 1), (16) (Bound 2), (18) (Bound 3), and the approximate bound (21) for  $n = |J_i| = 10, 100, 2,000$ . In Figure 1 we compare Bounds 1 and 2 for  $n = 10$ , which clearly shows that Bound 2 dominates Bound 1 (in this case there is no need to calculate Bound 3 and the approximate bound as  $n$  is small). In Figure 2 we compare all bounds for  $n = 100$ . It is clear that Bound 3, which is simple to compute, is identical to Bound 2, and both Bounds 2 and 3 dominate Bound 1 by an order of magnitude. The approximate bound provides a reasonable approximation to Bound 2. In Figure 3 we compare Bounds 1 and 3 and the approximate bound for  $n = 2,000$ . Bound 3 is identical to the approximate bound, and both dominate Bound 1 by an order of magnitude. In summary, in the remainder of the paper we will use Bound 3, as it is simple to compute, it is a true bound (as opposed to the approximate bound), and it dominates Bound 1. To amplify this point, Table 1 illustrates the choice of  $\Gamma_i$  as a


function of  $n = |J_i|$  so that the probability that a constraint is violated is less than 1%, where we used Bounds 1, 2, 3, and the approximate bound to evaluate the probability. It is clear that using Bounds 2, 3, or the approximate bound gives essentially identical values of  $\Gamma_i$ , while using Bound 1 leads to unnecessarily higher values of  $\Gamma_i$ . For  $|J_i| = 200$ , we need to use  $\Gamma = 33.9$ , i.e., only 17% of the number of uncertain data, to guarantee violation probability of less than 1%. For constraints with a lower number of uncertain data, such as  $|J_i| = 5$ , it is necessary to ensure full protection, which is equivalent to Soyster's method. Clearly, for constraints with a large number of uncertain data, the proposed approach is capable of delivering less conservative solutions compared to Soyster's method.

#### 4.1. On the Conservatism of Robust Solutions


We have argued so far that the linear optimization framework of our approach has some computational advantages over the conic quadratic framework of Ben-Tal and Nemirovski (1998, 1999, 2000), El-Ghaoui et al. (1998), and El-Ghaoui and Lebret (1997), especially with respect to discrete optimization problems. Our objective in this section is to provide some insight, but not conclusive evidence, on the degree of conservatism for both approaches.

Given a constraint  $\mathbf{a}'\mathbf{x} \leq b$ , with  $\mathbf{a} \in [\bar{\mathbf{a}} - \hat{\mathbf{a}}, \bar{\mathbf{a}} + \hat{\mathbf{a}}]$ , the robust counterpart of Ben-Tal and Nemirovski (1998, 1999, 2000), El-Ghaoui et al. (1998), and El-Ghaoui and

**Figure 2.** Comparison of probability bounds for  $n = |J_i| = 100$ .



**Figure 3.** Comparison of probability bounds for  $n = |J_i| = 2,000$ .



**Table 1.** Choice of  $\Gamma_i$  as a function of  $n = |J_i|$  so that the probability of constraint violation is less than 1%.

$ J_i $	$\Gamma_i$ from Bound 1	$\Gamma_i$ from Bounds 2, 3	$\Gamma_i$ from Approx.
5	5	5	5
10	9.6	8.2	8.4
100	30.3	24.3	24.3
200	42.9	33.9	33.9
2,000	135.7	105	105

Le Bret (1998) in its simplest form of ellipsoidal uncertainty (Formulation (3) includes combined interval and ellipsoidal uncertainty) is:

$$\bar{\mathbf{a}}'\mathbf{x} + \Omega\|\hat{\mathbf{A}}\mathbf{x}\| \leq b,$$

where  $\hat{\mathbf{A}}$  is a diagonal matrix with elements  $\hat{a}_i$  in the diagonal. Ben-Tal and Nemirovski (2000) show that under the model of data uncertainty  $\mathbf{U}$  for  $\mathbf{a}$ , the probability that the constraint is violated is bounded above by  $\exp(-\Omega^2/2)$ .

The robust counterpart of the current approach is

$$\bar{\mathbf{a}}'\mathbf{x} + \beta(\mathbf{x}, \Gamma) \leq b,$$

where we assumed that  $\Gamma$  is integral and

$$\beta(\mathbf{x}, \Gamma) = \max_{S, |S|=\Gamma} \sum_{i \in S} \hat{a}_i |x_i|.$$

From Equation (11), the probability that the constraint is violated under the model of data uncertainty  $\mathbf{U}$  for  $\mathbf{a}$  is bounded above by  $\exp(-\Gamma^2/(2n))$ . Note that we do not use the stronger bound (16) for simplicity.

Let us select  $\Gamma = \Omega\sqrt{n}$  so that the bounds for the probability of violation are the same for both approaches. The protection levels are  $\Omega\|\hat{\mathbf{A}}\mathbf{x}\|$  and  $\beta(\mathbf{x}, \Gamma)$ . We will compare the protection levels both from a worst- and an average-case point of view in order to obtain some insight on the degree of conservatism. To simplify the exposition we define  $y_i = \hat{a}_i |x_i|$ . We also assume without loss of generality that  $y_1 \geq y_2 \geq \dots \geq y_n \geq 0$ . Then the two protection levels become  $\Omega\|\mathbf{y}\|$  and  $\sum_{i=1}^{\Gamma} y_i$ .

For  $\Gamma = \theta\sqrt{n}$ , and  $y_1^0 = \dots = y_{\Gamma}^0 = 1$ ,  $y_k^0 = 0$  for  $k \geq \Gamma + 1$ , we have  $\sum_{i=1}^{\Gamma} y_i^0 = \Gamma = \theta\sqrt{n}$ , while  $\Omega\|\mathbf{y}\| = \theta^{3/2}n^{1/4}$ ; i.e., in this example the protection level of the conic quadratic framework is asymptotically smaller than our framework by a multiplicative factor of  $n^{1/4}$ . This order of the magnitude is, in fact, worst possible, since

$$\sum_{i=1}^{\Gamma} y_i \leq \sqrt{\Gamma}\|\mathbf{y}\| = \sqrt{\frac{n}{\Gamma}}(\Omega\|\mathbf{y}\|),$$

which for  $\Gamma = \theta\sqrt{n}$  leads to

$$\sum_{i=1}^{\Gamma} y_i \leq \frac{n^{1/4}}{\theta^{1/2}}(\Omega\|\mathbf{y}\|).$$

Moreover, we have

$$\begin{aligned} \Omega\|\mathbf{y}\| &\leq \Omega \sqrt{\sum_{i=1}^{\Gamma} y_i^2 + y_{\Gamma}^2(n - \Gamma)} \\ &\leq \Omega \sqrt{\sum_{i=1}^{\Gamma} y_i^2 + \sum_{i=1}^{\Gamma} y_i^2 \left(\frac{n - \Gamma}{\Gamma^2}\right)} \\ &= \frac{\Gamma}{\sqrt{n}} \sqrt{\sum_{i=1}^{\Gamma} y_i^2 \sqrt{1 + \frac{n - \Gamma}{\Gamma^2}}} \\ &\leq \sqrt{\frac{\Gamma^2 + n - \Gamma}{n}} \sum_{i=1}^{\Gamma} y_i. \end{aligned}$$

If we select  $\Gamma = \theta\sqrt{n}$ , which makes the probability of violation  $\exp(-\theta^2/2)$ , we obtain that

$$\Omega\|\mathbf{y}\| \leq \sqrt{1 + \theta^2} \sum_{i=1}^{\Gamma} y_i.$$

Thus, in the worst case the protection level of our framework can only be smaller than the conic quadratic framework by a multiplicative factor of a constant. We conclude that in the worst case the protection level for the conic quadratic framework can be smaller than our framework by a factor of  $n^{1/4}$ , while the protection of our framework can be smaller than the conic quadratic framework by at most a constant.

Let us compare the protection levels on average, however. In order to obtain some insight, let us assume that  $y_i$  are independently and uniformly distributed in  $[0, 1]$ . Simple calculations show that for the case in question ( $\Omega = \Gamma/\sqrt{n}$ ,  $\Gamma = \theta\sqrt{n}$ )

$$E[\Omega\|\mathbf{y}\|] = \Theta(\sqrt{n}), \quad E\left[\max_{S, |S|=\Gamma} \sum_{i \in S} y_i\right] = \Theta(\sqrt{n}),$$

which implies that on average the two protection levels are of the same order of magnitude.

It is admittedly unclear whether it is the worst or the average case we presented which is more relevant, and thus the previous discussion is inconclusive. It is fair to say, however, that both approaches allow control of the degree of conservatism by adjusting the parameters  $\Gamma$  and  $\Omega$ . Moreover, we think that the ultimate criterion for comparing the degree of conservatism of these methods will be computation in real problems.

## 4.2. Local Sensitivity Analysis of the Protection Level

Given the solution of Problem (7), it is desirable to estimate the change in the objective function value with respect to the change of the protection level  $\Gamma_i$ . In this way we can assess the price of increasing or decreasing the protection level  $\Gamma_i$  of any constraint. Note that when  $\Gamma_i$  changes, only one parameter in the coefficient matrix in Problem (7)

changes. Thus, we can use results from sensitivity analysis (see Freund 1985 for a comprehensive analysis) to understand the effect of changing the protection level under nondegeneracy assumptions.

**THEOREM 4.** *Let  $\mathbf{z}^*$  and  $\mathbf{q}^*$  be the optimal nondegenerate primal and dual solutions for the linear optimization Problem (7) (under nondegeneracy, the primal and dual optimal solutions are unique). Then, the derivative of the objective function value with respect to protection level  $\Gamma_i$  of the  $i$ th constraint is*

$$-z_i^* q_i^*, \quad (22)$$

where  $z_i^*$  is the optimal primal variable corresponding to the protection level  $\Gamma_i$  and  $q_i^*$  is the optimal dual variable of the  $i$ th constraint.

**PROOF.** We transform Problem (7) in standard form,

$$\begin{aligned} G(\Gamma_i) = \text{maximize} \quad & \mathbf{c}'\mathbf{x} \\ \text{subject to} \quad & \mathbf{A}\mathbf{x} + \Gamma_i z_i \mathbf{e}_i = \mathbf{b} \\ & \mathbf{x} \geq \mathbf{0}, \end{aligned}$$

where  $\mathbf{e}_i$  is a unit vector with a one in the  $i$ th position. Let  $\mathbf{B}$  be the optimal basis, which is unique under primal and dual nondegeneracy. If the column  $\Gamma_i \mathbf{e}_i$  corresponding to the variable  $z_i$  is not in the basis, then  $z_i^* = 0$ . In this case, under dual nondegeneracy all reduced costs associated with the nonbasic variables are strictly negative, and thus a marginal change in the protection level does not affect the objective function value. Equation (22) correctly indicates the zero variation.

If the column  $\Gamma_i \mathbf{e}_i$  corresponding to the variable  $z_i$  is in the basis, and the protection level  $\Gamma_i$  changes by  $\Delta\Gamma_i$ , then  $\mathbf{B}$  becomes  $\mathbf{B} + \Delta\Gamma_i \mathbf{e}_i \mathbf{e}_i'$ . By the *Matrix Inversion Lemma* we have:

$$(\mathbf{B} + \Delta\Gamma_i \mathbf{e}_i \mathbf{e}_i')^{-1} = \mathbf{B}^{-1} - \frac{\Delta\Gamma_i \mathbf{B}^{-1} \mathbf{e}_i \mathbf{e}_i' \mathbf{B}^{-1}}{1 + \Delta\Gamma_i \mathbf{e}_i' \mathbf{B}^{-1} \mathbf{e}_i}.$$

Under primal and dual nondegeneracy, for small changes  $\Delta\Gamma_i$ , the new solutions preserve primal and dual feasibility. Therefore, the corresponding change in the objective function value is,

$$\begin{aligned} G(\Gamma_i + \Delta\Gamma_i) - G(\Gamma_i) &= -\frac{\Delta\Gamma_i \mathbf{c}_b' \mathbf{B}^{-1} \mathbf{e}_i \mathbf{e}_i' \mathbf{B}^{-1} \mathbf{b}}{1 + \Delta\Gamma_i \mathbf{e}_i' \mathbf{B}^{-1} \mathbf{e}_i} \\ &= -\frac{\Delta\Gamma_i z_i^* q_i^*}{1 + \Delta\Gamma_i \mathbf{e}_i' \mathbf{B}^{-1} \mathbf{e}_i}, \end{aligned}$$

where  $\mathbf{c}_b$  is the part of the vector  $\mathbf{c}$  corresponding to the columns in  $\mathbf{B}$ . Thus,

$$G'(\Gamma_i) = \lim_{\Delta\Gamma_i \rightarrow 0} \frac{G(\Gamma_i + \Delta\Gamma_i) - G(\Gamma_i)}{\Delta\Gamma_i} = -z_i^* q_i^*. \quad \square$$

**REMARK.** An attractive aspect of Equation (22) is its simplicity, as it only involves the primal optimal solution corresponding to the protection level  $\Gamma_i$  and the dual optimal solution corresponding to the  $i$ th constraint.

## 5. Correlated Data

So far we have assumed that the data are independently uncertain. It is possible, however, that the data are correlated. In particular, we envision that there are few sources of data uncertainty that affect all the data. More precisely, we assume that the model of data uncertainty is as follows.

**Correlated Model of Data Uncertainty C.** Consider a particular row  $i$  of the matrix  $\mathbf{A}$  and let  $J_i$  the set of coefficients in row  $i$  that are subject to uncertainty. Each entry  $a_{ij}$ ,  $j \in J_i$  is modeled as

$$\tilde{a}_{ij} = a_{ij} + \sum_{k \in K_i} \tilde{\eta}_{ik} g_{kj}$$

and  $\tilde{\eta}_{ik}$  are independent and symmetrically distributed random variables in  $[-1, 1]$ .

Note that under this model there are only  $|K_i|$  sources of data uncertainty that affect the data in row  $i$ . Note that these sources of uncertainty affect all the entries  $a_{ij}$ ,  $j \in J_i$ . For example, if  $|K_i| = 1$ , then all data in a row are affected by a single random variable. For a concrete example, consider a portfolio construction problem in which returns of various assets are predicted from a regression model. In this case, there are a few sources of uncertainty that globally affect all the assets classes.

Analogously to (4), we propose the following robust formulation:

$$\begin{aligned} \text{maximize} \quad & \mathbf{c}'\mathbf{x} \\ \text{subject to} \quad & \sum_j a_{ij} x_j + \max_{\{S_i \cup \{t_i\} | S_i \subseteq K_i, |S_i| = \lfloor \Gamma_i \rfloor, t_i \in K_i \setminus S_i\}} \left\{ \sum_{k \in S_i} \left| \sum_{j \in J_i} g_{kj} x_j \right| \right. \\ & \left. + (\Gamma_i - \lfloor \Gamma_i \rfloor) \left| \sum_{j \in J_i} g_{t_i j} x_j \right| \right\} \leq b_i \quad \forall i \\ & \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}, \end{aligned} \quad (23)$$

which can be written as a linear optimization problem as follows:

$$\begin{aligned} \text{maximize} \quad & \mathbf{c}'\mathbf{x} \\ \text{subject to} \quad & \sum_j a_{ij} x_j + z_i \Gamma_i + \sum_{k \in K_i} p_{ik} \leq b_i \quad \forall i \\ & z_i + p_{ik} \geq y_{ik} \quad \forall i, k \in K_i \\ & -y_{ik} \leq \sum_{j \in J_i} g_{kj} x_j \leq y_{ik} \quad \forall i, k \in K_i \\ & l_j \leq x_j \leq u_j \quad \forall j \\ & p_{ik}, y_{ik} \geq 0 \quad \forall i, k \in K_i \\ & z_i \geq 0 \quad \forall i. \end{aligned} \quad (24)$$

Analogously to Theorem 3, we can show that the probability that the  $i$ th constraint is violated,  $k$  is at most  $B(|K_i|, \Gamma_i)$ , defined in Equation (16).

## 6. Experimental Results

In this section, we present three experiments illustrating our robust solution to problems with data uncertainty. In the first experiment, we show that our methods extend to discrete optimization problems with data uncertainty, by solving a knapsack problem with uncertain weights. The second example is a simple portfolio optimization problem from Ben-Tal and Nemirovski (1999), which has data uncertainty in the objective function. In the last experiment we apply our method to a problem PILOT4 from the well-known Net Lib collection to examine the effectiveness of our approach to real-world problems.

### 6.1. The Robust Knapsack Problem

The proposed robust formulation in Theorem 1 in the case in which the nominal problem is a mixed integer programming (MIP) model—i.e., some of the variables in the vector  $\mathbf{x}$  take integer values—is still an MIP formulation, and thus can be solved in the same way that the nominal problem can be solved. Moreover, both the deterministic guarantee as well as the probabilistic guarantee (Theorem 3) that our approach provides is still valid. As a result, our approach applies for addressing data uncertainty for MIPs. To the best of our knowledge, there is no prior research in robust discrete optimization that is both tractable computationally and involves a probabilistic guarantee for constraint violation.

In this section, we apply our approach to zero-one knapsack problems that are subject to data uncertainty. Our objective in this section is to examine whether our approach is computationally tractable and whether it succeeds in reducing the price of robustness.

The zero-one knapsack problem is the following discrete optimization problem:

$$\begin{aligned} & \text{maximize} && \sum_{i \in N} c_i x_i \\ & \text{subject to} && \sum_{i \in N} w_i x_i \leq b \\ & && x_i \in \{0, 1\}. \end{aligned}$$

Although the knapsack problem is *NP-hard*, for problems of moderate size, it is often solved to optimality using state-of-the-art MIP solvers. For this experiment, we use CPLEX 6.0 to solve to optimality a random knapsack problem of size  $|N| = 200$ .

Regarding the uncertainty model for data, we assume the weights  $\tilde{w}_i$  are uncertain, independently distributed, and follow symmetric distributions in  $[w_i - \delta_i, w_i + \delta_i]$ . An application of this problem is to maximize the total value of goods to be loaded on a cargo that has strict weight restrictions. The weight of the individual item is assumed to be uncertain, independent of other weights, and follows a symmetric distribution. In our robust model, we want to maximize the total value of the goods but allow a maximum

of 1% chance of constraint violation. The robust model of Theorem 1 is as follows:

$$\begin{aligned} & \text{maximize} && \sum_{i \in N} c_i x_i \\ & \text{subject to} && DS \sum_{i \in N} w_i x_i + \zeta(x, \Gamma) \leq b \\ & && x_i \in \{0, 1\}. \end{aligned}$$

where

$$\zeta(x, \Gamma) = \max_{\{S \cup \{t\} | S \subseteq N, |S| = \lfloor \Gamma \rfloor, t \in N \setminus S\}} \left\{ \sum_{j \in S} \delta_j x_j + (\Gamma - \lfloor \Gamma \rfloor) \delta_t x_t \right\}.$$

For the random knapsack example, we set the capacity limit  $b$  to 4,000, the nominal weight  $w_i$  being randomly chosen from the set  $\{20, 21, \dots, 29\}$ , and the cost  $c_i$  randomly chosen from the set  $\{16, 17, \dots, 77\}$ . We set the weight uncertainty  $\delta_i$  to equal 10% of the nominal weight. The time to solve the robust discrete problems to optimality using CPLEX 6.0 on a Pentium II 400 PC ranges from 0.05 to 50 seconds.

Figure 4 illustrates the effect of the protection level on the objective function value. In the absence of protection to the capacity constraint, the optimal value is 5,592. However, with maximum protection, that is, admitting Soyster's (1973) method, the optimal value is reduced by 5.5% to 5,283. In Figure 5, we plot the optimal value with respect to the approximate probability bound of constraint violation. In Table 2, we present a sample of the objective function value and the probability bound of constraint violation.


It is interesting to note that the optimal value is marginally affected when we increase the protection level. For instance, to have a probability guarantee of at most 0.57% chance of constraint violation, we only reduce the objective by 1.54%. We can summarize the key insights in this example:

1. Our approach succeeds in reducing the price of robustness; that is, we do not heavily penalize the objective function value in order to protect ourselves against constraint violation.
2. The proposed robust approach is computationally tractable in that the problem can be solved in reasonable computational times.


### 6.2. A Simple Portfolio Problem

In this section, we consider a portfolio construction problem consisting of a set of  $N$  stocks ( $|N| = n$ ). Stock  $i$  has return  $\tilde{p}_i$  which is of course uncertain. The objective is to determine the fraction  $x_i$  of wealth invested in stock  $i$  so as to maximize the portfolio value  $\sum_{i=1}^n \tilde{p}_i x_i$ . We model the uncertain return  $\tilde{p}_i$  as a random variable that has an arbitrary symmetric distribution in the interval  $[p_i - \sigma_i, p_i + \sigma_i]$ , where  $p_i$  is the expected return and  $\sigma_i$  is a measure of the uncertainty of the return of stock  $i$ . We further assume that the returns  $\tilde{p}_i$  are independent.

**Figure 4.** Optimal value of the robust knapsack formulation as a function of  $\Gamma$ .



**Figure 5.** Optimal value of the robust knapsack formulation as a function of the probability bound of constraint violation given in Equation (18).



**Table 2.** Results of robust knapsack solutions.

$\Gamma$	Probability Bound	Optimal Value	Reduction (%)
2.8	$4.49 \times 10^{-1}$	5,585	0.13
14.1	$1.76 \times 10^{-1}$	5,557	0.63
25.5	$4.19 \times 10^{-2}$	5,531	1.09
36.8	$5.71 \times 10^{-3}$	5,506	1.54
48.1	$4.35 \times 10^{-4}$	5,481	1.98
59.4	$1.82 \times 10^{-5}$	5,456	2.43
70.7	$4.13 \times 10^{-7}$	5,432	2.86
82.0	$5.04 \times 10^{-9}$	5,408	3.29
93.3	$3.30 \times 10^{-11}$	5,386	3.68
104.7	$1.16 \times 10^{-13}$	5,364	4.08
116.0	$2.22 \times 10^{-16}$	5,342	4.47

The classical approach in portfolio construction is to use quadratic optimization and solve the following problem:

$$\begin{aligned} & \text{maximize} && \sum_{i=1}^n p_i x_i - \phi \sum_{i=1}^n \sigma_i^2 x_i^2 \\ & \text{subject to} && \sum_{i=1}^n x_i = 1 \\ & && x_i \geq 0, \end{aligned}$$

where we interpret  $\sigma_i$  as the standard deviation of the return for stock  $i$ , and  $\phi$  is a parameter that controls the trade-off between risk and return. Applying our approach, we will solve instead the following problem (which can be reformulated as a linear optimization problem as in Theorem 7):

$$\begin{aligned} & \text{maximize} && z \\ & \text{subject to} && z \leq \sum_{i=1}^n p_i x_i - \beta(\mathbf{x}, \Gamma) \\ & && \sum_{i=1}^n x_i = 1 \\ & && x_i \geq 0, \end{aligned} \tag{25}$$

where

$$\beta(\mathbf{x}, \Gamma) = \max_{\{S \cup \{t\} | S \subseteq N, |S| = \lfloor \Gamma \rfloor, t \in N \setminus S\}} \left\{ \sum_{j \in S} \sigma_j x_j + (\Gamma - \lfloor \Gamma \rfloor) \sigma_t x_t \right\}.$$

In this setting,  $\Gamma$  is the protection level of the actual portfolio return in the following sense. Let  $\mathbf{x}^*$  be an optimal solution of Problem (25) and let  $z^*$  be the optimal solution value of Problem (25). Then,  $\mathbf{x}^*$  satisfies that  $\Pr(\tilde{\mathbf{p}}' \mathbf{x}^* < z^*)$  is less than or equal to the bound in Equation (18). Ben-Tal and Nemirovski (1999) consider the same portfolio problem using  $n = 150$ ,

$$p_i = 1.15 + i \frac{0.05}{150}, \quad \sigma_i = \frac{0.05}{450} \sqrt{2in(n+1)}.$$

Note that in this experiment, stocks with higher returns are also more risky.

**Optimization Results.** Let  $\mathbf{x}^*(\Gamma)$  be an optimal solution to Problem (25) corresponding to the protection level  $\Gamma$ . A classical measure of risk is the *standard deviation*,

$$w(\Gamma) = \sqrt{\sum_{i \in N} \sigma_i^2 (x_i^*(\Gamma))^2}.$$

We first solved Problem (25) for various levels of  $\Gamma$ . Figure 6 illustrates the performance of the robust solution as a function of the protection level  $\Gamma$ , while Figure 7 shows the solution itself for various levels of the protection level. The solution exhibits some interesting “phase transitions” as the protection level increases:

1. For  $\Gamma \leq 17$ , both the expected return as well as the risk-adjusted return (the objective function value) gradually decrease. Starting with  $\Gamma = 0$ , for which the solution consists of the stock 150 that has the highest expected return, the portfolio becomes gradually more diversified, putting more weight on stocks with higher ordinal numbers. This can be seen for example for  $\Gamma = 10$  in Figure 7.

2. For  $17 < \Gamma \leq 41$ , the risk-adjusted return continues to gradually decrease as the protection level increases, while the expected return is insensitive to the protection level. In this range,  $x_i^* = \sum_i (1/\sigma_i) / \sigma_i$ ; i.e., the portfolio is fully diversified.


3. For  $\Gamma \geq 41$ , there is a sudden phase transition (see Figure 6). The portfolio consists of only stock 1, which is the one that has the largest risk-adjusted return  $p_i - \sigma_i$ . This is exactly the solution given by the Soyster method as well. In this range, both the expected and the risk-adjusted returns are insensitive to  $\Gamma$ .

**Simulation Results.** To examine the quality of the robust solution, we run 10,000 simulations of random yields and compare robust solutions generated by varying the protection level  $\Gamma$ . As we have discussed, for the worst-case simulation, we consider the distribution with  $\tilde{p}_i$  taking with probability 1/2 the values at  $p_i \pm \sigma_i$ . In Figure 8, we compare the theoretical bound in Equation (18) with the fraction of the simulated portfolio returns falling below the optimal solution,  $z^*$ . The empirical results suggest that the theoretical bound is close to the empirically observed values.


In Table 3, we present the results of the simulation indicating the trade-off between risk and return. The corresponding plots are also presented in Figures 9 and 10. As expected, as the protection level increases, the expected and maximum returns decrease, while the minimum returns increase. For instance, with  $\Gamma \geq 15$ , the minimum return is maintained above 12% for all simulated portfolios.


This example suggests that our approach captures the trade-off between risk and return, very much like the mean variance approach, but does so in a linear framework. Additionally, the robust approach provides a deterministic guarantee about the return of the portfolio as well as a probabilistic guarantee that is valid for all symmetric distributions.

**Figure 6.** The return and the objective function value (risk-adjusted return) as a function of the protection level  $\Gamma$ .



**Figure 7.** The solution of the portfolio for various protection levels.



**Figure 8.** Simulation study of the probability of underperforming the nominal return as a function of  $\Gamma$ .

### 6.3. Robust Solutions of a Real-World Linear Optimization Problem

As noted in Ben-Tal and Nemirovski (2000), optimal solutions of linear optimization problems may become severely infeasible if the nominal data are slightly perturbed. In this experiment, we applied our method to the problem PILOT4 from the Net Lib library of problems. Problem PILOT4 is a linear optimization problem with 411 rows, 1,000 columns, 5,145 nonzero elements, and optimum objective value,  $-2,581.1392613$ . It contains coefficients such as  $717.562256$ ,  $-1.078783$ ,  $-3.053161$ ,  $-0.549569$ ,  $-22.634094$ , and  $-39.874283$ , which seem unnecessarily precise. In our study, we assume that the coefficients of this type that participate in the inequalities of the formula-

**Table 3.** Simulation results given by the robust solution.

$\Gamma$	Prob. Violation	Exp. Return	Min. Return	Max. Return	$w(\Gamma)$
0	0.5325	1.200	0.911	1.489	0.289
5	0.3720	1.184	1.093	1.287	0.025
10	0.2312	1.178	1.108	1.262	0.019
15	0.1265	1.172	1.121	1.238	0.015
20	0.0604	1.168	1.125	1.223	0.013
25	0.0250	1.168	1.125	1.223	0.013
30	0.0089	1.168	1.125	1.223	0.013
35	0.0028	1.168	1.125	1.223	0.013
40	0.0007	1.168	1.125	1.223	0.013
45	0.0002	1.150	1.127	1.174	0.024

tion have a maximum 2% deviation from the corresponding nominal values. Table 4 presents the distributions of the number of uncertain data in the problem. We highlight that each of the constraints has, at most, 29 uncertain data.

We solve the robust Problem (7) and report the results in Table 5. In Figure 11, we present the efficient frontier of the probability of constraint violation and cost.


We note that the cost of full protection (Soyster's method) is equal to  $-2,397.5798763$ . In this example, we observe that relaxing the need of full protection still leads to a high increase in the cost unless one is willing to accept unrealistically high probabilities for constraint violation. We attribute this to the fact that there are very few uncertain coefficients in each constraint (Table 4), and thus probabilistic protection is quite close to deterministic protection.

## 7. Conclusions


The major insights from our analysis are:

1. Our proposed robust methodology provides solutions that ensure deterministic and probabilistic guarantees that constraints will be satisfied as data change.
2. Under the proposed method, the protection level determines probability bounds of constraint violation, which do not depend on the solution of the robust model.
3. The method naturally applies to discrete optimization problems.

**Figure 9.** Empirical result of expected, maximum, and minimum yield.



**Figure 10.** Trade-offs between probability of underperforming and returns.



**Table 4.** Distributions of  $|J_i|$  in PILOT4.

$ J_i $	# Constraints	$ J_i $	# Constraints
1	21	24	12
11	4	25	4
14	4	26	8
15	4	27	8
16	4	28	4
17	8	29	8
22	4		

4. Comparing Figure 5, for a problem with  $n = 200$  uncertain coefficients, and Figure 11, for a problem in which the maximum number of uncertain coefficients per row is 29, we observe that in order to have probability of violation of the constraints  $10^{-4}$  we have an objective function change of 2% in the former case and 7% in the latter case. We feel that this is indicative of the fact that the attractiveness of the method increases as the number of uncertain data increases.

### 8. Appendix: Proof of Theorem 3

In this appendix we present the proof of Theorem 3.

(a) The proof follows from Proposition 2, Parts (a) and (b). To simplify the exposition, we will drop the subscript  $i$ , which represents the index of the constraint. We prove the bound in (16) by induction on  $n$ . We define the

**Table 5.** The trade-off between optimal cost and robustness.

Optimal Value	% Change	Prob. Violation
-2,486.0334864	3.68	0.5
-2,450.4323768	5.06	0.421
-2,433.4959374	5.72	0.345
-2,413.2013332	6.51	0.159
-2,403.1494574	6.90	0.0228
-2,399.2592198	7.05	0.00135
-2,397.8404967	7.10	$3.17 \times 10^{-5}$
-2,397.5798996	7.11	$2.87 \times 10^{-7}$
-2,397.5798763	7.11	$9.96 \times 10^{-8}$

auxiliary quantities:


$$\nu(\Gamma, n) = \frac{\Gamma + n}{2}, \quad \mu(\Gamma, n) = \nu(\Gamma, n) - \lfloor \nu(\Gamma, n) \rfloor,$$

$$\Upsilon(s, n) = \frac{1}{2^n} \sum_{l=s}^n \binom{n}{l}.$$

The induction hypothesis is formulated as follows:

$$\Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma\right) \leq \begin{cases} (1 - \mu(\Gamma, n))\Upsilon(\lfloor \nu(\Gamma, n) \rfloor, n) \\ + \mu(\Gamma, n)\Upsilon(\lfloor \nu(\Gamma, n) \rfloor + 1, n) & \text{if } \Gamma \in [1, n] \\ 0 & \text{if } \Gamma > n. \end{cases}$$

**Figure 11.** The trade-off between cost and robustness.



For  $n = 1$ , then  $\Gamma = 1$ , and so  $\nu(1, 1) = 1$ ,  $\mu(1, 1) = 0$ , and  $\Upsilon(1, 1) = 1/2$ , leading to:

$$\begin{aligned} \Pr(\eta_1 \geq \Gamma) &\leq \Pr(\eta_1 \geq 0) \\ &\leq \frac{1}{2} \\ &= (1 - \mu(1, 1))\Upsilon(\lfloor \nu(1, 1) \rfloor, 1) \\ &\quad + \mu(1, 1)\Upsilon(\lfloor \nu(1, 1) \rfloor + 1, 1). \end{aligned}$$

Assuming the induction hypothesis holds for  $n$ , we have

$$\begin{aligned} \Pr\left(\sum_{j=1}^{n+1} \gamma_j \eta_j \geq \Gamma\right) &= \int_{-1}^1 \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma - \gamma_{n+1} \eta_{n+1} \mid \eta_{n+1} = \eta\right) dF_{\eta_{n+1}}(\eta) \\ &= \int_{-1}^1 \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma - \gamma_{n+1} \eta\right) dF_{\eta_{n+1}}(\eta) \quad (26) \end{aligned}$$

$$\begin{aligned} &= \int_0^1 \left[ \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma - \gamma_{n+1} \eta\right) \right. \\ &\quad \left. + \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma + \gamma_{n+1} \eta\right) \right] dF_{\eta_{n+1}}(\eta) \quad (27) \end{aligned}$$

$$\begin{aligned} &\leq \max_{\phi \in [0, \gamma_{n+1}]} \left\{ \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma - \phi\right) \right. \\ &\quad \left. + \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma + \phi\right) \right\} \int_0^1 dF_{\eta_{n+1}}(\eta) \\ &= \frac{1}{2} \max_{\phi \in [0, \gamma_{n+1}]} \left\{ \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma - \phi\right) \right. \\ &\quad \left. + \Pr\left(\sum_{j=1}^n \gamma_j \eta_j \geq \Gamma + \phi\right) \right\} \end{aligned}$$

$$\leq \frac{1}{2} \max_{\phi \in [0, \gamma_{n+1}]} \Psi_n(\phi) \quad (28)$$

$$\leq \frac{1}{2} \Psi_n(1) \quad (29)$$

$$\begin{aligned} &= (1 - \mu(\Gamma, n+1))\Upsilon(\lfloor \nu(\Gamma, n+1) \rfloor, n+1) \\ &\quad + \mu(\Gamma, n+1)\Upsilon(\lfloor \nu(\Gamma, n+1) \rfloor + 1, n+1), \quad (30) \end{aligned}$$

where

$$\begin{aligned} \Psi_n(\phi) &= (1 - \mu(\Gamma - \phi, n))\Upsilon(\lfloor \nu(\Gamma - \phi, n) \rfloor, n) \\ &\quad + \mu(\Gamma - \phi, n)\Upsilon(\lfloor \nu(\Gamma - \phi, n) \rfloor + 1, n) \\ &\quad + (1 - \mu(\Gamma + \phi, n))\Upsilon(\lfloor \nu(\Gamma + \phi, n) \rfloor, n) \\ &\quad + \mu(\Gamma + \phi, n)\Upsilon(\lfloor \nu(\Gamma + \phi, n) \rfloor + 1, n). \end{aligned}$$

Equations (26) and (27) follow from the assumption that  $\eta_j$ s are independent, symmetrically distributed random variables in  $[-1, 1]$ . Inequality (28) represents the induc-

tion hypothesis. Equation (30) follows from:

$$\begin{aligned} \Psi_n(1) &= (1 - \mu(\Gamma - 1, n))\Upsilon(\lfloor \nu(\Gamma - 1, n) \rfloor, n) \\ &\quad + \mu(\Gamma - 1, n)\Upsilon(\lfloor \nu(\Gamma - 1, n) \rfloor + 1, n) \\ &\quad + (1 - \mu(\Gamma + 1, n))\Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor, n) \\ &\quad + \mu(\Gamma + 1, n)\Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor + 1, n) \quad (31) \end{aligned}$$

$$\begin{aligned} &= (1 - \mu(\Gamma + 1, n))(\Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor - 1, n) \\ &\quad + \Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor, n)) \\ &\quad + \mu(\Gamma + 1, n)(\Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor, n) \\ &\quad + \Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor + 1, n)) \quad (32) \end{aligned}$$

$$\begin{aligned} &= 2\{(1 - \mu(\Gamma + 1, n))\Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor, n+1) \\ &\quad + \mu(\Gamma + 1, n)\Upsilon(\lfloor \nu(\Gamma + 1, n) \rfloor + 1, n+1)\} \quad (33) \end{aligned}$$

$$\begin{aligned} &= 2\{(1 - \mu(\Gamma, n+1))\Upsilon(\lfloor \nu(\Gamma, n+1) \rfloor, n+1) \\ &\quad + \mu(\Gamma, n+1)\Upsilon(\lfloor \nu(\Gamma, n+1) \rfloor + 1, n+1)\}. \quad (34) \end{aligned}$$

Equations (31) and (32) follow from noting that  $\mu(\Gamma - 1, n) = \mu(\Gamma + 1, n)$  and  $\lfloor \nu(\Gamma - 1, n) \rfloor = \lfloor \nu(\Gamma + 1, n) \rfloor - 1$ . Equation (33) follows from the claim that  $\Upsilon(s, n) + \Upsilon(s + 1, n) = 2\Upsilon(s + 1, n + 1)$ , which is presented next:

$$\begin{aligned} \Upsilon(s, n) + \Upsilon(s + 1, n) &= \frac{1}{2^n} \left\{ \sum_{l=s}^n \binom{n}{l} + \sum_{l=s+1}^n \binom{n}{l} \right\} \\ &= \frac{1}{2^n} \left( \sum_{l=s}^{n-1} \left[ \binom{n}{l} + \binom{n}{l+1} \right] + 1 \right) \\ &= \frac{1}{2^n} \left( \sum_{l=s}^{n-1} \binom{n+1}{l+1} + 1 \right) \\ &= \frac{1}{2^n} \sum_{l=s+1}^{n+1} \binom{n+1}{l} \\ &= 2\Upsilon(s + 1, n + 1), \end{aligned}$$

and Equation (34) follows from  $\mu(\Gamma + 1, n) = \mu(\Gamma, n + 1) = (\Gamma + n + 1)/2$ .

We are left to show that  $\Psi_n(\phi)$  is a monotonically non-decreasing function in the domain  $\phi \in [0, 1]$ , which implies that for any  $\phi_1, \phi_2 \in [0, 1]$  such that  $\phi_1 > \phi_2$ ,  $\Psi_n(\phi_1) - \Psi_n(\phi_2) \geq 0$ . We fix  $\Gamma$  and  $n$ . To simplify the notation we use:  $\mu(\phi) = \mu(\Gamma + \phi, n) = (\Gamma + \phi + n)/2$ ,  $\nu(\phi) = \nu(\Gamma + \phi, n)$ . For any choice of  $\phi_1$  and  $\phi_2$ , we have  $\rho = \lfloor \nu_{-\phi_1} \rfloor \leq \lfloor \nu_{-\phi_2} \rfloor \leq \lfloor \nu_{\phi_2} \rfloor \leq \lfloor \nu_{\phi_1} \rfloor \leq \rho + 1$ . Therefore, we consider the following cases:

For  $\rho = \lfloor \nu_{-\phi_1} \rfloor = \lfloor \nu_{-\phi_2} \rfloor = \lfloor \nu_{\phi_2} \rfloor = \lfloor \nu_{\phi_1} \rfloor$ ,

$$\begin{aligned} \mu_{-\phi_1} - \mu_{-\phi_2} &= -\frac{\phi_1 - \phi_2}{2} \\ \mu_{\phi_1} - \mu_{\phi_2} &= \frac{\phi_1 - \phi_2}{2} \\ \Psi_n(\phi_1) - \Psi_n(\phi_2) &= \frac{\phi_1 - \phi_2}{2} \{ \Upsilon(\rho, n) - \Upsilon(\rho + 1, n) - \Upsilon(\rho, n) + \Upsilon(\rho + 1, n) \} \\ &= 0. \end{aligned}$$

For  $\rho = \lfloor \nu_{-\phi_1} \rfloor = \lfloor \nu_{-\phi_2} \rfloor = \lfloor \nu_{\phi_2} \rfloor - 1 = \lfloor \nu_{\phi_1} \rfloor - 1$ ,

$$\begin{aligned} \mu_{-\phi_1} - \mu_{-\phi_2} &= -\frac{\phi_1 - \phi_2}{2} \\ \mu_{\phi_1} - \mu_{\phi_2} &= \frac{\phi_1 - \phi_2}{2} \\ \Psi_n(\phi_1) - \Psi_n(\phi_2) &= \frac{\phi_1 - \phi_2}{2} \{ \Upsilon(\rho, n) - 2\Upsilon(\rho + 1, n) + \Upsilon(\rho + 2, n) \} \\ &= \frac{\phi_1 - \phi_2}{2^{n+1}(n+1)} \binom{n+1}{\rho+1} (1+2\rho-n) \\ &\geq \frac{\phi_1 - \phi_2}{2^{n+1}(n+1)} \binom{n+1}{\rho+1} \\ &\quad \cdot \left( 1 + 2 \left\lfloor \frac{1+n-\phi_1}{2} \right\rfloor - n \right) \\ &\geq 0. \end{aligned}$$

For  $\rho = \lfloor \nu_{-\phi_1} \rfloor = \lfloor \nu_{-\phi_2} \rfloor - 1 = \lfloor \nu_{\phi_2} \rfloor - 1 = \lfloor \nu_{\phi_1} \rfloor - 1$ ,

$$\begin{aligned} \mu_{-\phi_1} - \mu_{-\phi_2} &= -\frac{\phi_1 - \phi_2}{2} + 1 \\ \mu_{\phi_1} - \mu_{\phi_2} &= \frac{\phi_1 - \phi_2}{2} \\ \Psi_n(\phi_1) - \Psi_n(\phi_2) &= (1 - \mu_{-\phi_1}) \{ \Upsilon(\rho, n) - 2\Upsilon(\rho + 1, n) + \Upsilon(\rho + 2, n) \} \\ &\geq 0. \end{aligned}$$

For  $\rho = \lfloor \nu_{-\phi_1} \rfloor = \lfloor \nu_{-\phi_2} \rfloor = \lfloor \nu_{\phi_2} \rfloor = \lfloor \nu_{\phi_1} \rfloor - 1$ ,

$$\begin{aligned} \mu_{-\phi_1} - \mu_{-\phi_2} &= -\frac{\phi_1 - \phi_2}{2} \\ \mu_{\phi_1} - \mu_{\phi_2} &= \frac{\phi_1 - \phi_2}{2} - 1 \\ \Psi_n(\phi_1) - \Psi_n(\phi_2) &= \mu_{\phi_1} \{ \Upsilon(\rho, n) - 2\Upsilon(\rho + 1, n) + \Upsilon(\rho + 2, n) \} \\ &\geq 0. \end{aligned}$$

(b) Let  $\eta_j$  obey a discrete probability distribution  $\Pr(\eta_j = 1) = 1/2$  and  $\Pr(\eta_j = -1) = 1/2$ ,  $\gamma_j = 1$ ,  $\Gamma \geq 1$  is integral and  $\Gamma + n$  is even. Let  $S_n$  obey a binomial distribution with parameters  $n$  and  $1/2$ . Then,

$$\begin{aligned} \Pr\left(\sum_{j=1}^n \eta_j \geq \Gamma\right) &= \Pr(S_n - (n - S_n) \geq \Gamma) \\ &= \Pr(2S_n - n \geq \Gamma) \\ &= \Pr\left(S_n \geq \frac{n + \Gamma}{2}\right) \\ &= \frac{1}{2^n} \sum_{l=(n+\Gamma)/2}^n \binom{n}{l}, \end{aligned} \quad (35)$$

which implies that the bound (16) is indeed tight.

(c) From Equation (17), we need to find an upper bound for the function  $1/2^n \binom{n}{l}$ . From Robbins (1955) we obtain for  $n \geq 1$ ,

$$\begin{aligned} \sqrt{2\pi} n^{n+1/2} \exp(-n + 1/(12n + 1)) \\ \leq n! \leq \sqrt{2\pi} n^{n+1/2} \exp(-n + 1/(12n)), \end{aligned}$$

we can establish for  $l \in \{1, \dots, n-1\}$ ,

$$\begin{aligned} \frac{1}{2^n} \binom{n}{l} &= \frac{n!}{2^n (n-l)! l!} \\ &\leq \frac{1}{\sqrt{2\pi}} \sqrt{\frac{n}{(n-l)l}} \\ &\quad \cdot \exp\left(\frac{1}{12n} - \frac{1}{12(n-l)+1} - \frac{1}{12l+1}\right) \\ &\quad \cdot \left(\frac{n}{2(n-l)}\right)^n \left(\frac{n-l}{l}\right)^l \\ &\leq \frac{1}{\sqrt{2\pi}} \sqrt{\frac{n}{(n-l)l}} \left(\frac{n}{2(n-l)}\right)^n \left(\frac{n-l}{l}\right)^l \\ &= \frac{1}{\sqrt{2\pi}} \sqrt{\frac{n}{(n-l)l}} \\ &\quad \cdot \exp\left(n \log\left(\frac{n}{2(n-l)}\right) + l \log\left(\frac{n-l}{l}\right)\right), \end{aligned} \quad (36)$$

where Equation (36) follows from

$$\begin{aligned} \frac{1}{12(n-l)+1} + \frac{1}{12l+1} &\geq \frac{2}{(12(n-l)+1) + (12l+1)} \\ &= \frac{2}{12n+2} > \frac{1}{12n}. \end{aligned}$$

For  $l=0$  and  $l=n$   $\binom{n}{l} = 1/2^n$ .

(d) Bound (16) can be written as

$$B(n, \Gamma_i) = (1 - \mu) \Pr(S_n \geq \lfloor \nu \rfloor) + \mu \Pr(S_n \geq \lfloor \nu \rfloor + 1),$$

where  $S_n$  represents a Binomial distribution with parameters  $n$  and  $1/2$ . Since  $\Pr(S_n \geq \lfloor \nu \rfloor + 1) \leq \Pr(S_n \geq \lfloor \nu \rfloor)$ , we have

$$\begin{aligned} \Pr(S_n \geq \nu + 1) &= \Pr(S_n \geq \lfloor \nu \rfloor + 1) \leq B(n, \Gamma_i) \\ &\leq \Pr(S_n \geq \lfloor \nu \rfloor) = \Pr(S_n \geq \nu), \end{aligned}$$

since  $S_n$  is a discrete distribution. For  $\Gamma_i = \theta \sqrt{n}$ , where  $\theta$  is a constant, we have

$$\Pr\left(\frac{S_n - n/2}{\sqrt{n}/2} \geq \theta + \frac{2}{\sqrt{n}}\right) \leq B(n, \Gamma_i) \leq \Pr\left(\frac{S_n - n/2}{\sqrt{n}/2} \geq \theta\right).$$

By the central limit theorem, we obtain that

$$\begin{aligned} \lim_{n \rightarrow \infty} \Pr\left(\frac{S_n - n/2}{\sqrt{n}/2} \geq \theta + \frac{2}{\sqrt{n}}\right) &= \lim_{n \rightarrow \infty} \Pr\left(\frac{S_n - n/2}{\sqrt{n}/2} \geq \theta\right) \\ &= 1 - \Phi(\theta), \end{aligned}$$

where  $\Phi(\theta)$  is the cumulative distribution function of a standard normal. Thus, for  $\Gamma_i = \theta \sqrt{n}$ , we have

$$\lim_{n \rightarrow \infty} B(n, \Gamma_i) = 1 - \Phi(\theta). \quad \square$$

## Acknowledgments

The authors thank the reviewers of the paper for several insightful comments. The research of Dimitris Bertsimas was partially supported by the Singapore-MIT alliance. The research of Melvyn Sim is supported by a graduate scholarship from the National University of Singapore.

## References

- Ben-Tal, A., A. Nemirovski. 1998. Robust convex optimization. *Math. Oper. Res.* **23** 769–805.
- Ben-Tal, A., A. Nemirovski. 1999. Robust solutions to uncertain programs. *Oper. Res. Lett.* **25** 1–13.
- Ben-Tal, A., A. Nemirovski. 2000. Robust solutions of linear programming problems contaminated with uncertain data. *Math. Programming* **88** 411–424.
- El-Ghaoui, L., H. Lebret. 1997. Robust solutions to least-square problems to uncertain data matrices. *SIAM J. Matrix Anal. Appl.* **18** 1035–1064.
- El-Ghaoui, L., F. Oustry, H. Lebret. 1998. Robust solutions to uncertain semidefinite programs. *SIAM J. Optim.* **9** 33–52.
- Freund, R. M. 1985. Postoptimal analysis of a linear program under simultaneous changes in matrix coefficients. *Math. Programming Stud.* **24** 1–13.
- Robbins, H. 1955. A remark of Stirling's formula. *Amer. Math. Monthly* **62** 26–29.
- Soyster, A. L. 1973. Convex programming with set-inclusive constraints and applications to inexact linear programming. *Oper. Res.* **21** 1154–1157.