

Queueing theory fundamentals

Stochastic Delays

Cathy Wu

1.041/1.200 Transportation: Foundations and Methods

References

1. Readings: Chapter 2 of Larson and Odoni (2007), Urban Operations Research.
2. Walrand, Jean. Probability in Electrical Engineering and Computer Science: An Application-Driven Course. Quorum Books (2014).
3. Slides adapted from Carolina Osorio

Outline

1. Delays in transportation
2. Queueing theory fundamentals
3. Problem: timing of a pedestrian crossing light

Outline

1. **Delays in transportation**
2. Queueing theory fundamentals
3. Problem: timing of a pedestrian crossing light

Tl;dr: Delay is a big AND growing problem.

Delays in transportation

- Congestion/delays arises across all transportation modes
- Sources: Urban mobility report (TTI, 2009); Transportation Vision for 2030 (US DOT, 2008)



Delays in transportation

- Urban congestion:

- “Urban congestion lead in 2007 to an estimated additional 4.2 billion hours of travel and 2.8 billion gallons of fuel with a cost of \$87.2 billion across urban areas in the US, an increase of more than 50% over the previous decade”

- Highway traffic:

- “Highway vehicle miles traveled are projected to grow 60%, from 2,952 billion miles traveled in 2005 to 4,733 billion miles traveled in 2030.”

Delays in transportation

■ Air traffic:

- “The airline industry’s on-time performance in the first seven months of 2007 was the worst on record, and nationally almost 30% of all flights are now cancelled or substantially delayed.”
- “Aircraft travel is projected to nearly double, and current forecasts estimate over 1.5 billion air passengers annually by 2030. This will place unparalleled demand on the air system.”

■ Freight traffic:

- “The U.S. transportation system currently moves over 50 million tons of freight on the US transportation network. (...) By 2035, tons transported overall are expected to double to over 100 million, placing incomparable pressure on our domestic transportation network.”



Google maps hack Artist creates traffic jams in Google Maps with a wagon full of phones

This is the kind of post-modern art we can get behind.



Igor Bonifacic, @igorbonifacic
February 3, 2020

40

Comments

15

Shares

[Video](#)



Simon Weckert

Importance of modeling delay

- Queueing leads to delay
- Given the high cost of delay, there is a need for methods to:
 - identify sources/causes of delay (e.g. bottlenecks)
 - quantify occurrence of delay: how often does it occur?
 - duration of delay: how long does it typically take for a queue to dissipate?
 - ultimately: identify ways of reducing delay or mitigating its impact
- Queueing theory:
 - predicts various characteristics of waiting lines (or queues) such as average waiting time, average queue length
 - allows accounting for the uncertainty in arrival patterns, service patterns and routing of vehicles/customers to a service location

Transportation applications of queueing models

- Road transportation
 - Left-turn bay length
 - Number of approach lanes at a signal controlled intersection
 - Number of toll booths
 - Number of border crossing booths
- Air transportation:
 - Airport runways: impacts of capacity improvements on flight delays
 - Airport parking facilities: number of parking fee collection booths
 - Airport check-in
- Maritime transportation
 - Analysis of flow at container terminals (e.g. entrance/exit gates, within yards)
- Operations / Control problems: e.g. signal control (traffic lights), scheduling of flight departures, priority boarding schemes
- Pedestrian flow through circulation systems (e.g. on MIT campus, Infinite Corridor, inside Stata Center)

Relevant for other fields

- Telecommunications
- Software architectures performance prediction
- Manufacturing systems
- Prisoner flow through a network of prisons with varying security levels
- Hospital patient flow within networks of hospital units
- Biology: ribosome flow within protein synthesis networks

Background for queuing theory

- To be useful, the assumed distributional form, should be:
 1. Sufficiently **realistic** (reasonable predictions)
 2. Sufficiently **simple** (mathematically tractable)
- Key tools
 - The Poisson process
 - The exponential distribution, which is intimately tied to the Poisson process
- Both play a central role in transportation modeling.
- They can often be used to model (approximately) the **occurrence of random events**, such as:
 - the arrival of uncoordinated demands at a transportation facility,
 - the passage of cars by an observation point on a highway with free-flowing traffic.
- The exponential r.v. is also often used to approximate **service times** at transportation facilities.


Outline

1. Delays in transportation
2. **Queueing theory fundamentals**
 - a. Exponential distribution
 - b. Poisson process
3. Problem: timing of a pedestrian crossing light

Exponential distribution

- Also called the negative exponential
- $T \sim \text{Exp}(\lambda)$
- λ is a parameter (a constant)
- PDF: $f_T(t) = \begin{cases} \lambda e^{-\lambda t} & t \geq 0 \\ 0 & t < 0 \end{cases}$
- $E[T] = \frac{1}{\lambda}$
- $\text{Var}[T] = \frac{1}{\lambda^2}$
- CDF: The probability that T will be less than or equal to any particular constant value, t , is equal to:


$$F_t(t) = P(T \leq t) = \int_{x=0}^t \lambda e^{-\lambda x} dx = 1 - e^{-\lambda t}$$



Exponential distribution: numerical example

Gap acceptance problem

- Suppose that T :
 - a) represents the times between successive passages (headways) of vehicles on a road; and
 - b) is described by an exponential pdf.
- A driver requires 5 sec to insert into mainstream traffic that flows at 400 veh/hr
- What is the probability that the next gap is acceptable?




Exponential dbn

- What are the implications of assuming $T \sim \text{Exp}(\lambda)$?

Main properties of the exponential dbn:

- Prop. 1) $f_T(t)$, the pdf, is a **strictly decreasing** function of t ($t \geq 0$)
 - $P(0 \leq T \leq \Delta t) > P(t \leq T \leq t + \Delta t)$, $t > 0$, $\Delta t > 0$
 - $P\left(0 < T < \frac{E[T]}{2}\right) = 0.393$
 - $P\left(\frac{E[T]}{2} < T < \frac{3E[T]}{2}\right) = 0.38$
 - So, T is more likely to be $< \frac{E[T]}{2}$ (small), than near $E[T]$, even though 2nd interval is twice as wide as the 1st.
- Implications in practice?



Exponential dbn

- Prop. 2) Lack of memory (“memoryless”)

$$P(T > t + s | T \geq s) = P(T > t) \quad t > 0, \quad s > 0$$

- Interpretation: The probability dbn of the remaining time until the occurrence of the next event (e.g. next vehicle arrival, next service completion) is always the same, **regardless of how much time has already passed**.
 - For inter-arrival times: the time until the next arrival is uninfluenced by when the last arrival occurred
 - For service times: if they differ from customer to customer, then “memoryless” may be a desirable property
- The exponential dbn is the only continuous dbn with this property.
- Also, a small interval is independent of the time:

$$P(T \leq t + \varepsilon | T > t) = \lambda \varepsilon + o(\varepsilon), \quad \varepsilon \ll 1$$

- Implications in practice?

Example: Jitney rider

- Carla waits on the side of the road for a **jitney**, which will transport her to the next town. This jitney travels along a fixed route between the edge of two towns.
 - A jitney is a form of unlicensed taxi, often unscheduled. Jitney service (in various forms) is prevalent in developing countries.
 - Trivia: Technically, Uber is considered an illegal taxi service in jurisdictions with medallion systems that restrict the number of legal cabs in operation.
- 1. Suppose that the interarrival time of the jitney service is an **exponentially distributed random variable with a mean of 10 minutes**. Carla has **already waited 15.5 minutes**. What is the expected additional time (conditional mean) that she will have to wait?
- 2. Now assume the jitney interarrival time is **uniformly distributed between 2 to 18 minutes**. What is the expected additional time (conditional mean) that Carla will have to wait?

Example: Jitney rider

1. Suppose that the interarrival time of the jitney service is an exponentially distributed random variable with a mean of 10 minutes. Carla has already waited 15.5 minutes. What is the expected additional time (conditional mean) that she will have to wait?
2. Now assume the jitney interarrival time is uniformly distributed between 2 to 18 minutes. What is the expected additional time (conditional mean) that Carla will have to wait?

Exponential dbn

- Prop. 2) Lack of memory (“memoryless”)

$$P(T > t + s | T \geq s) = P(T > t) \quad t > 0, \quad s > 0$$

Exponential dbn

- A small interval is independent of the time:

$$P(T \leq t + \varepsilon | T > t) = \lambda \varepsilon + o(\varepsilon), \quad \varepsilon \ll 1$$

Exponential dbn

- Prop. 3) Minimum of Exp. R.V.s yields an Exp. R.V.

$$T_i \sim \text{Exp}(\lambda_i), \quad \forall i \in 1, 2, \dots, n \text{ independent}$$

$$U = \min_{i=1, \dots, n} T_i$$

$$U \sim \text{Exp}\left(\sum_{i=1}^n \lambda_i\right)$$

- Inter-arrivals of different types, then the **time until the next arrival** follows an exponential dbn.
- Example: n check-in counters currently attending customers, then the **time until the next service completion** follows an exponential dbn.

Exponential dbn

- Prop. 4) Relationship to the **Erlang distribution**.

- k independent r.v's: $T_i \sim \text{Exp}(\lambda)$, $\forall i \in 1, 2, \dots, k$

Then, $E_k = \sum_{i=1}^k T_i$ follows a k^{th} -order Erlang dbn.

$$f_{E_k}(x) = \frac{\lambda^k x^{k-1} e^{-\lambda x}}{(k-1)!}, \quad x \geq 0, \quad k = 1, 2, \dots$$

- $E[E_k] = \frac{k}{\lambda}$
- $\text{Var}(E_k) = \frac{k}{\lambda^2}$
- Example: **total** waiting time of passengers for a bus
- Note (sanity check): the exponential dbn coincides with $k = 1$.

Exponential dbn


- Prop. 5) Relationship to the Poisson process.
- Suppose we have a sequence of demand arrivals (“events”) such that
 - a) successive demand inter-arrival times are mutually independent and
 - b) the demand inter-arrival times are all described by the same exponential pdf.Then, the **number of arrivals** constitutes a “Poisson process”.
- Stated the other way around:
The inter-arrival times between events in a Poisson process that occur at the rate of λ per unit of time are: (i) mutually independent; and (ii) described by an exponential pdf with parameter λ .

Outline

1. Delays in transportation
2. **Queueing theory fundamentals**
 - a. Exponential distribution
 - b. **Poisson process**
3. Problem: timing of a pedestrian crossing light

The Poisson process

- Count the number of events $N(t)$ that occur during time interval $[0, t]$
- Definition (Poisson process):
If successive inter-event (inter-arrival) times are independent and identically distributed as $Exp(\lambda)$, then: $\{N(t)\}$ is a Poisson process with rate λ
- $P(N(t) = n) = \frac{(\lambda t)^n e^{-\lambda t}}{n!}$: probability that n events take place during the time interval
- t : time interval length
- λ : arrival rate (veh/unit time), also referred to as the “intensity” of the arrivals of Poisson events
- $E[N(t)] = \lambda t, \text{Var}(N(T)) = \lambda t$



Source: wikipedia

Poisson process $\rightarrow$ Exponential dbn

- T is the time between successive arrivals, then during a time interval of length $t \leq T$, no arrivals occurred, so for all such t :

$$P(t \leq T) = P(N(t) = 0) = \frac{(\lambda t)^0 e^{-\lambda t}}{0!} = e^{-\lambda t}$$

- Cumulative dbn function:

$$F(t; \lambda) = P(0 \leq T \leq t) = 1 - e^{-\lambda t}$$

- Probability density function:

$$f(t; \lambda) = \frac{dF(t; \lambda)}{dt} = \lambda e^{-\lambda t}$$

- Inter-arrival times follow an exponential dbn, with parameter λ .

Inter-arrival times that are i.i.d. $Exp(\lambda)$ is equivalent to arrivals according to a Poisson process $P(\lambda)$

Fundamental properties of the Poisson process

1. For a sufficiently small Δt :

$$P(N(t + \Delta t) - N(t) = 1) \simeq \lambda \Delta t$$

$$P(N(t + \Delta t) - N(t) = 0) \simeq 1 - \lambda \Delta t$$

$$P(N(t + \Delta t) - N(t) \geq 2) \simeq 0$$

where these probabilities represent, respectively:

- the probability that exactly one event will occur in the next Δt
 - the probability that no event will occur in the next Δt
 - the probability that two or more events will occur in the next Δt
2. The number of events that occur in disjoint time intervals are mutually independent r.v.'s
 3. The number of events that occur during any pre-specified interval of length Δt does not depend on the “starting time” of the time interval or on the number of events recorded prior to the time interval

More properties of the Poisson process

- Given the relationship between the Poisson process and the exponential dbn, we have the following properties:
 1. (“Memoryless”) If we begin observing a Poisson process with rate λ at $t = 0$, the pdf for the time, X , until the next arrival is given by $f_X(t) = \lambda e^{-\lambda t}$, $t \geq 0$ no matter how long before $t = 0$ the last arrival occurred.
 2. “The sum of K independent Poisson processes is a Poisson process with a rate equal to the sum of the K rates”.
 3. Given two independent Poisson processes, with rates λ_1 and λ_2 , respectively. Let X_1 and X_2 , respectively, be the time until the next arrival from each process. Then, the probability that an arrival from Process 1 will take place before an arrival from Process 2 is equal to:

$$P(X_1 < X_2) = \frac{\lambda_1}{\lambda_1 + \lambda_2}$$

Poisson events and Bernoulli trials

- Note that by Property 3 in the previous slide, the probability that the next arrival from either process will be a process 1 arrival is given by:

$$p = \frac{\lambda_1}{\lambda_1 + \lambda_2}$$

and the probability that it will be a process 2 arrival by:

$$q = 1 - p = \frac{\lambda_2}{\lambda_1 + \lambda_2}$$

- Thus, the type of arrival (from process 1 or from process 2) that will occur when an arrival takes place is determined by a **Bernoulli trial** (“coin flip”).
- The probability that, out of n arrivals, m will be from process 1 is given by:

$$P(m \text{ successes in } n \text{ trials}) = C_m^n p^m (1 - p)^{n-m}, \quad 0 \leq m \leq n$$

Outline

1. Delays in transportation
2. Queueing theory fundamentals
3. **Problem: timing of a pedestrian crossing light**

Example: Timing of a pedestrian crossing light*

- Pedestrians approach from the left (resp. right) side of the crossing in a Poisson manner with arrival rate λ_L (resp. λ_R) arrivals per minute. Pedestrians wait until light turns green, referred to as a “dump”. Assume:
 - all pedestrians cross instantaneously
 - the dump duration is zero
 - left and right arrival processes are independent.
- We consider three operating rules:
 - Dump every t_1 minutes
 - Dump whenever the total number of waiting pedestrians equals n_0
 - Dump when the first pedestrian to arrive, after the previous dump, has waited for t_2 minutes
- For each rule, determine:
 1. The expected number of pedestrians crossing left to right on any dump
 2. The probability that zero pedestrians cross left to right on any particular dump
 3. The pdf for the time between dumps




FIGURE 2.14 Pedestrian crossing problem.


Rule C: Dump whenever the *first* pedestrian to arrive after the previous dump has waited T_0 minutes.

Example: Timing of a pedestrian crossing light*

- Arrival rates λ_L, λ_R
- We consider three operating rules:
 - A. Dump every t_1 minutes
 - B. Dump whenever the total number of waiting pedestrians equals n_0
 - C. Dump when the first pedestrian to arrive, after the previous dump, has waited for t_2 minutes

- For each rule, determine:

- 1. The expected number of pedestrians crossing left to right on any dump
- 2. The probability that zero pedestrians cross left to right on any particular dump
- 3. The pdf for the time between dumps



Example: Timing of a pedestrian crossing light*

- Arrival rates λ_L, λ_R
- We consider three operating rules:
 - A. Dump every t_1 minutes
 - B. Dump whenever the total number of waiting pedestrians equals n_0
 - C. Dump when the first pedestrian to arrive, after the previous dump, has waited for t_2 minutes

- For each rule, determine:

- 1. The expected number of pedestrians crossing left to right on any dump
- 2. The probability that zero pedestrians cross left to right on any particular dump
- 3. The pdf for the time between dumps

