


3134393684

Suppose  $\alpha^3 + 2\alpha^2 + 1 = 0$  and  $K = 2 + \alpha^2$ .

$$\sigma(n) = \begin{cases} \alpha^{24-9n} & \text{if } n = 4, 6, 8; \\ \alpha^{9n-6} & \text{if } n \text{ even otherwise;} \\ \alpha^{9n-10} & \text{if } n \text{ odd.} \end{cases}$$

$$z^4 - (22432i + 17212)z^3 + (24822589i + 14048442)z^2 + (195926151359i - 197283732239)z + (103093386360336i - 218552646456468)$$





---

ANDREW ZOMBIEBUYERS

---

$$\begin{pmatrix} 2 & 2 & 0 \\ 1 & 2 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad K = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$$




---


FEW = 74971	FOLK = 2014909	CALMLY = 5861084641
DRIVE = 80479069	BLACK = 24351365	TAXIS = 12013337
UP = 267	MAJOR = 682594	ROADS = 27609131
ON = 3053	QUIET = 94459883	HAZY = 1856077
NIGHTS = 188607503		

---

Let  $x_{12}$  be an equilateral triangle of area  $x_1$  and set  $x_7$  as the radius of its incircle  $x_{25}$ . Suppose that the origin  $x_{21}$  is contained within  $x_{25}$ , and choose a point  $(x_{18}, x_{23}, x_{17})$  with prime distance  $x_{20}$  from  $x_{21}$ . Let  $x_{14}$  be the group of isometries fixing  $x_{25}$  and set  $x_9, x_{24}$  and  $x_8$  as the length, width and height of the smallest bounding box  $x_4$  containing all translates of  $(x_{18}, x_{23}, x_{17})$  under  $x_{14}$ .

Now fix an irreducible polynomial  $x_{26}$  defining a field  $x_{15}$  of characteristic  $x_{20}$ . Let  $x_{19}$  be a vector space over  $x_{15}$  of even dimension  $x_2$  and choose a matrix  $x_{16}$  in  $x_{10}x_5(x_{19})$ . Then obviously  $(x_{16})_{x_{11}, x_{13}} < x_8$ .  
 $x_{22}x_6x_3$

		On $\cdot$		On $-$		On $\cup$	
$\rightarrow$ 0	$\cdot$	22	R	$-$ 22	R	$\cup$ 0	R
1	$\cup$	11	L	$\cup$ 9	L	$\cup$ 2	L
2	$\cup$	2	R	$\cup$ 2	R	$\cup$ 2	R
3	$\cup$	6	R	$\cdot$ 6	L	$\cup$ 6	R
4	$\cup$	X		$-$ X		$\cup$ 2	L
5	$\cdot$	25	R	$\cdot$ 28	R	$\cup$ 2	R
6	$\cdot$	X		$\cdot$ X		$\cdot$ X	
7	$\cdot$	6	R	$\cdot$ 6	R	$\cdot$ 6	R
8	$\cdot$	29	R	$\cdot$ 29	R	$\cdot$ 29	R
9	$\cdot$	10	L	$-$ 10	L	$\cup$ 2	L
10	$\cdot$	25	L	$-$ 25	L	$\cup$ 2	L
11	$\cdot$	13	L	$-$ 3	L	$\cup$ 2	L
12	$\cdot$	11	L	$-$ 4	L	$\cup$ 2	L
13	$\cdot$	6	L	$-$ 6	L	$\cup$ 25	R
14	$\cdot$	15	L	$-$ 17	L	$\cup$ 2	R
15	$\cdot$	X		$-$ 19	L	$\cup$ 2	R
16	$\cdot$	X		$-$ 18	L	$\cup$ 2	R
17	$\cdot$	20	L	$-$ X		$\cup$ 2	R
18	$\cdot$	X		$-$ 21	L	$\cup$ 2	R
19	$\cdot$	30	R	$-$ X		$\cup$ 2	R
20	$\cdot$	8	R	$-$ X		$\cup$ 2	R
21	$\cdot$	26	R	$-$ X		$\cup$ 2	R
22	$\cdot$	23	R	$-$ 23	R	$\cup$ 5	L
23	$\cdot$	24	R	$-$ 24	R	$\cup$ 12	L
24	$-$	14	L	$\cdot$ 16	L	$\cup$ 1	L
25	$-$	X		$-$ X		$-$ X	
26	$-$	27	R	$-$ 27	R	$-$ 27	R
27	$-$	4	R	$-$ 4	R	$-$ 4	R
28	$-$	29	R	$-$ 29	R	$-$ 29	R
29	$-$	6	R	$-$ 6	R	$-$ 6	R
30	$-$	7	R	$-$ 7	R	$-$ 7	R



$\sigma(x_7) = x_9$	$x_{22} \subset x_{15} \cup x_{21}$	$x_{10}, x_{16}, x_{20}, x_{23} \in \text{Stabilizer}(\sigma)$
$\sigma(x_{17}) = x_{25}$	$X_{13} \subset X_{18}$	$x_{10}, x_{12}, x_{20} \in \text{Stabilizer}(\tau\sigma)$
$\tau(x_4) = x_{25}$	$X_{14} \subset X_{10} \cup X_{15}$	$x_5, x_6, x_{15}, x_{24}, x_{26}, X_1, X_5, X_{15}, X_{19}, X_{26} \in \text{Stabilizer}(\tau)$
$\tau(x_8) = x_{17}$	$X_{17} \subset X_3$	$x_5, x_{24}, X_{11}, X_{12}, X_{15}, X_{18}, X_{19}, X_{25} \in \text{Stabilizer}(\tau\rho)$
$\rho(X_2) = X_6$	$X_{23} \subset X_{10} \cup X_{18}$	$X_7, X_{20}, X_{23} \in \text{Stabilizer}(\rho)$

$$\frac{x + 4x^3 - 8x^4 + 9x^5 - 8x^6 + 4x^7}{1 - 2x + x^2 + x^4 - 2x^5 + x^6}$$

