

5.73

Quiz 26

November 6, 2002

$$\mathbf{H}^{\text{SO}} = \frac{\zeta}{\hbar} \mathbf{L} \cdot \mathbf{S} = \frac{1}{2} \frac{\zeta}{\hbar} [\mathbf{J}^2 - \mathbf{L}^2 - \mathbf{S}^2]$$

$$\mathbf{H}^{\text{Zeeman}} = -\gamma B_z (\mathbf{L}_z + 2\mathbf{S}_z)$$

Using ladders plus orthogonality, you should have obtained

$$|{}^2F_{7/2}, M_J = 5/2\rangle = \left(\frac{6}{7}\right)^{1/2} |{}^2F, M_L = 2, M_S = 1/2\rangle + \left(\frac{1}{6}\right)^{1/2} |{}^2F, M_L = 3, M_S = -1/2\rangle$$

and

$$|{}^2F_{5/2}, M_J = 5/2\rangle = -\left(\frac{1}{7}\right)^{1/2} |{}^2F, M_L = 2, M_S = 1/2\rangle + \left(\frac{6}{7}\right)^{1/2} |{}^2F, M_L = 3, M_S = -1/2\rangle.$$

A. Compute $E^{\text{SO}}({}^2F_{7/2}) = \langle {}^2F_{7/2}, M_J = 5/2 | \mathbf{H}^{\text{SO}} | {}^2F_{7/2}, M_J = 5/2 \rangle$.

B. Compute $E^{\text{SO}}({}^2F_{5/2}) = \langle {}^2F_{5/2}, M_J = 5/2 | \mathbf{H}^{\text{SO}} | {}^2F_{5/2}, M_J = 5/2 \rangle$.

C. Compute $E^Z({}^2F_{5/2}, 5/2) = \langle {}^2F_{5/2}, M_J = 5/2 | \mathbf{H}^{\text{Zeeman}} | {}^2F_{5/2}, M_J = 5/2 \rangle$.

D. Compute the off-diagonal matrix element:

$$\mathbf{H}_{5/2, 5/2; 7/2, 5/2}^Z = \langle {}^2F_{5/2}, M_J = 5/2 | \mathbf{H}^{\text{Zeeman}} | {}^2F_{7/2}, M_J = 5/2 \rangle.$$

E. Use second order perturbation theory to derive the Zeeman tuning rate for the nominal ${}^2F_{5/2}, M_J = 5/2$ state. The Zeeman tuning rate is $\frac{dE}{dB_Z}$.

$$E^{(0)}({}^2F_{5/2}, M_J = 5/2) = E^{\text{SO}}({}^2F_{5/2}) + E^Z({}^2F_{5/2}, 5/2)$$

$$E^{(2)}({}^2F_{5/2}, M_J = 5/2) = \frac{|\mathbf{H}_{5/2, 5/2; 7/2, 5/2}^Z|^2}{E^{\text{SO}}({}^2F_{5/2}) - E^{\text{SO}}({}^2F_{7/2})}.$$