

# Insulin-like signalling to the maternal germline controls progeny response to osmotic stress

Nicholas O. Burton<sup>1</sup>, Tokiko Furuta<sup>2</sup>, Amy K. Webster<sup>3</sup>, Rebecca E. W. Kaplan<sup>3</sup>, L. Ryan Baugh<sup>3</sup>, Swathi Arur<sup>2</sup> and H. Robert Horvitz<sup>1,4</sup>

**In 1893 August Weismann proposed that information about the environment could not pass from somatic cells to germ cells<sup>1</sup>, a hypothesis now known as the Weismann barrier. However, recent studies have indicated that parental exposure to environmental stress can modify progeny physiology<sup>2–7</sup> and that parental stress can contribute to progeny disorders<sup>8</sup>. The mechanisms regulating these phenomena are poorly understood. We report that the nematode *Caenorhabditis elegans* can protect itself from osmotic stress by entering a state of arrested development and can protect its progeny from osmotic stress by increasing the expression of the glycerol biosynthetic enzyme GPDH-2 in progeny. Both of these protective mechanisms are regulated by insulin-like signalling: insulin-like signalling to the intestine regulates developmental arrest, while insulin-like signalling to the maternal germline regulates glycerol metabolism in progeny. Thus, there is a heritable link between insulin-like signalling to the maternal germline and progeny metabolism and gene expression. We speculate that analogous modulation of insulin-like signalling to the germline is responsible for effects of the maternal environment on human diseases that involve insulin signalling, such as obesity and type-2 diabetes<sup>8</sup>.**

Maternal exposure to a wide variety of environmental stresses alters progeny growth, development and physiology of diverse organisms<sup>2–7</sup> and is thought to be a contributing factor to several human pathologies, including obesity and diabetes<sup>8</sup>. The mechanisms by which the maternal environment can modify progeny biology are poorly understood. Parental exposure of the nematode *Caenorhabditis elegans* to mild osmotic stress can protect progeny from the effects of strong osmotic stress<sup>9</sup>. This finding and similar observations of other organisms<sup>2</sup> suggest that besides the potentially deleterious effects of maternal environmental stress on progeny, maternal exposure to environmental stress might epigenetically precondition progeny and protect them from similar environmental insults in the future. How


maternal exposure to environmental stress can protect progeny from future environmental stress remains largely unknown.

To determine how parental exposure to mild osmotic stress (300 mM NaCl) protects progeny from the effects of strong osmotic stress (500 mM NaCl) we first examined the effects of 500 mM NaCl on *C. elegans*. Embryos placed at 500 mM NaCl completed embryonic development and hatched but arrested development immediately after hatching (Fig. 1a and Supplementary Fig. 1a). These arrested animals were unable to move, feed or respond to touch (Supplementary Fig. 1a). However, when arrested animals were returned to normal osmotic conditions (50 mM NaCl), they regained mobility and resumed development (Fig. 1b). We made similar observations when NaBr, KCl or sucrose was used to cause osmotic stress (Supplementary Fig. 1b–d). We conclude that young *C. elegans* larvae can enter a state of immobile arrested development in response to osmotic stress.

*C. elegans* also arrests development in response to other environmental stresses, such as starvation<sup>10,11</sup>. These arrests are caused by the loss of insulin-like signalling via the insulin receptor DAF-2 and the consequent activation of the FOXO transcription factor DAF-16 (refs 10–12). We tested whether the activation of DAF-16 is required for developmental arrest in response to osmotic stress. Animals lacking DAF-16 were less likely than the wild type to arrest development in response to osmotic stress (Fig. 1a), whereas animals with increased DAF-16 activation, such as *daf-2* mutants<sup>12</sup>, were more likely to arrest development in response to osmotic stress (Fig. 1a). Genes in other stress response pathways, such as those that regulate the responses of *C. elegans* to oxidative stress<sup>13</sup> or infection<sup>14</sup>, were not required for developmental arrest in response to osmotic stress (Supplementary Fig. 1e,f). In addition, DAF-2 activity in the intestine was required for development at 300 mM NaCl (Fig. 1c), and exposure to 500 mM NaCl caused the translocation of DAF-16 from the cytoplasm to the nucleus (Fig. 1d). Like the developmental arrest caused by starvation<sup>10</sup>, the arrest caused by osmotic stress required dense-core vesicle release (Supplementary Fig. 1g); furthermore, we found that it is

<sup>1</sup>Howard Hughes Medical Institute, Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA. <sup>2</sup>Department of Genetics, UT MD Anderson Cancer Center, Houston, Texas 77030, USA. <sup>3</sup>Department of Biology Duke University, Durham, North Carolina 27708, USA.

<sup>4</sup>Correspondence should be addressed to H.R.H. (e-mail: [Horvitz@mit.edu](mailto:Horvitz@mit.edu))



**Figure 1** Insulin-like signalling to the intestine regulates developmental arrest in response to osmotic stress. **(a)** Percentage of wild-type, *daf-2(e1370)*, *age-1(hx546)*, *pdk-1(sa709)*, *daf-16(mu86)* and *daf-2(e1370); daf-16(mu86)* animals developing past the L1 larval stage (L2+) after 48 h. Error bars, s.d.  $n=3$  experiments of >100 animals;  $P<0.01$  for all genotypes (two-tailed  $t$ -test). **(b)** Percentage of wild-type, *daf-2(e1370)*, *daf-16(mu86)* and *daf-2(e1370); daf-16(mu86)* animals that resume development after 24 h of exposure to osmotic stress. Error bars, s.d.  $n=3$ ; see Supplementary Table 6;  $P<0.01$  for all genotypes (two-tailed  $t$ -test). **(c)** Percentage of wild-type and *daf-2(e1370)* animals developing past the


L1 larval stage at 300 mM NaCl after 48 h. Neuron-specific expression was driven by *Prgef-1*; intestine-specific expression was driven by *Pges-1*; muscle-specific expression was driven by *Pmyo-3*. Error bars, s.d.  $n=3$  experiments of >100 animals. **(d)** Confocal images of DAF-16::GFP after 6 h of exposure to 50 mM and 500 mM NaCl. Percentage values indicate the percentage of embryos resembling the representative image. See Supplementary Table 6. 3 experimental replicates of >100 animals. The quantified results are presented as mean  $\pm$  s.d. using two-tailed  $t$ -test; \*\*\*\* $P<0.0001$  were considered significant. See Statistics Source Data in Supplementary Table 6.

dense-core vesicle release from sensory neurons that was required for developmental arrest in response to osmotic stress (Supplementary Fig. 1g), and *daf-16* mutants, which are resistant to developmental arrest, showed an increased susceptibility to osmotic stress (Fig. 1b). These results indicate that like starvation, osmotic stress causes developmental arrest by inhibiting insulin-like signalling and that arrested development correlates with enhanced survival.

In contrast to starvation-induced developmental arrest, we found that developmental arrest in response to osmotic stress is regulated by a different insulin-like peptide, INS-3 (Supplementary Fig. 1h,i and Supplementary Table 1), than arrest in response to starvation (INS-4 and DAF-28; ref. 15) and that animals that arrest development in response to osmotic stress are immobile and unable to respond to touch, unlike arrest in response to starvation in which animals remain mobile<sup>10</sup>. In addition, we found that a majority (78%) of genes the expression of which reproducibly changed in response to osmotic stress were not affected by starvation (Supplementary Fig. 2 and Supplementary Table 2). These results suggest that these two arrest phenotypes are controlled by partially overlapping but distinct pathways.

Frazier and Roth (2009) found that parental exposure of *C. elegans* to mild osmotic stress protects progeny from the effects of strong osmotic stress and that this protection required DAF-2 activation<sup>9</sup>. These authors described an apparently anomalous result: *daf-2; daf-16* double mutants were significantly better at adaptation to osmotic

stress than both wild-type animals and either *daf-2* or *daf-16* single mutants<sup>9</sup>. We hypothesized that this anomaly might be at least in part caused by differing effects of maternal insulin-like signalling and progeny insulin-like signalling with respect to progeny response to osmotic stress. Specifically, we suspected that DAF-2 activation in embryos was required for adaptation to osmotic stress, consistent with both the earlier observations<sup>9</sup> and our findings (Fig. 1a). We also suspected that it was an inhibition rather than an activation of parental insulin-like signalling that resulted in the protection of progeny from osmotic stress, just as the inhibition of larval insulin-like signalling induced by osmotic stress protects larvae from osmotic stress by causing developmental arrest. To test this hypothesis, we crossed wild-type animals with *daf-2* and *daf-2; daf-16* double-mutant animals in normal osmotic conditions (50 mM NaCl) and assayed the response of their progeny to 500 mM NaCl. Approximately 60% of the progeny from the cross of wild-type males with *daf-2* mutant hermaphrodites hatched and developed at 500 mM NaCl (Fig. 2a). By contrast, the reciprocal cross of *daf-2* mutant males with wild-type hermaphrodites and the cross of wild-type males with *daf-2; daf-16* double-mutant hermaphrodites did not produce any progeny that hatched and developed at 500 mM NaCl (Fig. 2a). These results demonstrate that like parental exposure to mild osmotic stress, reduced maternal insulin-like signalling can protect progeny from the effects of strong osmotic stress. Importantly, these observations also




**Figure 2** Insulin-like signalling to the maternal germline regulates progeny response to osmotic stress. **(a)** Percentage of wild-type (WT), *daf-2(e1370)* and *daf-2(e1370); daf-16(mu86)* cross progeny failing to arrest development after 48 h at 500 mM NaCl. Males contained (*Pegl-1::4xNLS::GFP*); *him-5(e1490)*; *nls349 (Pceh-28::4xNLS::mCherry)* for the identification of cross progeny. The *pie-1* promoter was used to drive germline-specific expression of DAF-2 and the *mex-5* promoter was used to drive germline-specific expression of DAF-16. Error bars, s.d.  $n=7, 3, 6, 3, 3, 3$  and 3; see Supplementary Table 6. **(b)** Percentage of wild-type, *gpdh-1(ok1558)* and *gpdh-2(ok1733)* animals failing to arrest development at 500 mM NaCl

after 48 h. Error bars, s.d.  $n=3$  experiments of >100 animals. **(c)** Average fold change of 2 replicates of the 25 most upregulated genes in embryos in response to osmotic stress after 6 h. FPKM, fragments per kilobase of transcript per million mapped reads. **(d)** Percentage of wild-type, *daf-2(e1370)* and *gpdh-2(ok1733)* cross progeny failing to arrest development after 48 h at 500 mM NaCl. Males contained *otIs39 (Punc-47::GFP)*; *him-5(e1490)* for the identification of cross progeny. Error bars, s.d.  $n=3$  experiments of >20 animals. The quantified results are presented as mean  $\pm$  s.d. using ANOVA. \* $P < 0.05$ , \*\*\* $P < 0.001$ , \*\*\*\* $P < 0.0001$  were considered significant. NS, not significant. See Statistics Source Data in Supplementary Table 6.

indicate that there is a previously undescribed link between maternal insulin-like signalling and progeny physiology.

Parental exposure to osmotic stress has been hypothesized to protect progeny from the effects of osmotic stress by increasing the deposition of glycerol from mothers into embryos, since embryos from parents exposed to 300 mM NaCl have more glycerol than embryos from animals grown at normal osmotic conditions (50 mM NaCl)<sup>9</sup> and glycerol is known to be protective against various environmental stresses<sup>16,17</sup>. We confirmed that exposure of parents to 300 mM NaCl resulted in progeny that are resistant to the effects of 500 mM NaCl (Fig. 2b,c and Supplementary Table 3), and we discovered that the glycerol biosynthetic enzyme GPDH-2 is required for parental exposure to 300 mM NaCl to protect progeny from 500 mM NaCl (Fig. 2b) but does not affect the response to osmotic stress of animals with parents grown at 50 mM NaCl (Supplementary Fig. 3a). These observations are consistent with the hypothesis that an increased level of glycerol is required for

adaptation to osmotic stress. However, it remained unclear how these observations relate to our finding that reduced maternal insulin-like signalling can protect progeny from strong osmotic stress (Fig. 2a), since previous studies indicated that embryos from *daf-2* mutant hermaphrodites contain the same amount of glycerol as embryos from wild-type animals<sup>9</sup> and hence that *daf-2* mutant mothers do not deposit more glycerol into embryos. We hypothesized that reduced maternal insulin-like signalling protects progeny from the effects of osmotic stress not by increasing deposition of glycerol from mothers into embryos but rather by increasing glycerol production in embryos via GPDH-2. To test this hypothesis, we crossed *gpdh-2* mutant males with *daf-2; gpdh-2* double-mutant hermaphrodites. GPDH-2 was required for reduced maternal insulin-like signalling to protect progeny from developmental arrest (Fig. 2d). To test whether GPDH-2 functions maternally, we crossed wild-type males with *daf-2; gpdh-2* double-mutant hermaphrodites. GPDH-2 was not required in mothers to protect progeny from developmental arrest in




**Figure 3** Insulin-like signalling to the maternal germline modifies progeny response to osmotic stress by regulating the RAS–ERK-like pathway. **(a)** Percentage of wild-type and *lin-45*(*n2018*), *mek-2*(*ku114*), *mpk-1*(*n5639*) and *lin-45*(*n2018*); *gpdh-2*(*ok1733*) animals failing to arrest development at 500 mM NaCl after 48 h. Error bars, s.d.  $n=6$ , 6, 3, 3, and 3 experiments of >100 animals. See Supplementary Table 6. **(b)** Percentage of wild-type and *lin-45*(*n2018*) cross progeny failing to arrest development at 500 mM NaCl after 48 h. Males contained *otIs39* (*Punc-47::GFP*); *him-5*(*e1490*) for the identification of cross progeny. Error bars, s.d.  $n=3$  experiments of >20 animals. **(c)** Representative germlines dissected from wild-type animals exposed to either 50 mM NaCl or 300 mM NaCl and stained for DNA

(DAPI, white) and diphosphorylated MPK-1 (dpMPK-1) (red). Each condition was replicated 16 times. Asterisks, mitotic end of germline. **(d)** Relative expression of *gpdh-2* mRNA in wild-type and *lin-45*(*n2018*) embryos at 50 mM NaCl measured by qRT-PCR and normalized to the expression of the histone *his-24*. Error bars, s.d.  $n=3$  experiments from pellets of >1,000 embryos. **(e)** Glycerol-to-glucose ratio in wild-type and *lin-45*(*n2018*) mutant embryos at 50 mM NaCl. Error bars, s.d.  $n=3$  experiments from pellets of >1,000 embryos. The quantified results are presented as mean  $\pm$  s.d. using ANOVA **(a, b)** and two-tailed *t*-test **(d, e)**. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  were considered significant. NS, not significant. See Statistics Source Data in Supplementary Table 6.

response to osmotic stress (Fig. 2d). We conclude that the inhibition of maternal insulin-like signalling does not result in the increased deposition of glycerol into embryos but rather results in increased glycerol production in embryos. Importantly, these results reveal that there is a heritable link between maternal insulin-like signalling and progeny metabolism.

We hypothesized that the heritable effects of maternal insulin-like signalling on progeny might be mediated by insulin-like signalling to the germline. To test this hypothesis, we expressed rescuing copies of the wild-type *daf-2* or *daf-16* genes specifically in the germline. Germline-specific expression of either *daf-2* or *daf-16* was sufficient to rescue the effects of deficient maternal insulin-like signalling on progeny response to osmotic stress (Fig. 2a). In addition, overexpression of *daf-2* in the germline blocked the protective effects of parental exposure to 300 mM NaCl on progeny response to 500 mM NaCl (Supplementary Fig. 3b). These data suggest that maternal exposure to 300 mM NaCl inhibits insulin-like signalling to the germline and that this loss of insulin-like signalling to the germline protects progeny from the effects of osmotic stress.

Insulin-like signalling to the *C. elegans* germline both inhibits DAF-16<sup>18</sup> and activates the RAS–ERK pathway<sup>19</sup>, which includes the Raf protein LIN-45, the Mek protein MEK-2 and the Erk protein MPK-1<sup>20</sup>. We found that partial loss-of-function mutants in *lin-45*, *mek-2* or *mpk-1* (null mutants are lethal) did not arrest development at 500 mM NaCl (Fig. 3a and Supplementary Fig. 3c). In addition, treatment of wild-type animals with the MEK inhibitor U0126 (ref. 21) or RNAi knockdown of *mek-2* prevented developmental arrest in response to 500 mM NaCl (Supplementary Fig. 3d–f); RAS–ERK signalling functioned maternally to regulate progeny response to osmotic stress (Fig. 3b); GPDH-2 was required for reduced RAS–ERK signalling to protect animals from developmental arrest (Fig. 3a); and maternal exposure to osmotic stress inhibited MPK-1 activation in the germline (Fig. 3c and Supplementary Table 4), possibly by inhibiting the release of insulin-like peptides from sensory neurons (Supplementary Fig. 4a). We conclude that reduced insulin-like signalling from the soma (probably from sensory neurons) to the maternal germline protects progeny from the effects of osmotic stress by both activating DAF-16 and inactivating MPK-1.



**Figure 4** RAS–ERK signalling regulates *C. elegans* response to bacterial infection and starvation. **(a)** Percentage of animals expressing *lin-4::YFP* after 24 h of exposure to either *Escherichia coli* OP50 or *Pseudomonas aeruginosa* PA14. Error bars, s.d.  $n=3$  experiments of >100 animals. **(b)** Percentage of animals failing to arrest development after 24 h at 350 mM NaCl. Error bars, s.d.  $n=3$  experiments of >100 animals. **(c)** Percentage of wild-type, *lin-5*(*n2018*) and *daf-16*(*mu86*) mutants with a divided M-cell after 7 days without food in S-basal at 20 °C. Error bars, s.e.m.

$n=4$  experiments of >100 animals. **(d)** Model for how maternal exposure to osmotic stress inhibits DAF-2 activity in the germline and affects progeny response to osmotic stress. See text for details. Red, embryo; green, germline; purple, intestine. The quantified results are presented as mean  $\pm$  s.d. **(a,b)** and s.e.m. **(c)** using ANOVA **(a)** and two-tailed *t*-test **(b,c)**. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ , \*\*\*\* $P < 0.0001$  were considered significant. NS, not significant. See Statistics Source Data in Supplementary Table 6.

To further examine how reduced maternal insulin-like signalling via the RAS–ERK pathway modifies progeny physiology we performed RNA-seq of wild-type and *lin-5* mutant embryos. We identified a total of 616 genes upregulated more than twofold and 1,310 genes downregulated more than twofold in *lin-5* mutant embryos (Supplementary Table 5). Among the 616 upregulated genes was *gpdh-2*, and we confirmed that *gpdh-2* messenger RNA expression is upregulated approximately threefold in *lin-5* mutant embryos using quantitative PCR with reverse transcription (qRT-PCR) (Fig. 3d). To test whether increased GPDH-2 expression results in increased glycerol production, we compared the glycerol-to-glucose ratios in wild-type and *lin-5* mutant embryos by mass spectrometry. We observed an 82% increase in the glycerol-to-glucose ratio in *lin-5* mutant embryos compared with that in wild-type embryos (Fig. 3e). These data are consistent with our hypothesis that reduced insulin-like signalling to the maternal germline protects progeny from the effects of osmotic stress by increasing the expression in embryos of the rate-limiting glycerol biosynthetic enzyme GPDH-2.

We note that previous studies found that adaptation to osmotic stress resulted in up to a 1,000% increase in glycerol in *C. elegans*<sup>9</sup>,

significantly higher than the increase we observed in *lin-5* mutants. Our genetic data demonstrate that GPDH-2 is required for reduced RAS–ERK signalling in the maternal germline to protect progeny from developmental arrest (Fig. 3a). However, given the modest increase in glycerol levels in *lin-5* mutants it remains possible, and perhaps likely, that one or more of the additional 615 genes upregulated in *lin-5* mutants also contribute to the resistance of these animals to developmental arrest in response to osmotic stress.

We tested whether parental exposure to other environmental stresses, such as bacterial infection or starvation, could similarly modify progeny response to environmental stress via RAS–ERK signalling. Exposure of *C. elegans* to the opportunistic pathogen *Pseudomonas aeruginosa* PA14 slowed early larval development (Fig. 4a), parental exposure to PA14 enhanced this slowing of larval development (Fig. 4a), and this heritable slowing of larval development in response to bacterial infection required RAS–ERK signalling (Fig. 4a). However, maternal exposure to PA14 did not protect progeny from developmental arrest at 500 mM NaCl (Supplementary Fig. 4b) but rather resulted in progeny that were more sensitive to arrest in response to osmotic stress (Fig. 4b). In addition, we found that RAS–ERK signalling was

required for L1 arrest in response to starvation (Fig. 4c). Collectively, these results suggest that maternal exposure to environmental stress modifies progeny physiology via RAS–ERK signalling, but that the effects of these stresses on progeny are different for different environmental stresses.

In conclusion, we propose a model in which the inhibition of insulin-like signalling to both the intestine and the germline can enhance *C. elegans* survival during osmotic stress but in which the effects of inhibition of insulin-like signalling to these two tissues are distinct. Specifically, the loss of insulin-like signalling to the intestine enhances resistance to osmotic stress by promoting developmental arrest, whereas the loss of insulin-like signalling to the maternal germline enhances progeny resistance to osmotic stress by increasing glycerol synthesis in embryos (Fig. 4d). In this model, information about the maternal environment is inherited via germ cells to enhance progeny resistance to future environmental stress.

The salt concentrations at which *C. elegans* arrests development in response to osmotic stress are approximately those of seawater, 480 mM Na<sup>+</sup> and 559 mM Cl<sup>-</sup> (ref. 22). We speculate that both the state of immobile arrested development and the ability of parents to confer progeny resistance to osmotic stress evolved to enhance organismal survival in response to osmotic stress caused by seawater. The insulin signalling pathway is broadly conserved among metazoa, and we postulate that insulin signalling to the germline plays a role in several human developmental and metabolic abnormalities known to result from abnormal insulin signalling, such as intrauterine growth restriction, obesity and type-2 diabetes, all of which have been linked to maternal environmental stress<sup>8,23</sup>. Consistent with this hypothesis, a recent report found that feeding parental mice a high-fat diet causes epigenetic changes to oocytes that result in progeny that are more susceptible to both obesity and diabetes<sup>7</sup>. These observations of a maternal high-fat diet modifying progeny physiology via oocytes in mice are similar to our observations of maternal exposure to osmotic stress modifying progeny physiology via oocytes in *C. elegans*. We propose that modified insulin-like signalling to the mouse germline might be the mechanism underlying these epigenetic changes in mouse oocytes. □

## METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the [online version of this paper](#).

*Note: Supplementary Information is available in the online version of the paper*

## ACKNOWLEDGEMENTS

We thank E. J. Hubbard, K. Ashrafi, S. Mitani and the *Caenorhabditis* Genetic Center, which is funded by the NIH National Center for Research Resources (NCRR), for providing strains; N. An for strain management; and K. Burkhart, S. Luo, A. Doi, N. Paquin and A. Corrionero for helpful discussions. H.R.H. and

N.O.B. were supported by NIH grant GM024663 and NSF grant 1122374. T.F. and S.A. were supported by NIH grant GM98200 and ACS grant RSG014-044-DDC. L.R.B., A.K.W. and R.E.W.K. were supported by NIH grant GM117408. H.R.H. is an investigator of the Howard Hughes Medical Institute.

## AUTHOR CONTRIBUTIONS

N.O.B., T.F., A.K.W., R.E.W.K., S.A., L.R.B. and H.R.H. designed the experiments and analysed the data. N.O.B., T.F., A.K.W., R.E.W.K. and S.A. performed the experiments. N.O.B. and H.R.H. wrote the manuscript.

## COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Published online at <http://dx.doi.org/10.1038/ncb3470>

Reprints and permissions information is available online at [www.nature.com/reprints](http://www.nature.com/reprints)

- Weismann, A. *The Germ-plasm: A Theory of Heredity* (Charles Scribner's Sons, 1893).
- Dantzer, B. *et al.* Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal. *Science* **340**, 1215–1217 (2013).
- Radford, E. J. *et al.* *In utero* undernourishment perturbs the adult sperm methylome and intergenerational metabolism. *Science* **345**, <https://doi.org/10.1126/science.1255903> (2014).
- Dias, B. G. & Ressler, K. J. Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nat. Neurosci.* **17**, 89–96 (2014).
- Öst, A. *et al.* Paternal diet defines offspring chromatin state and intergenerational obesity. *Cell* **159**, 1352–1364 (2014).
- Carone, B. R. *et al.* Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell* **143**, 1084–1096 (2010).
- Huypens, P. *et al.* Epigenetic germline inheritance of diet-induced obesity and insulin resistance. *Nat. Genet.* **48**, 497–499 (2016).
- Gallo, L., Tran, M., Master, J., Moritz, K. & Wlodek, M. Maternal adaptations and inheritance in the transgenerational programming of adult disease. *Cell Tissue Res.* **349**, 863–880 (2012).
- Frazier III, H. N. & Roth, M. B. Adaptive sugar provisioning controls survival of *C. elegans* embryos in adverse environments. *Curr. Biol.* **19**, 859–863 (2009).
- Baugh, L. R. & Sternberg, P. W. DAF-16/FOXO regulates transcription of *cki-1/Cip/Kip* and repression of *lin-4* during *C. elegans* L1 arrest. *Curr. Biol.* **16**, 780–785 (2006).
- Hu, P. *J. Dauer* 1–19 (WormBook, 2007); <https://doi.org/10.1895/wormbook.1.144.1>
- Murphy, C. & Hu, P. J. *Insulin/insulin-like Growth Factor Signaling in C. elegans* (WormBook, 2013); <https://doi.org/10.1895/wormbook.1.164.1>
- An, J. H. & Blackwell, T. K. SKN-1 links *C. elegans* mesodermal specification to a conserved oxidative stress response. *Genes Dev.* **17**, 1882–1893 (2003).
- Kim, D. H. *et al.* A conserved p38 MAP kinase pathway in *Caenorhabditis elegans* innate immunity. *Science* **297**, 623–626 (2002).
- Chen, Y. & Baugh, L. R. *ins-4* and *daf-28* function redundantly to regulate *C. elegans* L1 arrest. *Dev. Biol.* **394**, 314–326 (2014).
- Lamitina, S. T., Morrison, R., Moeckel, G. W. & Strange, K. Adaptation of the nematode *Caenorhabditis elegans* to extreme osmotic stress. *Am. J. Physiol.* **286**, C785–C791 (2004).
- Lamitina, T., Huang, C. G. & Strange, K. Genome-wide RNAi screening identifies protein damage as a regulator of osmoprotective gene expression. *Proc. Natl Acad. Sci. USA* **103**, 12173–12178 (2006).
- Michaelson, D., Korta, D. Z., Capua, Y. & Hubbard, E. J. A. Insulin signaling promotes germline proliferation in *C. elegans*. *Development* **137**, 671–680 (2010).
- Lopez III, A. L. *et al.* DAF-2 and ERK couple nutrient availability to meiotic progression during *Caenorhabditis elegans* oogenesis. *Dev. Cell* **27**, 227–240 (2013).
- Sundaram, M. V. *RTK/Ras/MAPK Signaling* (WormBook, 2006); <https://dx.doi.org/10.1895/wormbook.1.80.1>
- Favata, M. F. *et al.* Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J. Biol. Chem.* **273**, 18623–18632 (1998).
- Pilson, E. Q. M. *An Introduction to the Chemistry of the Sea* 2nd edn, 67 (Cambridge Univ. Press, 2013).
- Prada, J. A. & Tsang, R. C. Biological mechanisms of environmentally induced causes of IUGR. *Eur. J. Clin. Nutr.* **52**, S21–7–28 (1998).

## METHODS

**Strains.** All *C. elegans* strains were cultured as described previously<sup>24</sup> and maintained at 20 °C unless noted otherwise. The Bristol strain N2 was the wild-type strain. Mutations used are:

LG1: *daf-16(mu86)*, *mek-2(ku114)*, *ins-18(ok1672)*, *ins-18(ok2478)*, *ins-26(tm1983)*, *ins-28(ok2722)*, *ins-29(tm1922)*, *ins-30(ok2343)*, *ins-33(tm2988)*, *ins-36(tm6125)*, *rrf-1(pk1417)*; *gpdh-1(ok1558)*

LGII: *ins-2(tm4467)*, *ins-3(ok2488)*, *ins-3(tm3608)*, *ins-4(ok3534)*, *ins-5(tm2560)*, *ins-6(tm2416)*, *ins-11(tm1053)*, *ins-12(tm2918)*, *ins-13(tm4856)*, *ins-14(tm4886)*, *ins-15(ok3444)*, *ins-19(tm5155)*, *ins-20(tm5634)*, *ins-31(ok3543)*, *ins-32(tm6109)*, *ins-37(tm6268)*, *age-1(hx546)*, *knuSi379 [Pmex-5::daf-16cDNA::GFP::nos-2 3' UTR,unc-119(+)]*

LGIII: *daf-2(e1370)*, *mpk-1(n5639)*, *ins-17(tm790)*, *ins-21(tm5180)*, *ins-22(tm4639)*; *gpdh-2(ok1733)*

LGIV: *lin-45(n2018)*, *ins-1(nj32)*, *ins-7(tm1907)*, *ins-8(tm4144)*, *ins-34(tm3095)*; *unc-31(ft1)*; *zIs356 [daf-16::GFP]*; *pmk-1(km25)*; *skn-1(zu67)*

LGv: *daf-28(tm2308)*, *ins-10(tm3498)*, *ins-27(ok2474)*, *ins-35(ok3297)*, *him-5(e1490)*

LGx: *ins-9(tm3618)*, *pdk-1(sa709)* *nIs349[Pceh-28::4xNLS::mCherry; lin-15(+)]*; unknown linkage: *vizIs23 [pie-1p::GFP::daf-2(WT)::pie-1 3'UTR + unc-119(+)]*, *vizIs22 [pie-1p::GFP::daf-2(WT)::pie-1 3'UTR + unc-119(+)]*, *nIs343 [Pegl-1::4xNLS::GFP; lin15(+)]*; *oIs39[unc-47p::GFP]*;

Extrachromosomal arrays: *naEx187[pGC467 (ins-3(+))*, *pRF4)rol-6(su1006)]*, *hpEx2906 [Prgef-1::daf-2; Pmyo-2::mCherry]*, *hpEx3369 [Pges-1::daf-2; Pmyo-2::mCherry]*, *hpEx2905 [Pmyo-3::daf-2; Pmyo-2::mCherry]* *fx325 [Posm-6::unc-31a; Pmyo-3::mCherry; unc-119(+)]*

**Assay for developmental arrest.** Approximately 200 developing eggs from mothers grown at 50 mM NaCl (unless otherwise noted) were collected and placed on standard NGM plates containing varying concentrations of NaCl at 25 °C for 48 h. After 48 h, animals that remained immobile and were not feeding were scored as arrested. Mobile animals that were feeding were scored as developing. Percentage of L2+ is defined by the percentage of animals mobile and feeding and that have developed past the L1 larval stage. Percentage failing to arrest is defined by the percentage of animals mobile and feeding (unlike animals normally arrested in response to osmotic stress) but includes L1-stage larvae.

**Assay for survival after arrest.** Approximately 100 developing eggs from mothers grown at 50 mM NaCl were collected and placed on standard NGM plates containing varying concentrations of NaCl at 20 °C for 24 h. After 24 h arrested animals were picked onto plates containing 50 mM NaCl and allowed to recover for 24 h. After 24 h the fraction that regained mobility and resumed development were scored as surviving and the fraction that failed to resume development and did not respond to touch were assumed to be dead.

**DAF-16::GFP localization.** Confocal microscopy was performed using a Zeiss LSM 800 instrument. The resulting images were prepared using ImageJ software (National Institutes of Health). Image acquisition settings were calibrated to minimize the number of saturated pixels and were kept constant throughout the experiment.

**mRNA expression analysis by RNA-seq and qRT-PCR.** L4-stage wild-type and *lin-45* animals were placed on standard NGM plates containing either 50 mM or 300 mM NaCl for 24 h. Developing eggs from these animals were collected and placed at either 50 mM or 500 mM NaCl for 6 h. After 6 h embryos were collected in M9, and RNA was extracted using TissueRuptor and the RNeasy Mini kit (QIAGEN). For RNA-seq, RNA integrity and concentration were checked on a Fragment Analyzer (Advanced Analytical). The mRNA was purified by polyA-tail enrichment, fragmented, and reverse transcribed into cDNA (Illumina TruSeq). cDNA samples were then end-repaired and adaptor-ligated using the SPRI-works Fragment Library System I (Beckman Coulter Genomics) and indexed during amplification. Libraries were quantified using the Fragment Analyzer (Advanced Analytical) and qPCR before being loaded for paired-end sequencing using the Illumina NextSeq. For qRT-PCR reverse transcription was performed using SuperScript III (Invitrogen), and quantitative PCR was performed using Applied Biosystems Real-Time PCR Instruments. All results are normalized to the mRNA levels of the histone gene *his-24*.

Primers

*gpdh-2*

forward: ttgtatccaaccgtccgtat reverse: cgaattgatgtggaacaacg

*his-24*

forward: atgatcaaggagccatcaa reverse: tgacattgatctggatgaca

**Assay for wild-type adaptation to osmotic stress.** Wild-type embryos were placed on standard NGM plates containing 300 mM NaCl at 20 °C for 60 h and allowed

to develop to adulthood. Embryos from these adult animals were extracted using a razor blade into M9 solution and pipetted onto plates containing 500 mM NaCl. These animals were allowed to develop for 48 h at 20 °C. Percentage failing to arrest is defined by the fraction of animals that were mobile and feeding.

**Cross progeny analysis.** Approximately 20 L4 hermaphrodites were crossed with wild-type or *daf-2(e1370)* males for 24 h at 25 °C. After 24 h, embryos were dissected from hermaphrodites using a razor blade and then placed on plates containing 500 mM NaCl at 20 °C for 24 h. Percentage of L2+ is defined by the percentage of animals that developed past the L1 larval stage. All males contained *nIs343 (Pegl-1::4xNLS::mCherry)*; *nIs349 (Pceh-28::4xNLS::mCherry)* for the identification of cross progeny.

**Photographs.** Photographs of animals and embryos on standard NGM plates containing either 300 mM or 500 mM NaCl were obtained using an AxioCam MRm camera (Zeiss).

**dpMPK-1 imaging.** Wild-type animals at the L4 stage were placed on either 50 mM or 300 mM NaCl NGM plates seeded with OP50 at 20 °C for 24 h, and germlines were extruded. Dissections were performed as described previously<sup>25,26</sup>. Briefly, dissections were performed within 5 min of adding levamisole to achieve optimal diphosphorylated MPK-1 (dpMPK-1) staining. The dissected germlines were then fixed in 3% paraformaldehyde for 10 min, followed by a post-fix in 100% methanol at -20 °C. The fixed germ lines were then processed for immunofluorescence staining as described previously<sup>27-29</sup>. Anti-MAP kinase was used at a dilution of 1:200 (Clone MAPK-YT, Sigma). Secondary antibodies were donkey anti-mouse Alexa Fluor 594 and used at a dilution of 1:400. Each gonad was photographed as a montage, with each image taken as a 0.15 μ section and captured with overlapping cell boundaries at ×63 magnification. Images were taken using a Zeiss Axio Imager upright microscope with AxioVs40 V4.8.2.0 microimaging software and an Axio MRm camera (Zeiss). Montages were then assembled using Adobe Photoshop CS5.1, and white levels were uniformly adjusted using Affinity Illustrator to reduce background.

**RNA-seq prep and data analysis.** For Supplementary Tables 1 and 3 RNA integrity and concentration were checked on a Fragment Analyzer (Advanced Analytical). The mRNA was purified by polyA-tail enrichment, fragmented, and reverse transcribed into cDNA (Illumina TruSeq). cDNA samples were then end-repaired and adaptor-ligated using the SPRI-works Fragment Library System I (Beckman Coulter Genomics) and indexed during amplification. Libraries were quantified using the Fragment Analyzer (Advanced Analytical) and qPCR before being loaded for paired-end sequencing using the Illumina NextSeq. For QC purposes, BEDTools<sup>30</sup> (version: 2.25.0) was used to count the reads falling into genes, coding regions, intronic regions, 5' or 3' UTRs, flanking 3-kb genic regions and intergenic regions. Other basic statistics, including mapping rate, ratio of sense versus anti-sense reads and rRNA percentages were also collected for each sample. The reads were first cleaned up by removing reads aligned to rRNAs (BWA 0.6.1<sup>31</sup> and BEDTools 2.25.0 were used). RSEM<sup>32</sup> (version 1.2.15) was used to estimate gene levels based on ce10 ensembl annotations downloaded from UCSC genome table browser<sup>33</sup>. The gene expression count table was then imported into DESeq<sup>33</sup> (version 1.10.1) for differential gene expression test. See the README tab in Supplementary Table 2 for detailed descriptions of RNA-seq and analysis for Supplementary Table 2.

**Pathogen exposure and development assay.** Wild-type embryos were placed onto either NGM plates seeded with *Escherichia coli* OP50 or slow-killing assay plates seeded with *P. aeruginosa* PA14 and allowed to grow at 25 °C for 72 h. After 72 h embryos were collected from adults and placed on new plates seeded with either OP50 or PA14 and placed at 25 °C.

**M-cell division in response to starvation.** M-cell division analysis was performed as described previously<sup>10</sup>.

**Statistics and reproducibility.** ANOVA analysis with *post hoc* *P*-value calculations was used for Fig. 2, Fig. 3a,b, Fig. 4a and Supplementary Figs 3b and 4b. Unpaired two-tailed Student's *t*-test was used for Fig. 1, Fig. 3d,e, Fig. 4b,c and Supplementary Figs 1, 3c-f; \* = *P* < 0.05, \*\* = *P* < 0.01, \*\*\* = *P* < 0.001, \*\*\*\* *P* < 0.0001. No statistical method was used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.

**Metabolite preparation for quantification.** Approximately 100 μl of concentrated embryos were collected by egg preparation and placed on normal NGM agar plates for 3 h to recover. After 3 h, embryos were collected in M9, pelleted, and frozen. Frozen embryos were resuspended in 400 μl PBS and homogenized by douncing.

Homogenized embryos were centrifuged at 800g for 2 min to remove undouced tissue and 200 µl of supernatant was mixed with 800 µl of methanol and dried to extract polar metabolites. Dried samples were stored at  $-80^{\circ}\text{C}$ .

**Metabolite profiling.** Liquid chromatography and mass spectrometry were performed as described previously<sup>33</sup>.

**Assay for L2-, L3- and L4-stage developmental arrest.** Approximately 50 L2-, L3- or L4-stage animals were placed onto standard NGM Petri plates containing 500 mM NaCl for 24 h. After 24 h animals that were immobile and not developing or responding to touch were moved to plates containing 50 mM NaCl. Percentage developing is defined by the percentage of animals that resumed development and mobility after returning to normal growth conditions.

**MEK inhibitor exposure.** The MEK inhibitor U0126 (U120 Sigma-Aldrich) was resuspended in dimethylsulfoxide and added to standard NGM Petri plates at a final concentration of 100 *muupM*. These plates were then seeded with *Escherichia coli* OP50. Wild-type embryos were placed on plates containing either dimethylsulfoxide alone or dimethylsulfoxide and the MEK inhibitor U0126. Animals were allowed to grow for 72 h at  $25^{\circ}\text{C}$ . After 72 h, embryos from adult animals were collected and placed onto Petri plates containing 500 mM NaCl at  $25^{\circ}\text{C}$  for 24 h and the fraction of animals mobile and developing was determined.

**mek-2 RNAi exposure.** Wild-type embryos fed *Escherichia coli* HT115 containing either the empty vector L4440 or a *mek-2* RNAi vector (Ahringer library—Source Biosciences) were grown at  $20^{\circ}\text{C}$  for 72 h. After 72 h, embryos were collected and placed onto Petri plates containing 500 mM NaCl for 24 h at  $25^{\circ}\text{C}$  and the fraction of animals mobile and developing was determined.

**Assay for daf-2 adaptation to osmotic stress.** L4 animals were placed on standard NGM plates containing 300 mM NaCl at  $25^{\circ}\text{C}$  overnight. Embryos from resulting

adult animals were extracted with a razor blade into M9 solution and pipetted onto plates containing 500 mM NaCl. These animals were allowed to develop for 48 h at  $20^{\circ}\text{C}$ . Percentage of L2+ is defined by the percentage of animals that developed past the L1 larval stage.

**Data availability.** RNA-seq data that support the findings of this study have been deposited in the Gene Expression Omnibus (GEO) under accession codes GSE91073 and GSE91039. Data supporting the findings of Figs 1b,c,d, 2a,b,d, 3a,b,d,e and 4a,b,c and Supplementary Fig. 1b,c,d,e,f,g, Supplementary Fig. 3b,d,e,f and Supplementary Fig. 4b,c,d are provided in Supplementary Table 6. All other relevant data are available from the authors on request and/or are included with the manuscript.

24. Brenner, S. The genetics of *Caenorhabditis elegans*. *Genetics* **77**, 71–94 (1974).
25. Arur, S. *et al.* MPK-1 ERK controls membrane organization in *C. elegans* oogenesis via a sex-determination module. *Dev. Cell* **20**, 677–688 (2011).
26. Arur, S. *et al.* Multiple ERK substrates execute single biological processes in *Caenorhabditis elegans* germ-line development. *Proc. Natl Acad. Sci. USA* **106**, 4776–4781 (2009).
27. Lee, M.-H. *et al.* Multiple functions and dynamic activation of MPK-1 extracellular signal-regulated kinase signalling in *Caenorhabditis elegans* germline development. *Genetics* **177**, 2039–2062 (2007).
28. Quinlan, A. R. BEDTools: the Swiss-army tool for genome feature analysis. *Curr. Protoc. Bioinformatics* **47**, 11.12.1–11.12.34 (2014).
29. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).
30. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).
31. Karolchik, D. *et al.* The UCSC Table Browser data retrieval tool. *Nucleic Acids Res.* **32**, D493–D496 (2004).
32. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol.* **11**, R106 (2010).
33. Birsoy, K. *et al.* An essential role of the mitochondrial electron transport chain in cell proliferation is to enable aspartate synthesis. *Cell* **162**, 540–551 (2015).