## 18.S097 PSET 3

## IAP 2022

## Due 1/23/2022

Review / helpful information:

• Let  $A \subset \mathbb{R}^n$  and  $B \subset \mathbb{R}^n$ . Then, we define

$$A + B := \{a + b \mid a \in A, b \in B\}.$$

• We define the  $\ell^p$  norm on a sequence  $a = \{a_n\}_n$  of real numbers as

$$\|a\|_p = \left(\sum_{j=1}^{\infty} |a_n|^p\right)^{1/p}$$

where  $1 \le p < \infty$ . We define  $\ell^p$  space as the space of (infinite) sequences  $x = \{x_n\}_n$  in  $\mathbb{R}$  such that  $\|x\|_p < \infty$ .

**1.** Let A be a closed subset of  $\mathbb{R}^n$  and let B be a compact subset of  $\mathbb{R}^n$ . Show that A + B is closed. Hint: Let  $\{a_n\}_n$  be a sequence in A + B such that  $a_n \to z \in \mathbb{R}^n$ . Show that  $z \in A + B$ .

**Remark 1.** In fact, you can show that the sum of a closed set and a compact subset in a "topological vector space" is a closed set, but that goes beyond the scope of this class.

Let (X, d) be a metric space and S ⊂ X. Then, a point x ∈ S is an isolated point if there exists an ε > 0 such that B<sub>ε</sub>(x) contains no other points of S. Show that a point x ∈ S is an isolated point if and only if give a sequence {a<sub>n</sub>} in S converges to x, it must be the case that there exists an N such that for all n ≥ N, a<sub>n</sub> = x.
Let X be a metric space. Show that a finite union of compact subsets in X is compact.
Let 1 ≤ p < ∞. Consider the set</li>

$$S := \{a = \{a_n\} \in \ell^p \mid ||a||_p \le 1\}$$

Explain why S is closed and bounded in  $\ell^p$  (under the metric induced by the norm in PSET 2), and prove that S is not a compact subset of  $\ell^p$ .

Hint: Let  $e_n = {\delta_{k,n}}_k \in S$  where

$$\delta_{k,n} = \begin{cases} 1 & k = n \\ 0 & k \neq n \end{cases}.$$

Show that  $\{e_n\}_n$  does not have a convergent subsequence in S. Make sure to explain why this shows S is not compact.

5. Consider the set

$$S := \{ f \in C^0([0,1]) \mid ||f||_{\infty} = \sup_{x \in [0,1]} |f(x)| \le 1 \}.$$

Prove that S is not compact.

Hint: consider the sequence  $f_n(x) = x^n$ .

**6.** (Optional) Consider the space  $\ell^2$ . Show that the set

$$A = \{a = \{a_k\}_k \in \ell^2 \mid |a_k| < k^{-3}\}$$

is a compact subset of  $\ell^2.$ 

7. (Optional) Consider the set of functions of the form

$$\sum_{n} a_n e^{inx}$$

with  $|a_n| < (1+|n|)^{-2}$ .

- (a) Show that every function in this set lies in  $C^0([0, 2\pi])$ .
- (b) Show that this set is compact in  $C^0([0, 2\pi])$ .